

Q.1 [14 marks]

Fill in the blanks/MCQs using appropriate choice from the given options.

Q1.1 [1 mark]

$$\begin{vmatrix} 5 & 1 \\ 2 & 3 \end{vmatrix} = \underline{\hspace{1cm}}$$

Answer: b. 13

Solution:

For 2×2 determinant $\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$

$$\begin{vmatrix} 5 & 1 \\ 2 & 3 \end{vmatrix} = (5 \times 3) - (1 \times 2) = 15 - 2 = 13$$

Q1.2 [1 mark]

$$\text{If } \begin{vmatrix} x & 1 \\ 2 & 1 \end{vmatrix} = 0 \text{ then } x = \underline{\hspace{1cm}}$$

Answer: b. 2

Solution:

$$\begin{vmatrix} x & 1 \\ 2 & 1 \end{vmatrix} = x \times 1 - 1 \times 2 = x - 2 = 0$$

Therefore, $x = 2$

Q1.3 [1 mark]

$$\text{If } f(x) = x^2 \text{ then } f(-1) = \underline{\hspace{1cm}}$$

Answer: a. 1

Solution:

$$f(x) = x^2$$

$$f(-1) = (-1)^2 = 1$$

Q1.4 [1 mark]

$$\log_{10} 1 = \underline{\hspace{1cm}}$$

Answer: b. 0

Solution:

By logarithm property: $\log_a 1 = 0$ for any base $a > 0$

Therefore, $\log_{10} 1 = 0$

Q1.5 [1 mark]

$$\sin \frac{\pi}{2} + \cos \frac{\pi}{2} = \underline{\hspace{1cm}}$$

Answer: c. 1

Solution:

$$\sin \frac{\pi}{2} = 1 \text{ and } \cos \frac{\pi}{2} = 0$$

$$\text{Therefore, } \sin \frac{\pi}{2} + \cos \frac{\pi}{2} = 1 + 0 = 1$$

Q1.6 [1 mark]

$$\tan^{-1}(1) = \underline{\hspace{1cm}}$$

Answer: a. $\frac{\pi}{4}$

Solution:

$$\tan \frac{\pi}{4} = 1$$

$$\text{Therefore, } \tan^{-1}(1) = \frac{\pi}{4}$$

Q1.7 [1 mark]

$$\frac{2\pi}{3} \text{ radian} = \underline{\hspace{1cm}} \text{ degree}$$

Answer: d. 120

Solution:

$$\text{To convert radians to degrees: degrees} = \text{radians} \times \frac{180}{\pi}$$

$$\frac{2\pi}{3} \times \frac{180}{\pi} = \frac{2 \times 180}{3} = \frac{360}{3} = 120^\circ$$

Q1.8 [1 mark]

$$\hat{i} \times \hat{j} = \underline{\hspace{1cm}}$$

Answer: c. \hat{k}

Solution:

By right-hand rule for cross product:

$$\hat{i} \times \hat{j} = \hat{k}$$

Q1.9 [1 mark]

$$|\hat{i} + \hat{j} + \hat{k}| = \underline{\hspace{1cm}}$$

Answer: d. $\sqrt{3}$

Solution:

$$|\hat{i} + \hat{j} + \hat{k}| = \sqrt{1^2 + 1^2 + 1^2} = \sqrt{3}$$

Q1.10 [1 mark]

Slope of line $2x + y - 3 = 0$ is $\underline{\hspace{1cm}}$

Answer: a. -2

Solution:

Convert to slope-intercept form: $y = -2x + 3$

Slope = coefficient of $x = -2$

Q1.11 [1 mark]

Radius of circle $x^2 + y^2 = 81$ is ___

Answer: b. 9

Solution:

Standard form: $x^2 + y^2 = r^2$

Here, $r^2 = 81$, so $r = 9$

Q1.12 [1 mark]

$$\lim_{n \rightarrow \infty} \frac{1}{n} = \underline{\hspace{1cm}}$$

Answer: c. 0

Solution:

As n approaches infinity, $\frac{1}{n}$ approaches 0

Q1.13 [1 mark]

$$\lim_{x \rightarrow 1} (x^2 + x + 1) = \underline{\hspace{1cm}}$$

Answer: a. 3

Solution:

Direct substitution: $(1)^2 + (1) + 1 = 1 + 1 + 1 = 3$

Q1.14 [1 mark]

$$\lim_{\theta \rightarrow 0} \frac{\tan \theta}{\theta} = \underline{\hspace{1cm}}$$

Answer: b. 1

Solution:

This is a standard limit: $\lim_{\theta \rightarrow 0} \frac{\tan \theta}{\theta} = 1$

Q.2 (A) [6 marks]

Attempt any two

Q2.1 [3 marks]

Find the value of $\begin{vmatrix} 1 & 3 & 1 \\ 2 & -1 & 0 \\ 4 & -2 & 5 \end{vmatrix}$

Answer:

Solution:

Using expansion along second row (has zero):

$$= -2 \begin{vmatrix} 3 & 1 \\ -2 & 5 \end{vmatrix} + (-1) \begin{vmatrix} 1 & 1 \\ 4 & 5 \end{vmatrix} + 0$$

$$= -2(15 + 2) - 1(5 - 4)$$

$$= -2(17) - 1(1)$$

$$= -34 - 1 = -35$$

Table:

Step	Calculation	Result
Minor 1	$(3 \times 5) - (1 \times -2)$	17
Minor 2	$(1 \times 5) - (1 \times 4)$	1
Final	$-2(17) - 1(1)$	-35

Q2.2 [3 marks]If $f(x) = x^3 + 5$ then find $f(0)$, $f(1)$ and $f(-1)$ **Answer:****Solution:**

Given: $f(x) = x^3 + 5$

$$f(0) = (0)^3 + 5 = 0 + 5 = 5$$

$$f(1) = (1)^3 + 5 = 1 + 5 = 6$$

$$f(-1) = (-1)^3 + 5 = -1 + 5 = 4$$

Table:

Input	Calculation	Output
$f(0)$	$0^3 + 5$	5
$f(1)$	$1^3 + 5$	6
$f(-1)$	$(-1)^3 + 5$	4

Q2.3 [3 marks]Prove that $\tan^{-1} \left(\frac{1}{2} \right) + \tan^{-1} \left(\frac{1}{3} \right) = \frac{\pi}{4}$ **Answer:****Solution:**Using formula: $\tan^{-1} a + \tan^{-1} b = \tan^{-1} \left(\frac{a+b}{1-ab} \right)$

Let $a = \frac{1}{2}$, $b = \frac{1}{3}$

$$\begin{aligned}\tan^{-1}\left(\frac{1}{2}\right) + \tan^{-1}\left(\frac{1}{3}\right) &= \tan^{-1}\left(\frac{\frac{1}{2} + \frac{1}{3}}{1 - \frac{1}{2} \times \frac{1}{3}}\right) \\ &= \tan^{-1}\left(\frac{\frac{5}{6}}{1 - \frac{1}{6}}\right) = \tan^{-1}\left(\frac{\frac{5}{6}}{\frac{5}{6}}\right) = \tan^{-1}(1) = \frac{\pi}{4}\end{aligned}$$

Hence proved.

Q.2 (B) [8 marks]

Attempt any two

Q2.1 [4 marks]

If $f(x) = \frac{x-1}{x+1}$ then prove that $f(x) \cdot f(-x) = 1$

Answer:

Solution:

Given: $f(x) = \frac{x-1}{x+1}$

First find $f(-x)$:

$$f(-x) = \frac{(-x)-1}{(-x)+1} = \frac{-x-1}{-x+1} = \frac{-(x+1)}{-(x-1)} = \frac{x+1}{x-1}$$

Now calculate $f(x) \cdot f(-x)$:

$$f(x) \cdot f(-x) = \frac{x-1}{x+1} \cdot \frac{x+1}{x-1} = \frac{(x-1)(x+1)}{(x+1)(x-1)} = 1$$

Hence proved.

Q2.2 [4 marks]

If $\log\left(\frac{x+y}{2}\right) = \frac{1}{2}(\log x + \log y)$ then prove that $x = y$

Answer:

Solution:

Given: $\log\left(\frac{x+y}{2}\right) = \frac{1}{2}(\log x + \log y)$

Using logarithm properties:

$$\frac{1}{2}(\log x + \log y) = \frac{1}{2}\log(xy) = \log\sqrt{xy}$$

So: $\log\left(\frac{x+y}{2}\right) = \log\sqrt{xy}$

Taking antilog: $\frac{x+y}{2} = \sqrt{xy}$

Squaring both sides: $\left(\frac{x+y}{2}\right)^2 = xy$

$$\frac{(x+y)^2}{4} = xy$$

$$(x+y)^2 = 4xy$$

$$x^2 + 2xy + y^2 = 4xy$$

$$x^2 - 2xy + y^2 = 0$$

$$(x-y)^2 = 0$$

Therefore, $x = y$. Hence proved.

Q2.3 [4 marks]

Solve $\log(x + 3) + \log(x - 3) = \log 27$

Answer:

Solution:

Given: $\log(x + 3) + \log(x - 3) = \log 27$

Using logarithm property: $\log a + \log b = \log(ab)$

$$\log[(x + 3)(x - 3)] = \log 27$$

Taking antilog: $(x + 3)(x - 3) = 27$

$$x^2 - 9 = 27$$

$$x^2 = 36$$

$$x = \pm 6$$

Check validity:

- For $x = 6$: $x + 3 = 9 > 0$ and $x - 3 = 3 > 0 \checkmark$
- For $x = -6$: $x + 3 = -3 < 0$ (invalid for logarithm)

Therefore, $x = 6$

Q.3 (A) [6 marks]

Attempt any two

Q3.1 [3 marks]

Prove that $\frac{\sin(\frac{\pi}{2} + \theta)}{\cos(\pi - \theta)} + \frac{\tan(\pi - \theta)}{\cot(\frac{3\pi}{2} - \theta)} + \frac{\operatorname{cosec}(\frac{\pi}{2} - \theta)}{\sec(\pi + \theta)} = -3$

Answer:

Solution:

Using trigonometric identities:

$$\sin\left(\frac{\pi}{2} + \theta\right) = \cos \theta$$

$$\cos(\pi - \theta) = -\cos \theta$$

$$\tan(\pi - \theta) = -\tan \theta$$

$$\cot\left(\frac{3\pi}{2} - \theta\right) = \tan \theta$$

$$\operatorname{cosec}\left(\frac{\pi}{2} - \theta\right) = \sec \theta$$

$$\sec(\pi + \theta) = -\sec \theta$$

Substituting:

$$\frac{\cos \theta}{-\cos \theta} + \frac{-\tan \theta}{\tan \theta} + \frac{\sec \theta}{-\sec \theta}$$

$$= -1 + (-1) + (-1) = -3$$

Hence proved.

Q3.2 [3 marks]

Prove that $\tan 55^\circ = \frac{\cos 10^\circ + \sin 10^\circ}{\cos 10^\circ - \sin 10^\circ}$

Answer:

Solution:

We know that $\tan 55^\circ = \tan(45^\circ + 10^\circ)$

Using formula: $\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$

$$\tan 55^\circ = \frac{\tan 45^\circ + \tan 10^\circ}{1 - \tan 45^\circ \tan 10^\circ} = \frac{1 + \tan 10^\circ}{1 - \tan 10^\circ}$$

Now, $\tan 10^\circ = \frac{\sin 10^\circ}{\cos 10^\circ}$

$$\tan 55^\circ = \frac{1 + \frac{\sin 10^\circ}{\cos 10^\circ}}{1 - \frac{\sin 10^\circ}{\cos 10^\circ}} = \frac{\cos 10^\circ + \sin 10^\circ}{\cos 10^\circ - \sin 10^\circ}$$

Hence proved.

Q3.3 [3 marks]

If $\vec{a} = 2\hat{i} + 3\hat{j} + \hat{k}$, $\vec{b} = \hat{i} + \hat{j} + \hat{k}$ **and** $\vec{c} = 3\hat{i} + \hat{j} + \hat{k}$ **then find** $2\vec{a} + \vec{b} - \vec{c}$

Answer:

Solution:

Given:

$$\vec{a} = 2\hat{i} + 3\hat{j} + \hat{k}$$

$$\vec{b} = \hat{i} + \hat{j} + \hat{k}$$

$$\vec{c} = 3\hat{i} + \hat{j} + \hat{k}$$

$$2\vec{a} = 2(2\hat{i} + 3\hat{j} + \hat{k}) = 4\hat{i} + 6\hat{j} + 2\hat{k}$$

$$2\vec{a} + \vec{b} - \vec{c} = (4\hat{i} + 6\hat{j} + 2\hat{k}) + (\hat{i} + \hat{j} + \hat{k}) - (3\hat{i} + \hat{j} + \hat{k})$$

$$= (4 + 1 - 3)\hat{i} + (6 + 1 - 1)\hat{j} + (2 + 1 - 1)\hat{k}$$

$$= 2\hat{i} + 6\hat{j} + 2\hat{k}$$

Q.3 (B) [8 marks]

Attempt any two

Q3.1 [4 marks]

Prove that $\frac{\sin(x-y)}{\cos x \cos y} + \frac{\sin(y-z)}{\cos y \cos z} + \frac{\sin(z-x)}{\cos z \cos x} = 0$

Answer:

Solution:

Using identity: $\sin(A - B) = \sin A \cos B - \cos A \sin B$

$$\frac{\sin(x-y)}{\cos x \cos y} = \frac{\sin x \cos y - \cos x \sin y}{\cos x \cos y} = \tan x - \tan y$$

Similarly:

$$\frac{\sin(y-z)}{\cos y \cos z} = \tan y - \tan z$$

$$\frac{\sin(z-x)}{\cos z \cos x} = \tan z - \tan x$$

Adding all three:

$$(\tan x - \tan y) + (\tan y - \tan z) + (\tan z - \tan x) = 0$$

Hence proved.

Q3.2 [4 marks]

Draw graph of $y = \cos x$ for $0 \leq x \leq \pi$

Answer:

Solution:




Table of values:

x	0	$\pi/4$	$\pi/2$	$3\pi/4$	π
y	1	$\sqrt{2}/2$	0	$-\sqrt{2}/2$	-1

Q3.3 [4 marks]

Find equation of line passing through (1, 2) and (-3, 1)

Answer:

Solution:

Given points: $(x_1, y_1) = (1, 2)$ and $(x_2, y_2) = (-3, 1)$

$$\text{Slope: } m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - 2}{-3 - 1} = \frac{-1}{-4} = \frac{1}{4}$$

Using point-slope form: $y - y_1 = m(x - x_1)$

$$y - 2 = \frac{1}{4}(x - 1)$$

$$4(y - 2) = x - 1$$

$$4y - 8 = x - 1$$

$$x - 4y + 7 = 0$$

Equation: $x - 4y + 7 = 0$

Q.4 (A) [6 marks]

Attempt any two

Q4.1 [3 marks]

Find unit vector perpendicular to $\vec{a} = \hat{i} - 3\hat{j} + \hat{k}$ and $\vec{b} = 2\hat{i} + \hat{j} + 2\hat{k}$

Answer:

Solution:

$$\text{Cross product: } \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -3 & 1 \\ 2 & 1 & 2 \end{vmatrix}$$

$$= \hat{i}[(-3)(2) - (1)(1)] - \hat{j}[(1)(2) - (1)(2)] + \hat{k}[(1)(1) - (-3)(2)]$$

$$= \hat{i}(-6 - 1) - \hat{j}(2 - 2) + \hat{k}(1 + 6)$$

$$= -7\hat{i} + 0\hat{j} + 7\hat{k}$$

$$\text{Magnitude: } |\vec{a} \times \vec{b}| = \sqrt{(-7)^2 + 0^2 + 7^2} = \sqrt{49 + 49} = 7\sqrt{2}$$

$$\text{Unit vector: } \hat{n} = \frac{-7\hat{i} + 7\hat{k}}{7\sqrt{2}} = \frac{-\hat{i} + \hat{k}}{\sqrt{2}}$$

Q4.2 [3 marks]

Forces (1, 2, 1) and (2, -1, 3) act on a particle and the particle moves from point (2, 3, 1) to (4, 6, 2). Find the work done.

Answer:

Solution:

$$\text{Resultant force: } \vec{F} = (1, 2, 1) + (2, -1, 3) = (3, 1, 4)$$

$$\text{Displacement: } \vec{s} = (4, 6, 2) - (2, 3, 1) = (2, 3, 1)$$

$$\text{Work done: } W = \vec{F} \cdot \vec{s} = (3)(2) + (1)(3) + (4)(1) = 6 + 3 + 4 = 13 \text{ units}$$

Q4.3 [3 marks]

Show that lines $2x - 3y + 5 = 0$ and $8x - 12y - 3 = 0$ are parallel lines.

Answer:

Solution:

For line $2x - 3y + 5 = 0$: slope $m_1 = \frac{2}{3}$

For line $8x - 12y - 3 = 0$: slope $m_2 = \frac{8}{12} = \frac{2}{3}$

Since $m_1 = m_2 = \frac{2}{3}$, the lines are parallel.

Table:

Line	Standard Form	Slope
Line 1	$2x - 3y + 5 = 0$	$\frac{2}{3}$
Line 2	$8x - 12y - 3 = 0$	$\frac{2}{3}$

Q.4 (B) [8 marks]

Attempt any two

Q4.1 [4 marks]

Show that angle between $\vec{a} = \hat{i} + \hat{j} - \hat{k}$ and $\vec{b} = 2\hat{i} - 2\hat{j} + \hat{k}$ is $\sin^{-1} \left(\frac{\sqrt{26}}{27} \right)$

Answer:

Solution:

$$\vec{a} \cdot \vec{b} = (1)(2) + (1)(-2) + (-1)(1) = 2 - 2 - 1 = -1$$

$$|\vec{a}| = \sqrt{1^2 + 1^2 + (-1)^2} = \sqrt{3}$$

$$|\vec{b}| = \sqrt{2^2 + (-2)^2 + 1^2} = \sqrt{9} = 3$$

$$\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} = \frac{-1}{\sqrt{3} \times 3} = \frac{-1}{3\sqrt{3}}$$

$$\sin^2 \theta = 1 - \cos^2 \theta = 1 - \frac{1}{27} = \frac{26}{27}$$

$$\text{Therefore, } \sin \theta = \frac{\sqrt{26}}{3\sqrt{3}} = \frac{\sqrt{26}}{\sqrt{27}}$$

$$\text{Hence, } \theta = \sin^{-1} \left(\frac{\sqrt{26}}{\sqrt{27}} \right)$$

Q4.2 [4 marks]

If $\vec{a} = (1, 1, 1)$, $\vec{b} = (2, 0, 1)$ and $\vec{c} = (-2, 1, 0)$ then find $\vec{a} \cdot (\vec{b} \times \vec{c})$

Answer:

Solution:

First find $\vec{b} \times \vec{c}$:

$$\vec{b} \times \vec{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 0 & 1 \\ -2 & 1 & 0 \end{vmatrix}$$

$$\begin{aligned}
&= \hat{i}(0 \times 0 - 1 \times 1) - \hat{j}(2 \times 0 - 1 \times (-2)) + \hat{k}(2 \times 1 - 0 \times (-2)) \\
&= \hat{i}(-1) - \hat{j}(2) + \hat{k}(2) \\
&= -\hat{i} - 2\hat{j} + 2\hat{k}
\end{aligned}$$

Now find $\vec{a} \cdot (\vec{b} \times \vec{c})$:

$$\begin{aligned}
\vec{a} \cdot (\vec{b} \times \vec{c}) &= (1, 1, 1) \cdot (-1, -2, 2) \\
&= (1)(-1) + (1)(-2) + (1)(2) = -1 - 2 + 2 = -1
\end{aligned}$$

Q4.3 [4 marks]

Evaluate $\lim_{\theta \rightarrow 0} \frac{\sin 4\theta}{\theta}$

Answer:

Solution:

$$\lim_{\theta \rightarrow 0} \frac{\sin 4\theta}{\theta} = \lim_{\theta \rightarrow 0} \frac{\sin 4\theta}{4\theta} \times 4$$

Using standard limit $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$:

Let $u = 4\theta$, then as $\theta \rightarrow 0$, $u \rightarrow 0$

$$\lim_{\theta \rightarrow 0} \frac{\sin 4\theta}{4\theta} = \lim_{u \rightarrow 0} \frac{\sin u}{u} = 1$$

$$\text{Therefore, } \lim_{\theta \rightarrow 0} \frac{\sin 4\theta}{\theta} = 4 \times 1 = 4$$

Q.5 (A) [6 marks]

Attempt any two

Q5.1 [3 marks]

Evaluate $\lim_{x \rightarrow 9} \frac{x^2 - 81}{x - 9}$

Answer:

Solution:

Direct substitution gives $\frac{0}{0}$ form.

Factor the numerator: $x^2 - 81 = (x - 9)(x + 9)$

$$\begin{aligned}
\lim_{x \rightarrow 9} \frac{x^2 - 81}{x - 9} &= \lim_{x \rightarrow 9} \frac{(x - 9)(x + 9)}{x - 9} \\
&= \lim_{x \rightarrow 9} (x + 9) = 9 + 9 = 18
\end{aligned}$$

Q5.2 [3 marks]

Evaluate $\lim_{x \rightarrow \infty} \left(1 + \frac{3}{x}\right)^{2x}$

Answer:

Solution:

$$\text{Let } y = \left(1 + \frac{3}{x}\right)^{2x}$$

Taking natural logarithm:

$$\ln y = 2x \ln \left(1 + \frac{3}{x}\right)$$

$$\text{As } x \rightarrow \infty, \frac{3}{x} \rightarrow 0$$

Using $\ln(1+u) \approx u$ for small u :

$$\ln y = 2x \times \frac{3}{x} = 6$$

Therefore, $y = e^6$

Q5.3 [3 marks]

Evaluate $\lim_{x \rightarrow 1} \frac{x-1}{x^2+x-2}$

Answer:

Solution:

Factor the denominator: $x^2 + x - 2 = (x+2)(x-1)$

$$\lim_{x \rightarrow 1} \frac{x-1}{x^2+x-2} = \lim_{x \rightarrow 1} \frac{x-1}{(x+2)(x-1)}$$

$$= \lim_{x \rightarrow 1} \frac{1}{x+2} = \frac{1}{1+2} = \frac{1}{3}$$

Q.5 (B) [8 marks]

Attempt any two

Q5.1 [4 marks]

Find the equation of line passing through the point (2, -3) and having slope 4.

Answer:

Solution:

Using point-slope form: $y - y_1 = m(x - x_1)$

Given: $(x_1, y_1) = (2, -3)$ and slope $m = 4$

$$y - (-3) = 4(x - 2)$$

$$y + 3 = 4x - 8$$

$$y = 4x - 11$$

Equation: $y = 4x - 11$ or $4x - y - 11 = 0$

Q5.2 [4 marks]

For what value of m , lines $7x + y - 1 = 0$ and $3x - my + 2 = 0$ are perpendicular to each other.

Answer:

Solution:

For perpendicular lines, product of slopes = -1

For line $7x + y - 1 = 0$: slope $m_1 = -7$

For line $3x - my + 2 = 0$: slope $m_2 = \frac{3}{m}$

Condition: $m_1 \times m_2 = -1$

$$(-7) \times \frac{3}{m} = -1$$

$$\frac{-21}{m} = -1$$

$$21 = m$$

Therefore, $m = 21$

Table:

Line	Standard Form	Slope
Line 1	$7x + y - 1 = 0$	-7
Line 2	$3x - my + 2 = 0$	$\frac{3}{m}$

Verification: When $m = 21$, slopes are -7 and $\frac{3}{21} = \frac{1}{7}$

Product: $(-7) \times \frac{1}{7} = -1 \checkmark$

Q5.3 [4 marks]

Find the centre and radius of the circle $4x^2 + 4y^2 + 8x - 12y - 3 = 0$

Answer:

Solution:

First, divide by 4 to get standard form:

$$x^2 + y^2 + 2x - 3y - \frac{3}{4} = 0$$

Complete the square for x and y terms:

$$x^2 + 2x = (x + 1)^2 - 1$$

$$y^2 - 3y = \left(y - \frac{3}{2}\right)^2 - \frac{9}{4}$$

Substituting:

$$(x + 1)^2 - 1 + \left(y - \frac{3}{2}\right)^2 - \frac{9}{4} - \frac{3}{4} = 0$$

$$(x + 1)^2 + \left(y - \frac{3}{2}\right)^2 = 1 + \frac{9}{4} + \frac{3}{4} = 1 + 3 = 4$$

Centre: $\left(-1, \frac{3}{2}\right)$

Radius: $r = \sqrt{4} = 2$

Table:

Component	Value
Centre (h,k)	$\left(-1, \frac{3}{2}\right)$
Radius	2
Standard Form	$(x + 1)^2 + \left(y - \frac{3}{2}\right)^2 = 4$

Formula Cheat Sheet

Determinants

- **2×2 Determinant:** $\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$
- **3×3 Determinant:** Expand along any row/column

Functions & Logarithms

- **Basic:** $\log_a 1 = 0, \log_a a = 1$
- **Properties:** $\log(ab) = \log a + \log b, \log\left(\frac{a}{b}\right) = \log a - \log b$

Trigonometry

- **Basic Values:** $\sin 0^\circ = 0, \sin 30^\circ = \frac{1}{2}, \sin 45^\circ = \frac{\sqrt{2}}{2}, \sin 60^\circ = \frac{\sqrt{3}}{2}, \sin 90^\circ = 1$
- **Conversion:** Radians to degrees: $\times \frac{180}{\pi}$
- **Identities:** $\sin^2 \theta + \cos^2 \theta = 1$
- **Inverse:** $\tan^{-1}(1) = \frac{\pi}{4}$

Vectors

- **Magnitude:** $|\vec{a}| = \sqrt{a_x^2 + a_y^2 + a_z^2}$
- **Dot Product:** $\vec{a} \cdot \vec{b} = a_x b_x + a_y b_y + a_z b_z$
- **Cross Product:** $\hat{i} \times \hat{j} = \hat{k}, \hat{j} \times \hat{k} = \hat{i}, \hat{k} \times \hat{i} = \hat{j}$
- **Work Done:** $W = \vec{F} \cdot \vec{s}$

Coordinate Geometry

- **Slope:** $m = \frac{y_2 - y_1}{x_2 - x_1}$
- **Point-Slope Form:** $y - y_1 = m(x - x_1)$
- **Parallel Lines:** Same slope
- **Perpendicular Lines:** Product of slopes = -1
- **Circle:** $(x - h)^2 + (y - k)^2 = r^2$

Limits

- **Standard Limits:** $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1, \lim_{x \rightarrow 0} \frac{\tan x}{x} = 1$
- **Factorization:** Use for $\frac{0}{0}$ forms
- **L'Hôpital's Rule:** For indeterminate forms

Problem-Solving Strategies

For Determinants:

1. Choose the row/column with most zeros for expansion
2. Use cofactor expansion systematically
3. Check calculations by expanding along different rows

For Functions:

1. Direct substitution first
2. Use function properties and definitions
3. Check domain restrictions

For Trigonometry:

1. Convert all angles to same unit (degrees or radians)
2. Use standard angle values
3. Apply appropriate identities
4. Simplify step by step

For Vectors:

1. Write components clearly
2. Use right-hand rule for cross products
3. Check units and directions
4. Verify with geometric interpretation

For Coordinate Geometry:



1. Plot points when possible
2. Use appropriate formulas based on given information
3. Check parallel/perpendicular conditions
4. Complete the square for circles

For Limits:



1. Try direct substitution first
2. Factor polynomials for $\frac{0}{0}$ forms
3. Use standard limit formulas
4. Apply L'Hôpital's rule for indeterminate forms

Common Mistakes to Avoid



Determinants:

-  Wrong sign in calculations
-  Follow cofactor signs carefully: $(-1)^{i+j}$



Logarithms:

-  $\log(a + b) = \log a + \log b$ (WRONG)
-  $\log(ab) = \log a + \log b$ (CORRECT)



Trigonometry:

-  Mixing degrees and radians
-  Convert to same unit first

Vectors:

-  $\vec{a} \times \vec{b} = \vec{b} \times \vec{a}$ (WRONG)
-  $\vec{a} \times \vec{b} = -(\vec{b} \times \vec{a})$ (CORRECT)

Slopes:

-  Confusing parallel and perpendicular conditions
-  Parallel: same slope, Perpendicular: product = -1

Limits:

-  Direct substitution without checking indeterminate forms
-  Check for $\frac{0}{0}$ or $\frac{\infty}{\infty}$ first

Exam Tips

Time Management:

- Spend 2 minutes per mark (14 marks = 28 minutes for Q1)
- Start with familiar questions
- Leave difficult problems for the end

Calculation Tips:

- Show all steps clearly
- Use tables for organized presentation
- Double-check arithmetic

- Write final answers clearly

Writing Strategy:

- Write given information first
- State formulas before using them
- Include units where applicable
- Box or underline final answers

Last-Minute Checks:

- Verify all calculations
- Check if answers are reasonable
- Ensure all parts are attempted
- Review question requirements

Mnemonic for Standard Angles:

"Some People Have Curly Brown Hair Through Proper Brushing"

- $\sin 0^\circ = 0$, $\pi/6 = 1/2$, $\text{Half} = \sqrt{2}/2$, Cos complement , etc.

Remember: Mathematics is about **understanding patterns**, not memorizing formulas. Practice regularly and **think step by step!**

Quick Reference Table

Topic	Key Formula	Example
Determinant 2×2	$ad - bc$	$\begin{vmatrix} 2 & 3 \\ 1 & 4 \end{vmatrix} = 8 - 3 = 5$
Slope	$\frac{y_2 - y_1}{x_2 - x_1}$	Points (1,2), (3,8): $m = \frac{8-2}{3-1} = 3$
Distance	$\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$	Between (0,0), (3,4): $d = 5$
Circle	$(x - h)^2 + (y - k)^2 = r^2$	Center (1,2), radius 3
Limit	$\lim_{x \rightarrow a} f(x)$	Direct substitution or factoring

Final Tip: Keep practicing and stay confident! 🎯