EQUISTASI

PATIENT GUIDE · PAIN

Chronic pain: how to use Equistasi®

Equistasi® is a medical device designed to help you manage chronic pain through focal vibrations that stimulate body awareness, helping to reduce pain. Here is how to use it effectively.

1 Where to apply the devices

1 On the painful spot


Apply the first device where you feel the most intense pain.

2·3 On the muscles involved

Apply the other two on the muscles that support the movement of the painful area: they help body awareness and reduce tension. In the app you will find the muscles for every area.

4 On the vertebra of the metamere

Apply one device on the vertebra of the **metamere** linked to the muscle area: press the vertebrae of that area and choose the **most tender** one (knee → L3-L4). You will find them in the app, **Metamere and dermatome** section.



Example: knee pain. 1 on the painful spot, 2 and 3 on the thigh muscles, 4 on the L3-L4 vertebra (back view).

How to attach it: clean and dry the skin well (no creams or oils) and fix each device with the transparent patch supplied, in contact with the skin.

2 Check that it works

After applying the devices, wait about **10 minutes**.

✓ Do you feel even slight relief?
It means the chosen points are correct.

↻ No relief after 10–20 minutes?
Move the devices to nearby points, along the muscles or on the spine.

For greater benefit: adjust the points until you find those that give you the most relief; combining the pain point with nearby points can give longer-lasting relief.

This information does not replace the advice of a doctor or physiotherapist. Always read the device instructions for use.

Equistasi S.r.l. · www.equistasi.com · info@equistasi.com
Rev. 1 – October 2026 · page 1 of 2

3 How long to keep it on

First application	Leave the devices on the painful spot and on the muscles in place **until the next day**; remove them about **an hour after waking up**.
Device on the vertebra	Start with **1 hour**; if needed, increase up to **2 hours**.
If the pain has gone	There is no need to continue: when the pain comes back, reapply the devices in the same way.
Weekly break	To keep it effective over time, take a break of **at least 2 days every week**. **M** **T** **W** **T** **F** S S

⚠ When to remove the devices

- Remove them **earlier** if you feel **discomfort of any kind**: it passes within a few minutes after removing them.
- During application the initial pain may decrease or disappear and then return as **mild discomfort**: this is the time to remove the devices. The discomfort quickly decreases after removal.
- If you feel **nausea, headache or dizziness**, remove the devices immediately and try again the next day.

4 In short



1. One device on the **most painful spot**, two on the **muscles** supporting the area and one on the **vertebra** of the metamere (1–2 hours).
2. After **10 minutes** check: if you feel no relief, move the devices to nearby points.
3. The first time keep them on **until the next morning**; then reapply only when the pain returns, with **2 days off** a week.

This information does not replace the advice of a doctor or physiotherapist. Always read the device instructions for use.