EQUISTASI

PATIENT GUIDE · PARKINSON

Parkinson's: balance and postural stability

Equistasi® generates focal mechanical stimulation intended to influence proprioceptive information; the response is individual. Follow the advice of your neurologist or physiotherapist.

1 Where to apply the 3 devices

1 Seventh cervical vertebra (C7)
At the back of the neck: the most prominent vertebra when you bend your head forward.

2 Right soleus
At the back of the leg, at the base of the calf.

3 Left soleus
In the same position on the other leg.

How to attach it: clean and dry the skin well (no creams or oils) and fix each device with the transparent patch supplied, in contact with the skin.



Back view. 1 on C7, 2 and 3 on both soleus muscles.

2 How long to keep it on

First day	**1 hour** in a row, preferably at the times of day when you move the most.
From the second day	**2 hours in the morning** and **2 hours in the afternoon**.
Every week	**5 days** of use and **2 days off**, so that the receptors do not get used to the stimulus. M T W T F S S

This information does not replace the advice of a doctor or physiotherapist. Always read the device instructions for use.

Equistasi S.r.l. · www.equistasi.com · info@equistasi.com

3 What studies show

- Significant reduction in the risk of falls (from 3.2 to 0.65 falls per month in multicentre studies).
- Better postural stability and proprioception thanks to continuous stimulation.
- Better walking: longer steps and fewer freezing episodes.

⚠ Warnings

- Remove them **earlier** if you feel **discomfort of any kind**: it passes within a few minutes after removing them.
- If **nausea, headache or dizziness** appear, remove the devices immediately and reduce the application times.
- The devices are **strictly personal**: they adapt to your body temperature and must not be shared.

Find the points with the app. Scan the QR code: in the Equistasi Point app you will find anatomical plates with the application points and video tutorials.

This information does not replace the advice of a doctor or physiotherapist. Always read the device instructions for use.