EQUISTASI


PATIENT GUIDE · MULTIPLE SCLEROSIS

Multiple sclerosis: how to use Equistasi®

Two application patterns based on the study by the University of Naples Federico II.
Your doctor or physiotherapist decides which one to use.

A Balance and step length


1. **C7**
   The most prominent vertebra at the base of the neck.
2. **L5**
   The last lumbar vertebra, low in the back.
3. **Calves**
   One device on each calf (triceps surae).



Back view. 1 C7, 2 L5, 3 both calves (4 devices).

B Spasticity in one leg

1. **C7**
   At the base of the neck.
2. **Gluteus medius**
   On the side of the hip, in the upper buttock, on the side of the affected leg.
3. **Quadriceps**
   In the lower part of the thigh of the affected leg, above the knee.
4. **Tibialis anterior or calf**
   In the lower part of the affected leg, at the front (tibialis anterior) or at the back (calf): your physiotherapist chooses based on ankle movement.



Back and front views (example: right leg).

This information does not replace the advice of a doctor or physiotherapist. Always read the device instructions for use.

Equistasi S.r.l. · www.equistasi.com · info@equistasi.com
Rev. 1 – October 2026 · page 1 of 2

1 How long to keep it on

Week 1	**1.5 hours a day** for 5 days, then 2 days off.
Weeks 2, 3 and 4	**2 hours a day** for 5 days, then 2 days off.
If you feel dizzy	Start with **15 minutes** and try again later; increase the time gradually over the following days. **M** **T** **W** **T** **F** S S

2 How to assess progress

Your physiotherapist can measure balance and walking with simple tests (such as the Timed Up and Go and the Romberg test), ideally recorded on video: before starting, after the first hour of application and at later check-ups.

3 If you notice no perceived benefit

- Talk to your doctor: some medicines may affect the response to the device.
- You can take a break of one or more weeks and then repeat the treatment.
- Your physiotherapist can consider different application points.

How to attach it: clean and dry the skin well (no creams or oils) and fix each device with the transparent patch supplied, in contact with the skin.

⚠ When to remove the devices

- Remove them **earlier** if you feel **discomfort of any kind**: it passes within a few minutes after removing them.
- If you feel **nausea, headache or dizziness**, remove the devices immediately and try again later for a shorter time.

Find the points with the app. Scan the QR code: in the Equistasi Point app you will find anatomical plates with the application points and video tutorials.

This information does not replace the advice of a doctor or physiotherapist. Always read the device instructions for use.