EQUISTASI

PATIENT GUIDE · PAIN

# Chronic pain: how to use Equistasi®

Equistasi® generates focal mechanical stimulation intended to influence proprioceptive information. In some people it may be associated with a change in pain perception: the response is individual. Here is how to use it.

## 1 Where to apply the devices

### 1 On the painful spot


Apply the first device where you feel the most intense pain.

### 2·3 On the muscles involved

Apply the other two on the muscles that support the movement of the painful area. In the app you will find the muscles for every area.

### 4 On the vertebra of the metamere

Apply one device on the vertebra of the **metamere** linked to the muscle area: press the vertebrae of that area and choose the **most tender** one (knee → L3-L4). You will find them in the app, **Metamere and dermatome** section.




**Example: knee pain.** 1 on the painful spot, 2 and 3 on the thigh muscles, 4 on the L3-L4 vertebra (back view).

**How to attach it:** clean and dry the skin well (no creams or oils) and fix each device with the transparent patch supplied, in contact with the skin.

## 2 Check that it works

After applying the devices, wait about **10 minutes**.

**✓ Do you notice any perceived benefit?**

The chosen points suit you: the response is individual.

**↻ No perceived benefit after 10–20 minutes?**

Move the devices to nearby points, along the muscles or on the spine.

**Individual response:** you can adjust the points until you find those with which you notice any perceived benefit, also combining the pain point with nearby points.

This information does not replace the advice of a doctor or physiotherapist. Always read the device instructions for use.

**Equistasi S.r.l.** · www.equistasi.com · info@equistasi.com
Rev. 1 – October 2026 · page 1 of 2

## 3 How long to keep it on

| | |
|---|---|
| **First application** | Leave the devices on the painful spot and on the muscles in place **until the next day**; remove them about **an hour after waking up**. |
| **Device on the vertebra** | Start with **1 hour**; if needed, increase up to **2 hours**. |
| **If the pain has gone** | There is no need to continue: when the pain comes back, reapply the devices in the same way. |
| **Weekly break** | Take a break of **at least 2 days every week**.<br>**M** **T** **W** **T** **F** S S |

### ⚠ When to remove the devices

- Remove them **earlier** if you feel **discomfort of any kind**: it passes within a few minutes after removing them.
- During application your perception of pain may change; if **mild discomfort** then appears, this is the time to remove the devices. The discomfort quickly decreases after removal.
- If you feel **nausea, headache or dizziness**, remove the devices immediately and try again the next day.

## 4 In short



1. One device on the **most painful spot**, two on the **muscles** supporting the area and one on the **vertebra** of the metamere (1–2 hours).
2. After **10 minutes** check: if you notice no perceived benefit, move the devices to nearby points.
3. The first time keep them on **until the next morning**; then reapply only when the pain returns, with **2 days off** a week.

This information does not replace the advice of a doctor or physiotherapist. Always read the device instructions for use.