EQUISTASI

PATIENT GUIDE · PAIN

Pain and dermatomes: how to use Equistasi®

Each area of skin is connected to a nerve root that leaves the spine: this area is called a **dermatome**. Knowing it helps you personalise how you apply Equistasi®, going beyond the standard points.

1 Find the painful area

If you feel pain or changes in sensation (tingling, reduced feeling) in a specific area, look up the matching dermatome on the **map on page 3**. In the Equistasi Point app, in the **Metamere and dermatome** section, you will find the nerve root and the linked vertebrae for every area.


2 Where to apply the 3 devices

1 On the main point
Apply one device on the area of greatest tenderness or pain, within the affected dermatome.

2·3 On the supporting muscles
Apply the other two on the muscles that support the area (for example the quadriceps for L3-L4, the calves for S1-S2).

+ Or on the vertebra
You can also apply a device on the vertebra matching the dermatome (for example pain in L5 → lower back).

How to attach it: clean and dry the skin well (no creams or oils) and fix each device with the transparent patch supplied, in contact with the skin.



Example: L4-L5. 1 near the lumbar spine, 2 and 3 on the calves.

This information does not replace the advice of a doctor or physiotherapist. Always read the device instructions for use.

Equistasi S.r.l. · www.equistasi.com · info@equistasi.com
Rev. 1 – October 2026 · page 1 of 3

3 Practical examples

L4-L5 Pain or instability in the legs

One device near the matching lumbar spine and two on the calves (triceps surae).

C5-C7 Neck pain spreading to the arm

One device on C7 (the most prominent vertebra at the base of the neck) and two on the arms, in the areas of pain or muscle stiffness.

S1-S2 Walking problems in Parkinson's

Classic protocol: C7 and both calves (triceps surae), which correspond to dermatomes S1-S2.

4 How long to keep it on


After applying	Wait about **60 seconds** and reassess the pain.
On the vertebra	The first times keep the devices on for **1 hour**; if needed, then increase up to **2 hours**.
Every week	**5 days** of use and **2 days off**. **M** **T** **W** **T** **F** S S

⚠ When to remove the devices

- Remove them **earlier** if you feel **discomfort of any kind**: it passes within a few minutes after removing them.
- If you feel **nausea, headache or dizziness**, remove the devices immediately and try again the next day.

This information does not replace the advice of a doctor or physiotherapist. Always read the device instructions for use.

5 Dermatome map



Front and back. The map shows the dermatomes as separate areas: in reality each dermatome partly overlaps its neighbours.
Source: Frank H. Netter, *Atlas of Human Anatomy*, Elsevier.

Main dermatome levels

C5	Collarbones
C5, C6, C7	Outer side of the arms
C8, T1	Inner side of the arms
C6	Thumb
C6, C7, C8	Hand
C8	Ring and little fingers
T4	Nipple level
T10	Navel level
T12	Groin
L1, L2, L3, L4	Front and inner side of the legs
L4, L5, S1	Feet
L4	Inner side of the big toe
L5, S1, S2	Outer side and back of the legs
S1	Outer edge of the foot and little toe
S2, S3, S4, S5	Perineum

This information does not replace the advice of a doctor or physiotherapist. Always read the device instructions for use.