

MA579H Scientific Computing

Numerical Integration III

Rafikul Alam
Department of Mathematics
IIT Guwahati

Lecture outline

- Computation of nodes and weights for Gaussian quadrature

Construction of orthogonal polynomials

Let $\langle p, q \rangle$ be an inner product on $\mathcal{P}_n = \text{span}(1, x, \dots, x^n)$. For example,

Legendre: $\langle p, q \rangle := \int_{-1}^1 p(x)q(x)dx$

Construction of orthogonal polynomials

Let $\langle p, q \rangle$ be an inner product on $\mathcal{P}_n = \text{span}(1, x, \dots, x^n)$. For example,

Legendre: $\langle p, q \rangle := \int_{-1}^1 p(x)q(x)dx \longrightarrow \int_{-1}^1 f(x)dx$

Construction of orthogonal polynomials

Let $\langle p, q \rangle$ be an inner product on $\mathcal{P}_n = \text{span}(1, x, \dots, x^n)$. For example,

Legendre: $\langle p, q \rangle := \int_{-1}^1 p(x)q(x)dx \longrightarrow \int_{-1}^1 f(x)dx$

Laguerre: $\langle p, q \rangle := \int_0^\infty p(x)q(x)e^{-x}dx$

Construction of orthogonal polynomials

Let $\langle p, q \rangle$ be an inner product on $\mathcal{P}_n = \text{span}(1, x, \dots, x^n)$. For example,

Legendre: $\langle p, q \rangle := \int_{-1}^1 p(x)q(x)dx \longrightarrow \int_{-1}^1 f(x)dx$

Laguerre: $\langle p, q \rangle := \int_0^\infty p(x)q(x)e^{-x}dx \longrightarrow \int_0^\infty f(x)e^{-x}dx$

Construction of orthogonal polynomials

Let $\langle p, q \rangle$ be an inner product on $\mathcal{P}_n = \text{span}(1, x, \dots, x^n)$. For example,

Legendre: $\langle p, q \rangle := \int_{-1}^1 p(x)q(x)dx \longrightarrow \int_{-1}^1 f(x)dx$

Laguerre: $\langle p, q \rangle := \int_0^\infty p(x)q(x)e^{-x}dx \longrightarrow \int_0^\infty f(x)e^{-x}dx$

Chebyshev: $\langle p, q \rangle := \int_{-1}^1 p(x)q(x)(1-x^2)^{-1/2}dx$

Construction of orthogonal polynomials

Let $\langle p, q \rangle$ be an inner product on $\mathcal{P}_n = \text{span}(1, x, \dots, x^n)$. For example,

Legendre: $\langle p, q \rangle := \int_{-1}^1 p(x)q(x)dx \longrightarrow \int_{-1}^1 f(x)dx$

Laguerre: $\langle p, q \rangle := \int_0^\infty p(x)q(x)e^{-x}dx \longrightarrow \int_0^\infty f(x)e^{-x}dx$

Chebyshev: $\langle p, q \rangle := \int_{-1}^1 p(x)q(x)(1-x^2)^{-1/2}dx$
 $\longrightarrow \int_{-1}^1 f(x)(1-x^2)^{-1/2}dx$

are inner products on \mathcal{P}_n .

Construction of orthogonal polynomials

Let $\langle p, q \rangle$ be an inner product on $\mathcal{P}_n = \text{span}(1, x, \dots, x^n)$. For example,

Legendre: $\langle p, q \rangle := \int_{-1}^1 p(x)q(x)dx \longrightarrow \int_{-1}^1 f(x)dx$

Laguerre: $\langle p, q \rangle := \int_0^\infty p(x)q(x)e^{-x}dx \longrightarrow \int_0^\infty f(x)e^{-x}dx$

Chebyshev: $\langle p, q \rangle := \int_{-1}^1 p(x)q(x)(1-x^2)^{-1/2}dx$
 $\longrightarrow \int_{-1}^1 f(x)(1-x^2)^{-1/2}dx$

are inner products on \mathcal{P}_n .

Given an inner product on \mathcal{P}_n , there exist **orthogonal polynomials** $\phi_0(x), \phi_1(x), \dots, \phi_n(x)$ such that $\deg(\phi_\ell(x)) = \ell$ for $\ell = 0 : n$ and

$$\text{span}(1, x, \dots, x^j) = \text{span}(\phi_0(x), \phi_1(x), \dots, \phi_j(x)), \quad j = 0 : n.$$

Gauss quadrature

Gauss Legendre quadrature: Evaluate $\int_{-1}^1 f(x)dx$.

Gauss quadrature

Gauss Legendre quadrature: Evaluate $\int_{-1}^1 f(x)dx$.

Legendre polynomial (Rodrigues' formula): $P_n(x) := \frac{1}{2^n n!} \frac{d^n}{dx^n} (x^2 - 1)^n$.

Recurrence relation (monic Legendre):

$$P_{n+1}(x) = xP_n(x) - \frac{n^2}{4n^2 - 1} P_{n-1}(x).$$

Gauss quadrature

Gauss Legendre quadrature: Evaluate $\int_{-1}^1 f(x)dx$.

Legendre polynomial (Rodrigues' formula): $P_n(x) := \frac{1}{2^n n!} \frac{d^n}{dx^n} (x^2 - 1)^n$.

Recurrence relation (monic Legendre):

$$P_{n+1}(x) = xP_n(x) - \frac{n^2}{4n^2 - 1} P_{n-1}(x).$$

Gauss Laguerre quadrature: Evaluate $\int_0^\infty f(x)e^{-x}dx$.

Gauss quadrature

Gauss Legendre quadrature: Evaluate $\int_{-1}^1 f(x)dx$.

Legendre polynomial (Rodrigues' formula): $P_n(x) := \frac{1}{2^n n!} \frac{d^n}{dx^n} (x^2 - 1)^n$.

Recurrence relation (monic Legendre):

$$P_{n+1}(x) = xP_n(x) - \frac{n^2}{4n^2 - 1} P_{n-1}(x).$$

Gauss Laguerre quadrature: Evaluate $\int_0^\infty f(x)e^{-x}dx$.

Laguerre polynomial (Rodrigues formula) :

$$L_n(x) := \frac{e^x}{n!} \frac{d^n}{dx^n} (e^{-x} x^n) = \frac{1}{n!} \left(\frac{d}{dx} - 1 \right)^n x^n.$$

Recurrence relation (monic Laguerre):

$$L_{n+1}(x) = (x - (2n + 1))L_n(x) - n^2 L_{n-1}(x).$$

Gauss quadrature

Gauss Chebyshev quadrature: Evaluate $\int_{-1}^1 f(x)(1-x^2)^{-1/2} dx$.

Gauss quadrature

Gauss Chebyshev quadrature: Evaluate $\int_{-1}^1 f(x)(1-x^2)^{-1/2} dx$.

Chebyshev polynomial: For $x \in [-1, 1]$,

$$T_n(x) := \cos(n \cos^{-1} x)$$

is a Chebyshev polynomial of degree n .

Gauss quadrature

Gauss Chebyshev quadrature: Evaluate $\int_{-1}^1 f(x)(1-x^2)^{-1/2} dx$.

Chebyshev polynomial: For $x \in [-1, 1]$,

$$T_n(x) := \cos(n \cos^{-1} x)$$

is a Chebyshev polynomial of degree n .

Recurrence relation

$$T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x),$$

Gauss quadrature

Gauss Chebyshev quadrature: Evaluate $\int_{-1}^1 f(x)(1-x^2)^{-1/2} dx$.

Chebyshev polynomial: For $x \in [-1, 1]$,

$$T_n(x) := \cos(n \cos^{-1} x)$$

is a Chebyshev polynomial of degree n .

Recurrence relation

$$\begin{aligned} T_{n+1}(x) &= 2xT_n(x) - T_{n-1}(x), \\ \hat{T}_{n+1}(x) &= x\hat{T}_n(x) - \frac{1}{4}\hat{T}_{n-1}(x), \end{aligned}$$

where $\hat{T}_n(x) = T_n(x)/2^{n-1}$ is a monic Chebyshev polynomial.

Recurrence relation for orthogonal polynomials

Definition: A polynomial $p(x)$ is said to be **monic** if the coefficient of the highest power of x is 1.

Recurrence relation for orthogonal polynomials

Definition: A polynomial $p(x)$ is said to be **monic** if the coefficient of the highest power of x is 1.

Theorem: Let ϕ_{n+1} be monic polynomial of degree $n + 1$ such that $\phi_{n+1} \perp \mathcal{P}_n$. Then for $n = 0, 1, \dots$, we have

$$\phi_{n+1}(x) = (x - \alpha_n)\phi_n(x) - \beta_n\phi_{n-1}(x),$$

where $\phi_0(x) = 1, \phi_{-1}(x) = 0$ and

Recurrence relation for orthogonal polynomials

Definition: A polynomial $p(x)$ is said to be **monic** if the coefficient of the highest power of x is 1.

Theorem: Let ϕ_{n+1} be monic polynomial of degree $n+1$ such that $\phi_{n+1} \perp \mathcal{P}_n$. Then for $n = 0, 1, \dots$, we have

$$\phi_{n+1}(x) = (x - \alpha_n)\phi_n(x) - \beta_n\phi_{n-1}(x),$$

where $\phi_0(x) = 1, \phi_{-1}(x) = 0$ and

$$\alpha_n = \frac{\langle x\phi_n, \phi_n \rangle}{\langle \phi_n, \phi_n \rangle}, \quad \beta_n = \frac{\langle \phi_n, \phi_n \rangle}{\langle \phi_{n-1}, \phi_{n-1} \rangle} > 0.$$

Recurrence relation for orthogonal polynomials

Proof: Note that $\phi_{n+1} - x\phi_n \in \mathcal{P}_n$, hence

$$\phi_{n+1} - x\phi_n = \sum_{j=0}^n \frac{\langle \phi_{n+1} - x\phi_n, \phi_j \rangle}{\langle \phi_j, \phi_j \rangle} \phi_j$$

Recurrence relation for orthogonal polynomials

Proof: Note that $\phi_{n+1} - x\phi_n \in \mathcal{P}_n$, hence

$$\begin{aligned}\phi_{n+1} - x\phi_n &= \sum_{j=0}^n \frac{\langle \phi_{n+1} - x\phi_n, \phi_j \rangle}{\langle \phi_j, \phi_j \rangle} \phi_j = - \sum_{j=0}^n \frac{\langle x\phi_n, \phi_j \rangle}{\langle \phi_j, \phi_j \rangle} \phi_j \\ &= - \frac{\langle x\phi_n, \phi_n \rangle}{\langle \phi_n, \phi_n \rangle} \phi_n - \frac{\langle x\phi_n, \phi_{n-1} \rangle}{\langle \phi_{n-1}, \phi_{n-1} \rangle} \phi_{n-1}\end{aligned}$$

Recurrence relation for orthogonal polynomials

Proof: Note that $\phi_{n+1} - x\phi_n \in \mathcal{P}_n$, hence

$$\begin{aligned}\phi_{n+1} - x\phi_n &= \sum_{j=0}^n \frac{\langle \phi_{n+1} - x\phi_n, \phi_j \rangle}{\langle \phi_j, \phi_j \rangle} \phi_j = - \sum_{j=0}^n \frac{\langle x\phi_n, \phi_j \rangle}{\langle \phi_j, \phi_j \rangle} \phi_j \\ &= - \frac{\langle x\phi_n, \phi_n \rangle}{\langle \phi_n, \phi_n \rangle} \phi_n - \frac{\langle x\phi_n, \phi_{n-1} \rangle}{\langle \phi_{n-1}, \phi_{n-1} \rangle} \phi_{n-1}\end{aligned}$$

because $\sum_{j=0}^{n-2} \frac{\langle x\phi_n, \phi_j \rangle}{\langle \phi_j, \phi_j \rangle} \phi_j = 0$ as $x\phi_j \in \mathcal{P}_{n-1}$ for $j = 0, \dots, n-2$.

Recurrence relation for orthogonal polynomials

Proof: Note that $\phi_{n+1} - x\phi_n \in \mathcal{P}_n$, hence

$$\begin{aligned}\phi_{n+1} - x\phi_n &= \sum_{j=0}^n \frac{\langle \phi_{n+1} - x\phi_n, \phi_j \rangle}{\langle \phi_j, \phi_j \rangle} \phi_j = - \sum_{j=0}^n \frac{\langle x\phi_n, \phi_j \rangle}{\langle \phi_j, \phi_j \rangle} \phi_j \\ &= - \frac{\langle x\phi_n, \phi_n \rangle}{\langle \phi_n, \phi_n \rangle} \phi_n - \frac{\langle x\phi_n, \phi_{n-1} \rangle}{\langle \phi_{n-1}, \phi_{n-1} \rangle} \phi_{n-1}\end{aligned}$$

because $\sum_{j=0}^{n-2} \frac{\langle x\phi_n, \phi_j \rangle}{\langle \phi_j, \phi_j \rangle} \phi_j = 0$ as $x\phi_j \in \mathcal{P}_{n-1}$ for $j = 0, \dots, n-2$.

This shows that $\phi_{n+1}(x) = (x - \alpha_n)\phi_n(x) - \beta_n\phi_{n-1}(x)$, where

$$\alpha_n = \frac{\langle x\phi_n, \phi_n \rangle}{\langle \phi_n, \phi_n \rangle}, \quad \beta_n = \frac{\langle \phi_n, x\phi_{n-1} \rangle}{\langle \phi_{n-1}, \phi_{n-1} \rangle} = \frac{\langle \phi_n, \phi_n \rangle}{\langle \phi_{n-1}, \phi_{n-1} \rangle} > 0.$$

Recurrence relation for orthogonal polynomials

Proof: Note that $\phi_{n+1} - x\phi_n \in \mathcal{P}_n$, hence

$$\begin{aligned}\phi_{n+1} - x\phi_n &= \sum_{j=0}^n \frac{\langle \phi_{n+1} - x\phi_n, \phi_j \rangle}{\langle \phi_j, \phi_j \rangle} \phi_j = - \sum_{j=0}^n \frac{\langle x\phi_n, \phi_j \rangle}{\langle \phi_j, \phi_j \rangle} \phi_j \\ &= - \frac{\langle x\phi_n, \phi_n \rangle}{\langle \phi_n, \phi_n \rangle} \phi_n - \frac{\langle x\phi_n, \phi_{n-1} \rangle}{\langle \phi_{n-1}, \phi_{n-1} \rangle} \phi_{n-1}\end{aligned}$$

because $\sum_{j=0}^{n-2} \frac{\langle x\phi_n, \phi_j \rangle}{\langle \phi_j, \phi_j \rangle} \phi_j = 0$ as $x\phi_j \in \mathcal{P}_{n-1}$ for $j = 0, \dots, n-2$.

This shows that $\phi_{n+1}(x) = (x - \alpha_n)\phi_n(x) - \beta_n\phi_{n-1}(x)$, where

$$\alpha_n = \frac{\langle x\phi_n, \phi_n \rangle}{\langle \phi_n, \phi_n \rangle}, \quad \beta_n = \frac{\langle \phi_n, x\phi_{n-1} \rangle}{\langle \phi_{n-1}, \phi_{n-1} \rangle} = \frac{\langle \phi_n, \phi_n \rangle}{\langle \phi_{n-1}, \phi_{n-1} \rangle} > 0.$$

Indeed $x\phi_{n-1} = \phi_n + \text{lower order terms}$, we have $\langle \phi_n, x\phi_{n-1} \rangle = \langle \phi_n, \phi_n \rangle$ which yields β_n . ■

Zeros of orthogonal polynomials

Theorem: Let $\phi_0(x), \dots, \phi_{n+1}(x)$ be orthogonal polynomials such that

$$\phi_{n+1}(x) = (x - \alpha_n)\phi_n(x) - \beta_n\phi_{n-1}(x).$$

Define the Jacobi matrix

$$A = \begin{bmatrix} \alpha_0 & \sqrt{\beta_1} & & & \\ \sqrt{\beta_1} & \alpha_1 & \sqrt{\beta_2} & & \\ & \sqrt{\beta_2} & \ddots & \ddots & \\ & & \ddots & \ddots & \sqrt{\beta_n} \\ & & & \sqrt{\beta_n} & \alpha_n \end{bmatrix}.$$

Zeros of orthogonal polynomials

Theorem: Let $\phi_0(x), \dots, \phi_{n+1}(x)$ be orthogonal polynomials such that

$$\phi_{n+1}(x) = (x - \alpha_n)\phi_n(x) - \beta_n\phi_{n-1}(x).$$

Define the Jacobi matrix

$$A = \begin{bmatrix} \alpha_0 & \sqrt{\beta_1} & & & \\ \sqrt{\beta_1} & \alpha_1 & \sqrt{\beta_2} & & \\ & \sqrt{\beta_2} & \ddots & \ddots & \\ & & \ddots & \ddots & \sqrt{\beta_n} \\ & & & \sqrt{\beta_n} & \alpha_n \end{bmatrix}.$$

Then $\phi_{n+1}(x_j) = 0 \iff \det(A - x_j I) = 0$.

Zeros of orthogonal polynomials

Theorem: Let $\phi_0(x), \dots, \phi_{n+1}(x)$ be orthogonal polynomials such that

$$\phi_{n+1}(x) = (x - \alpha_n)\phi_n(x) - \beta_n\phi_{n-1}(x).$$

Define the Jacobi matrix

$$A = \begin{bmatrix} \alpha_0 & \sqrt{\beta_1} & & & \\ \sqrt{\beta_1} & \alpha_1 & \sqrt{\beta_2} & & \\ & \sqrt{\beta_2} & \ddots & \ddots & \\ & & \ddots & \ddots & \sqrt{\beta_n} \\ & & & \sqrt{\beta_n} & \alpha_n \end{bmatrix}.$$

Then $\phi_{n+1}(x_j) = 0 \iff \det(A - x_j I) = 0$. Let $\mathbf{v}_0, \dots, \mathbf{v}_n$ be orthonormal eigenvectors of A . Then

Zeros of orthogonal polynomials

Theorem: Let $\phi_0(x), \dots, \phi_{n+1}(x)$ be orthogonal polynomials such that

$$\phi_{n+1}(x) = (x - \alpha_n)\phi_n(x) - \beta_n\phi_{n-1}(x).$$

Define the Jacobi matrix

$$A = \begin{bmatrix} \alpha_0 & \sqrt{\beta_1} & & & \\ \sqrt{\beta_1} & \alpha_1 & \sqrt{\beta_2} & & \\ & \sqrt{\beta_2} & \ddots & \ddots & \\ & & \ddots & \ddots & \sqrt{\beta_n} \\ & & & \sqrt{\beta_n} & \alpha_n \end{bmatrix}.$$

Then $\phi_{n+1}(x_j) = 0 \iff \det(A - x_j I) = 0$. Let $\mathbf{v}_0, \dots, \mathbf{v}_n$ be orthonormal eigenvectors of A . Then $w_j = \langle \mathbf{1}, \mathbf{1} \rangle (\mathbf{e}_1^\top \mathbf{v}_j)^2$, $j = 0 : n$, are the required weights for the Gaussian quadrature rule.

Zeros of orthogonal polynomials

Theorem: Let $\phi_0(x), \dots, \phi_{n+1}(x)$ be orthogonal polynomials such that

$$\phi_{n+1}(x) = (x - \alpha_n)\phi_n(x) - \beta_n\phi_{n-1}(x).$$

Define the Jacobi matrix

$$A = \begin{bmatrix} \alpha_0 & \sqrt{\beta_1} & & & \\ \sqrt{\beta_1} & \alpha_1 & \sqrt{\beta_2} & & \\ & \sqrt{\beta_2} & \ddots & \ddots & \\ & & \ddots & \ddots & \sqrt{\beta_n} \\ & & & \sqrt{\beta_n} & \alpha_n \end{bmatrix}.$$

Then $\phi_{n+1}(x_j) = 0 \iff \det(A - x_j I) = 0$. Let $\mathbf{v}_0, \dots, \mathbf{v}_n$ be orthonormal eigenvectors of A . Then $w_j = \langle \mathbf{1}, \mathbf{1} \rangle (\mathbf{e}_1^\top \mathbf{v}_j)^2$, $j = 0 : n$, are the required weights for the Gaussian quadrature rule. Matlab code

```
[V, D] = eig(A); x = diag(D); w = transpose(inp * V(1, :).^2);
```

where $\text{inp} = \langle \mathbf{1}, \mathbf{1} \rangle$

Zeros of orthogonal polynomials

Proof: Define $\psi_j(x) := \phi_j(x)/\|\phi_j\|$. Dividing the recurrence relation by $\|\phi_n\|$ and noting that $\sqrt{\beta_n} = \|\phi_n\|/\|\phi_{n-1}\|$, we have

Zeros of orthogonal polynomials

Proof: Define $\psi_j(x) := \phi_j(x)/\|\phi_j\|$. Dividing the recurrence relation by $\|\phi_n\|$ and noting that $\sqrt{\beta_n} = \|\phi_n\|/\|\phi_{n-1}\|$, we have

$$\sqrt{\beta_{n+1}}\psi_{n+1}(x) = (x - \alpha_n)\psi_n(x) - \sqrt{\beta_n}\psi_{n-1}$$

which gives

Zeros of orthogonal polynomials

Proof: Define $\psi_j(x) := \phi_j(x)/\|\phi_j\|$. Dividing the recurrence relation by $\|\phi_n\|$ and noting that $\sqrt{\beta_n} = \|\phi_n\|/\|\phi_{n-1}\|$, we have

$$\sqrt{\beta_{n+1}}\psi_{n+1}(x) = (x - \alpha_n)\psi_n(x) - \sqrt{\beta_n}\psi_{n-1}$$

which gives

$$x\psi_n(x) = \sqrt{\beta_{n+1}}\psi_{n+1}(x) + \alpha_n\psi_n + \sqrt{\beta_n}\psi_{n-1}.$$

Zeros of orthogonal polynomials

Proof: Define $\psi_j(x) := \phi_j(x)/\|\phi_j\|$. Dividing the recurrence relation by $\|\phi_n\|$ and noting that $\sqrt{\beta_n} = \|\phi_n\|/\|\phi_{n-1}\|$, we have

$$\sqrt{\beta_{n+1}}\psi_{n+1}(x) = (x - \alpha_n)\psi_n(x) - \sqrt{\beta_n}\psi_{n-1}$$

which gives

$$x\psi_n(x) = \sqrt{\beta_{n+1}}\psi_{n+1}(x) + \alpha_n\psi_n + \sqrt{\beta_n}\psi_{n-1}.$$

$$x \begin{bmatrix} \psi_0(x) \\ \psi_1(x) \\ \vdots \\ \vdots \\ \psi_n(x) \end{bmatrix} = \begin{bmatrix} \alpha_0 & \sqrt{\beta_1} & & & \\ \sqrt{\beta_1} & \alpha_1 & \sqrt{\beta_2} & & \\ & \sqrt{\beta_2} & \ddots & \ddots & \\ & & \ddots & \ddots & \sqrt{\beta_n} \\ & & & \sqrt{\beta_n} & \alpha_n \end{bmatrix} \begin{bmatrix} \psi_0(x) \\ \psi_1(x) \\ \vdots \\ \vdots \\ \psi_n(x) \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ \sqrt{\beta_{n+1}}\psi_{n+1}(x) \end{bmatrix}.$$

Zeros of orthogonal polynomials

Proof: Define $\psi_j(x) := \phi_j(x)/\|\phi_j\|$. Dividing the recurrence relation by $\|\phi_n\|$ and noting that $\sqrt{\beta_n} = \|\phi_n\|/\|\phi_{n-1}\|$, we have

$$\sqrt{\beta_{n+1}}\psi_{n+1}(x) = (x - \alpha_n)\psi_n(x) - \sqrt{\beta_n}\psi_{n-1}$$

which gives

$$x\psi_n(x) = \sqrt{\beta_{n+1}}\psi_{n+1}(x) + \alpha_n\psi_n + \sqrt{\beta_n}\psi_{n-1}.$$

$$x \begin{bmatrix} \psi_0(x) \\ \psi_1(x) \\ \vdots \\ \vdots \\ \psi_n(x) \end{bmatrix} = \begin{bmatrix} \alpha_0 & \sqrt{\beta_1} & & & \\ \sqrt{\beta_1} & \alpha_1 & \sqrt{\beta_2} & & \\ & \sqrt{\beta_2} & \ddots & \ddots & \\ & & \ddots & \ddots & \sqrt{\beta_n} \\ & & & \sqrt{\beta_n} & \alpha_n \end{bmatrix} \begin{bmatrix} \psi_0(x) \\ \psi_1(x) \\ \vdots \\ \vdots \\ \psi_n(x) \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ \sqrt{\beta_{n+1}}\psi_{n+1}(x) \end{bmatrix}.$$

This shows that

$$\phi_{n+1}(x_j) = 0 \iff \psi_{n+1}(x_j) = 0 \iff \det(A - x_j I) = 0.$$

Zeros of orthogonal polynomials

Proof: Define $\psi_j(x) := \phi_j(x)/\|\phi_j\|$. Dividing the recurrence relation by $\|\phi_n\|$ and noting that $\sqrt{\beta_n} = \|\phi_n\|/\|\phi_{n-1}\|$, we have

$$\sqrt{\beta_{n+1}}\psi_{n+1}(x) = (x - \alpha_n)\psi_n(x) - \sqrt{\beta_n}\psi_{n-1}$$

which gives

$$x\psi_n(x) = \sqrt{\beta_{n+1}}\psi_{n+1}(x) + \alpha_n\psi_n + \sqrt{\beta_n}\psi_{n-1}.$$

$$x \begin{bmatrix} \psi_0(x) \\ \psi_1(x) \\ \vdots \\ \vdots \\ \psi_n(x) \end{bmatrix} = \begin{bmatrix} \alpha_0 & \sqrt{\beta_1} & & & \\ \sqrt{\beta_1} & \alpha_1 & \sqrt{\beta_2} & & \\ & \sqrt{\beta_2} & \ddots & \ddots & \\ & & \ddots & \ddots & \sqrt{\beta_n} \\ & & & \sqrt{\beta_n} & \alpha_n \end{bmatrix} \begin{bmatrix} \psi_0(x) \\ \psi_1(x) \\ \vdots \\ \vdots \\ \psi_n(x) \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ \sqrt{\beta_{n+1}}\psi_{n+1}(x) \end{bmatrix}.$$

This shows that

$$\phi_{n+1}(x_j) = 0 \iff \psi_{n+1}(x_j) = 0 \iff \det(A - x_j I) = 0.$$

Proof

Observe that $\mathbf{u}_j := [\psi_0(x_j) \ \cdots \ \psi_n(x_j)]^\top$ is an eigenvector of A corresponding to the eigenvalue x_j . Hence $\mathbf{u}_j = \mathbf{v}_j d_j$ for some $d_j \in \mathbb{R}$.

Proof

Observe that $\mathbf{u}_j := [\psi_0(x_j) \ \cdots \ \psi_n(x_j)]^\top$ is an eigenvector of A corresponding to the eigenvalue x_j . Hence $\mathbf{u}_j = \mathbf{v}_j d_j$ for some $d_j \in \mathbb{R}$.

Now $\psi_0(x) = \phi_0(x)/\|\phi_0\| = 1/\|\phi_0\|$ shows that

$$d_j \mathbf{e}_1^\top \mathbf{v}_j = \mathbf{e}_1^\top \mathbf{u}_j = \psi_0(x_j) = 1/\|\phi_0\| \implies d_j = 1/(\|\phi_0\| \mathbf{e}_1^\top \mathbf{v}_j).$$

Proof

Observe that $\mathbf{u}_j := [\psi_0(x_j) \ \cdots \ \psi_n(x_j)]^\top$ is an eigenvector of A corresponding to the eigenvalue x_j . Hence $\mathbf{u}_j = \mathbf{v}_j d_j$ for some $d_j \in \mathbb{R}$.

Now $\psi_0(x) = \phi_0(x)/\|\phi_0\| = 1/\|\phi_0\|$ shows that

$$d_j \mathbf{e}_1^\top \mathbf{v}_j = \mathbf{e}_1^\top \mathbf{u}_j = \psi_0(x_j) = 1/\|\phi_0\| \implies d_j = 1/(\|\phi_0\| \mathbf{e}_1^\top \mathbf{v}_j).$$

Set $U := [\mathbf{u}_0 \ \cdots \ \mathbf{u}_n]$ and $V := [\mathbf{v}_0 \ \cdots \ \mathbf{v}_n]$. Then $U = VD$, where $D = \text{diag}(d_0, \dots, d_n)$.

Proof

Observe that $\mathbf{u}_j := [\psi_0(x_j) \ \cdots \ \psi_n(x_j)]^\top$ is an eigenvector of A corresponding to the eigenvalue x_j . Hence $\mathbf{u}_j = \mathbf{v}_j d_j$ for some $d_j \in \mathbb{R}$.

Now $\psi_0(x) = \phi_0(x)/\|\phi_0\| = 1/\|\phi_0\|$ shows that

$$d_j \mathbf{e}_1^\top \mathbf{v}_j = \mathbf{e}_1^\top \mathbf{u}_j = \psi_0(x_j) = 1/\|\phi_0\| \implies d_j = 1/(\|\phi_0\| \mathbf{e}_1^\top \mathbf{v}_j).$$

Set $U := [\mathbf{u}_0 \ \cdots \ \mathbf{u}_n]$ and $V := [\mathbf{v}_0 \ \cdots \ \mathbf{v}_n]$. Then $U = VD$, where $D = \text{diag}(d_0, \dots, d_n)$.

By exactness of $G_n(f)$, we have

$$w_0 \psi_0(x_0) + \cdots + w_n \psi_0(x_n) = G_n(\psi_0) = \int_a^b \psi_0(x) \mu(x) dx = \langle 1, 1 \rangle / \|\phi_0\|$$

$$w_0 \psi_j(x_0) + \cdots + w_n \psi_j(x_n) = G_n(\psi_j) = \int_a^b \psi_j(x) \mu(x) dx = \langle \psi_j, 1 \rangle = 0$$

for $j = 1 : n$.

Proof

Observe that $\mathbf{u}_j := [\psi_0(x_j) \ \cdots \ \psi_n(x_j)]^\top$ is an eigenvector of A corresponding to the eigenvalue x_j . Hence $\mathbf{u}_j = \mathbf{v}_j d_j$ for some $d_j \in \mathbb{R}$.

Now $\psi_0(x) = \phi_0(x)/\|\phi_0\| = 1/\|\phi_0\|$ shows that

$$d_j \mathbf{e}_1^\top \mathbf{v}_j = \mathbf{e}_1^\top \mathbf{u}_j = \psi_0(x_j) = 1/\|\phi_0\| \implies d_j = 1/(\|\phi_0\| \mathbf{e}_1^\top \mathbf{v}_j).$$

Set $U := [\mathbf{u}_0 \ \cdots \ \mathbf{u}_n]$ and $V := [\mathbf{v}_0 \ \cdots \ \mathbf{v}_n]$. Then $U = VD$, where $D = \text{diag}(d_0, \dots, d_n)$.

By exactness of $G_n(f)$, we have

$$w_0 \psi_0(x_0) + \cdots + w_n \psi_0(x_n) = G_n(\psi_0) = \int_a^b \psi_0(x) \mu(x) dx = \langle 1, 1 \rangle / \|\phi_0\|$$

$$w_0 \psi_j(x_0) + \cdots + w_n \psi_j(x_n) = G_n(\psi_j) = \int_a^b \psi_j(x) \mu(x) dx = \langle \psi_j, 1 \rangle = 0$$

for $j = 1 : n$. Thus $U\mathbf{w} = (\langle 1, 1 \rangle / \|\phi_0\|) \mathbf{e}_1$, where $\mathbf{w} := [w_0 \ \cdots \ w_n]^\top$.

Proof

Since V is orthogonal, we have $U^{-1} = (VD)^{-1} = D^{-1}V^{\top}$. Hence

$$\mathbf{w} = (\langle \mathbf{1}, \mathbf{1} \rangle / \|\phi_0\|) U^{-1} \mathbf{e}_1 = (\langle \mathbf{1}, \mathbf{1} \rangle / \|\phi_0\|) D^{-1} V^{\top} \mathbf{e}_1.$$

Proof

Since V is orthogonal, we have $U^{-1} = (VD)^{-1} = D^{-1}V^T$. Hence

$$\mathbf{w} = (\langle \mathbf{1}, \mathbf{1} \rangle / \|\phi_0\|) U^{-1} \mathbf{e}_1 = (\langle \mathbf{1}, \mathbf{1} \rangle / \|\phi_0\|) D^{-1} V^T \mathbf{e}_1.$$

Now using the fact that $d_j = 1/(\|\phi_0\| \mathbf{e}_1^T \mathbf{v}_j)$, we have

$$\begin{aligned} w_j &= \mathbf{e}_j^T \mathbf{w} = (\langle \mathbf{1}, \mathbf{1} \rangle / \|\phi_0\|) \mathbf{e}_j^T D^{-1} V^T \mathbf{e}_1 \\ &= (\langle \mathbf{1}, \mathbf{1} \rangle / (\|\phi_0\| d_j)) \mathbf{e}_j^T V^T \mathbf{e}_1 = \langle \mathbf{1}, \mathbf{1} \rangle (\mathbf{e}_1^T V \mathbf{e}_j)^2 \\ &= \langle \mathbf{1}, \mathbf{1} \rangle (\mathbf{e}_1^T \mathbf{v}_j)^2. \quad \blacksquare \end{aligned}$$

Thus the weights w and nodes x can be computed in MATLAB as

```
[V, D] = eig(A); x = diag(D); w = transpose(inp*V(1, :).^2);
```

where $\text{inp} = \langle \mathbf{1}, \mathbf{1} \rangle$

Gauss Legendre quadrature

Evaluate $\int_{-1}^1 f(x)dx$.

Gauss Legendre quadrature

Evaluate $\int_{-1}^1 f(x)dx$. Inner product: $\langle f, g \rangle = \int_{-1}^1 f(x)g(x)dx$.

Gauss Legendre quadrature

Evaluate $\int_{-1}^1 f(x)dx$. Inner product: $\langle f, g \rangle = \int_{-1}^1 f(x)g(x)dx$.

Recurrence relation ([monic Legendre polynomial](#)):

$$P_{n+1}(x) = xP_n(x) - \frac{n^2}{4n^2 - 1}P_{n-1}(x).$$

Gauss Legendre quadrature

Evaluate $\int_{-1}^1 f(x)dx$. Inner product: $\langle f, g \rangle = \int_{-1}^1 f(x)g(x)dx$.

Recurrence relation ([monic Legendre polynomial](#)):

$$P_{n+1}(x) = xP_n(x) - \frac{n^2}{4n^2 - 1}P_{n-1}(x).$$

Here $\alpha_n := 0$ for $n = 0, 1, \dots$ and $\beta_n := \frac{n^2}{4n^2 - 1}$ for $n = 1, 2, \dots$

$$A = \begin{bmatrix} 0 & \frac{1}{\sqrt{3}} & & & \\ \frac{1}{\sqrt{3}} & 0 & \frac{2}{\sqrt{15}} & & \\ & \frac{2}{\sqrt{15}} & \ddots & \ddots & \\ & & \ddots & \ddots & \frac{n}{\sqrt{4n^2-1}} \\ & & & \frac{n}{\sqrt{4n^2-1}} & 0 \end{bmatrix}.$$

Gauss Legendre quadrature

Evaluate $\int_{-1}^1 f(x)dx$. Inner product: $\langle f, g \rangle = \int_{-1}^1 f(x)g(x)dx$.

Recurrence relation ([monic Legendre polynomial](#)):

$$P_{n+1}(x) = xP_n(x) - \frac{n^2}{4n^2 - 1}P_{n-1}(x).$$

Here $\alpha_n := 0$ for $n = 0, 1, \dots$ and $\beta_n := \frac{n^2}{4n^2 - 1}$ for $n = 1, 2, \dots$

$$A = \begin{bmatrix} 0 & \frac{1}{\sqrt{3}} & & & \\ \frac{1}{\sqrt{3}} & 0 & \frac{2}{\sqrt{15}} & & \\ & \frac{2}{\sqrt{15}} & \ddots & \ddots & \\ & & \ddots & \ddots & \frac{n}{\sqrt{4n^2-1}} \\ & & & \frac{n}{\sqrt{4n^2-1}} & 0 \end{bmatrix}.$$

Thus the weights w and nodes x can be computed in MATLAB as

```
[V, D] = eig(A); x = diag(D); w = transpose(2*V(1, :).^2);
```


Gauss Laguerre quadrature

Evaluate $\int_0^\infty f(x)e^{-x}dx$.

Gauss Laguerre quadrature

Evaluate $\int_0^\infty f(x)e^{-x}dx$. Inner product $\langle f, g \rangle = \int_0^\infty f(x)g(x)e^{-x}dx$.

Gauss Laguerre quadrature

Evaluate $\int_0^\infty f(x)e^{-x}dx$. Inner product $\langle f, g \rangle = \int_0^\infty f(x)g(x)e^{-x}dx$.

Recurrence relation (monic Laguerre):

$$L_{n+1}(x) = (x - (2n + 1))L_n(x) - n^2L_{n-1}(x).$$

Here $\alpha_n := 2n + 1$ for $n = 0, 1, \dots$ and $\beta_n := n^2$ for $n = 1, 2, \dots$

Gauss Laguerre quadrature

Evaluate $\int_0^\infty f(x)e^{-x}dx$. Inner product $\langle f, g \rangle = \int_0^\infty f(x)g(x)e^{-x}dx$.

Recurrence relation (monic Laguerre):

$$L_{n+1}(x) = (x - (2n + 1))L_n(x) - n^2L_{n-1}(x).$$

Here $\alpha_n := 2n + 1$ for $n = 0, 1, \dots$ and $\beta_n := n^2$ for $n = 1, 2, \dots$

$$A = \begin{bmatrix} 1 & 1 & & & \\ 1 & 3 & 2 & & \\ & 2 & \ddots & \ddots & \\ & & \ddots & \ddots & n \\ & & & n & 2n+1 \end{bmatrix}.$$

Gauss Laguerre quadrature

Evaluate $\int_0^\infty f(x)e^{-x}dx$. Inner product $\langle f, g \rangle = \int_0^\infty f(x)g(x)e^{-x}dx$.

Recurrence relation (monic Laguerre):

$$L_{n+1}(x) = (x - (2n + 1))L_n(x) - n^2L_{n-1}(x).$$

Here $\alpha_n := 2n + 1$ for $n = 0, 1, \dots$ and $\beta_n := n^2$ for $n = 1, 2, \dots$

$$A = \begin{bmatrix} 1 & 1 & & & \\ 1 & 3 & 2 & & \\ & 2 & \ddots & \ddots & \\ & & \ddots & \ddots & n \\ & & & n & 2n+1 \end{bmatrix}.$$

Thus the weights w and nodes x can be computed in MATLAB as

```
[V, D] = eig(A); x = diag(D); w = transpose(V(1, :).^2);
```

Gauss Chebyshev quadrature

Evaluate $\int_{-1}^1 f(x)(1-x^2)^{-1/2}dx$.

Gauss Chebyshev quadrature

Evaluate $\int_{-1}^1 f(x)(1-x^2)^{-1/2} dx$.

Inner product $\langle f, g \rangle = \int_{-1}^1 f(x)g(x)(1-x^2)^{-1/2} dx$.

Gauss Chebyshev quadrature

Evaluate $\int_{-1}^1 f(x)(1-x^2)^{-1/2}dx$.

Inner product $\langle f, g \rangle = \int_{-1}^1 f(x)g(x)(1-x^2)^{-1/2}dx$.

Recurrence relation (monic Chebyshev polynomial)

$$\hat{T}_{n+1}(x) = x\hat{T}_n(x) - \frac{1}{4}\hat{T}_{n-1}(x).$$

Gauss Chebyshev quadrature

Evaluate $\int_{-1}^1 f(x)(1-x^2)^{-1/2}dx$.

Inner product $\langle f, g \rangle = \int_{-1}^1 f(x)g(x)(1-x^2)^{-1/2}dx$.

Recurrence relation (monic Chebyshev polynomial)

$$\hat{T}_{n+1}(x) = x\hat{T}_n(x) - \frac{1}{4}\hat{T}_{n-1}(x).$$

Here $\alpha_n = 0$ for $n = 0, 1, \dots$ and $\beta_n = 1/4$ for $n = 1, 2, \dots$

$$A = \begin{bmatrix} 0 & \frac{1}{2} & & & \\ \frac{1}{2} & 0 & \frac{1}{2} & & \\ & \frac{1}{2} & \ddots & \ddots & \\ & & \ddots & \ddots & \frac{1}{2} \\ & & & \frac{1}{2} & 0 \end{bmatrix}.$$

Gauss Chebyshev quadrature

Evaluate $\int_{-1}^1 f(x)(1-x^2)^{-1/2}dx$.

Inner product $\langle f, g \rangle = \int_{-1}^1 f(x)g(x)(1-x^2)^{-1/2}dx$.

Recurrence relation (monic Chebyshev polynomial)

$$\hat{T}_{n+1}(x) = x\hat{T}_n(x) - \frac{1}{4}\hat{T}_{n-1}(x).$$

Here $\alpha_n = 0$ for $n = 0, 1, \dots$ and $\beta_n = 1/4$ for $n = 1, 2, \dots$

$$A = \begin{bmatrix} 0 & \frac{1}{2} & & & \\ \frac{1}{2} & 0 & \frac{1}{2} & & \\ & \frac{1}{2} & \ddots & \ddots & \\ & & \ddots & \ddots & \frac{1}{2} \\ & & & \frac{1}{2} & 0 \end{bmatrix}.$$

Thus the weights w and nodes x can be computed in MATLAB as

```
[V, D] = eig(A); x = diag(D); w = transpose(pi*V(1, :).^2);
```