

MA579H Scientific Computing

Solutions of Nonlinear Equations I

Rafikul Alam
Department of Mathematics
IIT Guwahati

Outline

- Bisection method
- Regula-Falsi method
- Secant method

Zeros of Bessel's function

The Bessel function of the first kind

$$J_{\alpha}(x) = \sum_{m=0}^{\infty} \frac{(-1)^m}{m! \Gamma(m + \alpha + 1)} \left(\frac{x}{2}\right)^{2m+\alpha}$$

is a solution of the Bessel's differential equation

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + (x^2 - \alpha^2)y = 0,$$

where $\Gamma(z)$ is the gamma function.

Zeros of Bessel's function

The Bessel function of the first kind

$$J_{\alpha}(x) = \sum_{m=0}^{\infty} \frac{(-1)^m}{m! \Gamma(m + \alpha + 1)} \left(\frac{x}{2}\right)^{2m+\alpha}$$

is a solution of the Bessel's differential equation

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + (x^2 - \alpha^2)y = 0,$$

where $\Gamma(z)$ is the gamma function. The positive zeros of $J_{\alpha}(x)$ are resonance frequencies.

Zeros of Bessel's function

The Bessel function of the first kind

$$J_{\alpha}(x) = \sum_{m=0}^{\infty} \frac{(-1)^m}{m! \Gamma(m + \alpha + 1)} \left(\frac{x}{2}\right)^{2m+\alpha}$$


is a solution of the Bessel's differential equation

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + (x^2 - \alpha^2)y = 0,$$

where $\Gamma(z)$ is the gamma function. The positive zeros of $J_{\alpha}(x)$ are resonance frequencies.


Plot of $J_3(x)$

```
J3 = @(x) besselj(3,x);  
fplot(J3,[0 20]), grid on  
xlabel('x'), ylabel('J_3(x)')  
title('Bessel function')
```



Plot of zeros of $J_3(x)$

```
z = [];  
for guess = [6,10,13,16,19]  
    z = [z;fzero(J3,guess)];  
end  
hold on, plot(z,J3(z),'o')  
title('Bessel roots')
```



Zeros of $J_3(x)$

Zeros of $J_3(x)$ computed by fzero in $[0, 20]$.

z =

6.380161895923984e+00

9.761023129981670e+00

1.301520072169843e+01

1.622346616031877e+01

1.940941522643502e+01

Nonlinear equations

Single nonlinear equation in one variable:

Given $f : [a, b] \rightarrow \mathbb{R}$, solve $f(x) = 0$.

Nonlinear equations

Single nonlinear equation in one variable:

Given $f : [a, b] \rightarrow \mathbb{R}$, solve $f(x) = 0$.

Example: Consider the nonlinear equation

$$x^2 - \sin(x) - .5 = 0$$

for which $x = 1.1961$ is an approximate solution.

Nonlinear equations

Single nonlinear equation in one variable:

Given $f : [a, b] \rightarrow \mathbb{R}$, solve $f(x) = 0$.

Example: Consider the nonlinear equation

$$x^2 - \sin(x) - .5 = 0$$

for which $x = 1.1961$ is an approximate solution.

System of nonlinear equations in several variables:

Given $F : U \subset \mathbb{R}^n \rightarrow \mathbb{R}^n$, solve $F(\mathbf{x}) = 0$.

Nonlinear equations

Single nonlinear equation in one variable:

Given $f : [a, b] \rightarrow \mathbb{R}$, solve $f(x) = 0$.

Example: Consider the nonlinear equation

$$x^2 - \sin(x) - .5 = 0$$

for which $x = 1.1961$ is an approximate solution.

System of nonlinear equations in several variables:

Given $F : U \subset \mathbb{R}^n \rightarrow \mathbb{R}^n$, solve $F(\mathbf{x}) = 0$.

Example: Consider the system of nonlinear equations in two dimensions

$$\begin{aligned}x_1^2 - x_2 + 0.25 &= 0 \\ -x_1 + x_2^2 + 0.25 &= 0\end{aligned}$$

for which $\mathbf{x} = [0.5 \ 0.5]^\top$ is a solution.

Convergence of Iterative Methods

An iterative method with initial guess x_0 for solving $f(x) = 0$ produces a sequence $\{x_n\}_{n=1}^{\infty}$ with a hope that x_n converges to a solution α .

Convergence of Iterative Methods

An iterative method with initial guess x_0 for solving $f(x) = 0$ produces a sequence $\{x_n\}_{n=1}^{\infty}$ with a hope that x_n converges to a solution α .

Rate of convergence: Let x_n be the n^{th} iterate of an iterative algorithm that converges to α . The order of convergence is said to be

- **linear** if there exists $0 < r < 1$ such that $\lim_{n \rightarrow \infty} \frac{\|x_{n+1} - \alpha\|}{\|x_n - \alpha\|} = r$. Then r is called the rate of convergence.

Example: $1, 10, 10^{-1}, 10^{-2}, \dots$

Convergence of Iterative Methods

An iterative method with initial guess x_0 for solving $f(x) = 0$ produces a sequence $\{x_n\}_{n=1}^{\infty}$ with a hope that x_n converges to a solution α .

Rate of convergence: Let x_n be the n^{th} iterate of an iterative algorithm that converges to α . The order of convergence is said to be

- **linear** if there exists $0 < r < 1$ such that $\lim_{n \rightarrow \infty} \frac{\|x_{n+1} - \alpha\|}{\|x_n - \alpha\|} = r$. Then r is called the rate of convergence.

Example: $1, 10, 10^{-1}, 10^{-2}, \dots$

- **quadratic** if $\lim_{n \rightarrow \infty} \frac{\|x_{n+1} - \alpha\|}{\|x_n - \alpha\|^2} = r$ for some real number $r > 0$.

Example: $10^{-1}, 10^{-2}, 10^{-4}, 10^{-8} \dots$

Convergence of Iterative Methods

An iterative method with initial guess x_0 for solving $f(x) = 0$ produces a sequence $\{x_n\}_{n=1}^{\infty}$ with a hope that x_n converges to a solution α .

Rate of convergence: Let x_n be the n^{th} iterate of an iterative algorithm that converges to α . The order of convergence is said to be

- **linear** if there exists $0 < r < 1$ such that $\lim_{n \rightarrow \infty} \frac{\|x_{n+1} - \alpha\|}{\|x_n - \alpha\|} = r$. Then r is called the rate of convergence.

Example: $1, 10, 10^{-1}, 10^{-2}, \dots$

- **quadratic** if $\lim_{n \rightarrow \infty} \frac{\|x_{n+1} - \alpha\|}{\|x_n - \alpha\|^2} = r$ for some real number $r > 0$.

Example: $10^{-1}, 10^{-2}, 10^{-4}, 10^{-8} \dots$

- **cubic** if $\lim_{n \rightarrow \infty} \frac{\|x_{n+1} - \alpha\|}{\|x_n - \alpha\|^3} = r$ for some real number $r > 0$.

Example: $10^{-1}, 10^{-8}, 10^{-24}, 10^{-128} \dots$

Bisection method

Let $f : \mathbb{R} \longrightarrow \mathbb{R}$ be continuous.

Bisection method

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous.

Bisection method begins with an interval $[a, b]$ containing a zero of f and repeatedly halves its length until a solution is found with a desired accuracy.

Bisection method

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous.

Bisection method begins with an interval $[a, b]$ containing a zero of f and repeatedly halves its length until a solution is found with a desired accuracy.

Input: $a, b \in \mathbb{R}$ such that $f(a)f(b) < 0$.

Output Approximate solution m of $f(x) = 0$.

Bisection method

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous.

Bisection method begins with an interval $[a, b]$ containing a zero of f and repeatedly halves its length until a solution is found with a desired accuracy.

Input: $a, b \in \mathbb{R}$ such that $f(a)f(b) < 0$.

Output Approximate solution m of $f(x) = 0$.

- **while** ($|b - a| > \mathbf{tol}$) **do** % e.g., $\mathbf{tol} = \mathbf{eps} * \mathbf{abs}(b)$
- $m = a + (b - a)/2$

Bisection method

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous.

Bisection method begins with an interval $[a, b]$ containing a zero of f and repeatedly halves its length until a solution is found with a desired accuracy.

Input: $a, b \in \mathbb{R}$ such that $f(a)f(b) < 0$.

Output Approximate solution m of $f(x) = 0$.

- **while** ($|b - a| > \mathbf{tol}$) **do** % e.g., $\mathbf{tol} = \mathbf{eps} * \mathbf{abs}(b)$
- $m = a + (b - a)/2$
- **if** $\mathbf{sign}(f(a)) = \mathbf{sign}(f(m))$ **then**

Bisection method

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous.

Bisection method begins with an interval $[a, b]$ containing a zero of f and repeatedly halves its length until a solution is found with a desired accuracy.

Input: $a, b \in \mathbb{R}$ such that $f(a)f(b) < 0$.

Output Approximate solution m of $f(x) = 0$.

- **while** ($|b - a| > \mathbf{tol}$) **do** % e.g., $\mathbf{tol} = \mathbf{eps} * \mathbf{abs}(b)$
- $m = a + (b - a)/2$
- **if** $\mathbf{sign}(f(a)) = \mathbf{sign}(f(m))$ **then**
- $a = m$ **else** $b = m$

Bisection method

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous.

Bisection method begins with an interval $[a, b]$ containing a zero of f and repeatedly halves its length until a solution is found with a desired accuracy.

Input: $a, b \in \mathbb{R}$ such that $f(a)f(b) < 0$.

Output Approximate solution m of $f(x) = 0$.

- **while** ($|b - a| > \mathbf{tol}$) **do** % e.g., $\mathbf{tol} = \mathbf{eps} * \mathbf{abs}(b)$
- $m = a + (b - a)/2$
- **if** $\mathbf{sign}(f(a)) = \mathbf{sign}(f(m))$ **then**
- $a = m$ **else** $b = m$

If x_n denote the midpoint after n -th iteration and if $\alpha \in [a, b]$ is a zero then $|x_n - \alpha| \leq 2^{-n}(b - a)$.

Bisection method

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous.

Bisection method begins with an interval $[a, b]$ containing a zero of f and repeatedly halves its length until a solution is found with a desired accuracy.

Input: $a, b \in \mathbb{R}$ such that $f(a)f(b) < 0$.

Output Approximate solution m of $f(x) = 0$.

- **while** ($|b - a| > \mathbf{tol}$) **do** % e.g., $\mathbf{tol} = \mathbf{eps} * \mathbf{abs}(b)$
- $m = a + (b - a)/2$
- **if** $\mathbf{sign}(f(a)) = \mathbf{sign}(f(m))$ **then**
- $a = m$ **else** $b = m$

If x_n denote the midpoint after n -th iteration and if $\alpha \in [a, b]$ is a zero then $|x_n - \alpha| \leq 2^{-n}(b - a)$. Set $\epsilon_n := 2^{-n}(b - a)$.

Bisection method

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous.

Bisection method begins with an interval $[a, b]$ containing a zero of f and repeatedly halves its length until a solution is found with a desired accuracy.

Input: $a, b \in \mathbb{R}$ such that $f(a)f(b) < 0$.

Output Approximate solution m of $f(x) = 0$.

- **while** ($|b - a| > \text{tol}$) **do** % e.g., $\text{tol} = \text{eps} * \text{abs}(b)$
- $m = a + (b - a)/2$
- **if** $\text{sign}(f(a)) = \text{sign}(f(m))$ **then**
- $a = m$ **else** $b = m$

If x_n denote the midpoint after n -th iteration and if $\alpha \in [a, b]$ is a zero then $|x_n - \alpha| \leq 2^{-n}(b - a)$. Set $\epsilon_n := 2^{-n}(b - a)$. Then

$$|x_n - \alpha| \leq \epsilon_n \text{ and } \frac{\epsilon_{n+1}}{\epsilon_n} = \frac{1}{2} < 1.$$

Thus the convergence is **linear** and the rate of convergence is $1/2$.


Example

Solution of $x^2 - \sin(x) - .5 = 0$ using bisection method.

Example

Solution of $x^2 - \sin(x) - .5 = 0$ using bisection method.

```
f = @(x) x.^2 -sin(x) -0.5;  
fplot(f, [0, 2], 'LineWidth', 2), grid on  
hold on  
xc = bisection(f, 0, 2, 1e-12)  
plot( xc, f(xc), 'o', 'LineWidth', 2)  
xc = 1.196082033297898e+00
```



Bisection method

A solution is correct to p decimal places if the error is less than $\frac{10^{-p}}{2}$.

Bisection method

A solution is correct to p decimal places if the error is less than $\frac{10^{-p}}{2}$.

How many iterations are necessary for the Bisection Method to provide solution correct to p decimal places?

Bisection method

A solution is correct to p decimal places if the error is less than $\frac{10^{-p}}{2}$.

How many iterations are necessary for the Bisection Method to provide solution correct to p decimal places?

The desired accuracy is assured if $2^{-n}(b - a) < \frac{10^{-p}}{2}$ which gives

$$n > 1 + \log_2(10^p(b - a))$$

regardless of function f involved.

Bisection method

A solution is correct to p decimal places if the error is less than $\frac{10^{-p}}{2}$.

How many iterations are necessary for the Bisection Method to provide solution correct to p decimal places?

The desired accuracy is assured if $2^{-n}(b - a) < \frac{10^{-p}}{2}$ which gives

$$n > 1 + \log_2(10^p(b - a))$$

regardless of function f involved.

- Bisection method makes no use of magnitudes of function values, only their signs.

Bisection method

A solution is correct to p decimal places if the error is less than $\frac{10^{-p}}{2}$.

How many iterations are necessary for the Bisection Method to provide solution correct to p decimal places?

The desired accuracy is assured if $2^{-n}(b - a) < \frac{10^{-p}}{2}$ which gives

$$n > 1 + \log_2(10^p(b - a))$$

regardless of function f involved.

- Bisection method makes no use of magnitudes of function values, only their signs.
- Bisection method is guaranteed to converge, but does so slowly as the convergence is linear.

Bisection method

A solution is correct to p decimal places if the error is less than $\frac{10^{-p}}{2}$.

How many iterations are necessary for the Bisection Method to provide solution correct to p decimal places?

The desired accuracy is assured if $2^{-n}(b - a) < \frac{10^{-p}}{2}$ which gives

$$n > 1 + \log_2(10^p(b - a))$$

regardless of function f involved.

- Bisection method makes no use of magnitudes of function values, only their signs.
- Bisection method is guaranteed to converge, but does so slowly as the convergence is linear.
- The number of iteration n required for $|x_n - \alpha| < \mathbf{tol}$ is given by

$$n \geq \log_2 \left(\frac{b - a}{\mathbf{tol}} \right),$$

regardless of function f involved.

Regula-Falsi method

Again we assume that the interval $[a, b]$ is such that $f(a)f(b) < 0$.

Regula-Falsi method

Again we assume that the interval $[a, b]$ is such that $f(a)f(b) < 0$. Starting with $a_1 = a$ and $b_1 = b$, we generate a sequence of nested intervals $[a_n, b_n]$ such that $f(a_n)f(b_n) < 0$ for $n = 1, 2, \dots$

Regula-Falsi method

Again we assume that the interval $[a, b]$ is such that $f(a)f(b) < 0$. Starting with $a_1 = a$ and $b_1 = b$, we generate a sequence of nested intervals $[a_n, b_n]$ such that $f(a_n)f(b_n) < 0$ for $n = 1, 2, \dots$

But unlike bisection method, we will not choose the midpoint $x_n := (a_n + b_n)/2$ to determine the next interval $[a_{n+1}, b_{n+1}]$.

Regula-Falsi method

Again we assume that the interval $[a, b]$ is such that $f(a)f(b) < 0$. Starting with $a_1 = a$ and $b_1 = b$, we generate a sequence of nested intervals $[a_n, b_n]$ such that $f(a_n)f(b_n) < 0$ for $n = 1, 2, \dots$

But unlike bisection method, we will not choose the midpoint $x_n := (a_n + b_n)/2$ to determine the next interval $[a_{n+1}, b_{n+1}]$.

Instead, we choose x_n to be the root of the interpolating polynomial $p(x)$ that interpolates the data $(a_n, f(a_n)), (b_n, f(b_n))$ (secant cuts the x -axis)

Regula-Falsi method

Again we assume that the interval $[a, b]$ is such that $f(a)f(b) < 0$. Starting with $a_1 = a$ and $b_1 = b$, we generate a sequence of **nested intervals** $[a_n, b_n]$ such that $f(a_n)f(b_n) < 0$ for $n = 1, 2, \dots$.

But unlike bisection method, we will not choose the **midpoint** $x_n := (a_n + b_n)/2$ to determine the next interval $[a_{n+1}, b_{n+1}]$.

Instead, we choose x_n to be the root of the interpolating polynomial $p(x)$ that interpolates the data $(a_n, f(a_n)), (b_n, f(b_n))$ (secant cuts the x -axis)

$$p(x) = \frac{(x - b_n)}{a_n - b_n} f(a_n) + \frac{(x - a_n)}{b_n - a_n} f(b_n).$$

Regula-Falsi method

Again we assume that the interval $[a, b]$ is such that $f(a)f(b) < 0$. Starting with $a_1 = a$ and $b_1 = b$, we generate a sequence of **nested intervals** $[a_n, b_n]$ such that $f(a_n)f(b_n) < 0$ for $n = 1, 2, \dots$

But unlike bisection method, we will not choose the **midpoint** $x_n := (a_n + b_n)/2$ to determine the next interval $[a_{n+1}, b_{n+1}]$.


Instead, we choose x_n to be the root of the interpolating polynomial $p(x)$ that interpolates the data $(a_n, f(a_n)), (b_n, f(b_n))$ (secant cuts the x -axis)

$$p(x) = \frac{(x - b_n)}{a_n - b_n} f(a_n) + \frac{(x - a_n)}{b_n - a_n} f(b_n).$$

Now $p(x) = 0$ yields

$$x = a_n - \frac{(a_n - b_n)}{f(a_n) - f(b_n)} f(a_n) = a_n - \frac{f(a_n)}{f[a_n, b_n]}.$$

Regula-Falsi method



Regula-Falsi method (cont.)

Thus we have the following iterative method:

- For $n = 1, 2, \dots$ do
- $$x_n = a_n - \frac{(a_n - b_n)}{f(a_n) - f(b_n)} f(a_n)$$

Regula-Falsi method (cont.)

Thus we have the following iterative method:

- For $n = 1, 2, \dots$ do
- $x_n = a_n - \frac{(a_n - b_n)}{f(a_n) - f(b_n)} f(a_n)$
- if $\text{sign}(f(a_n)) = \text{sign}(f(x_n))$ then

Regula-Falsi method (cont.)

Thus we have the following iterative method:

- For $n = 1, 2, \dots$ do
- $x_n = a_n - \frac{(a_n - b_n)}{f(a_n) - f(b_n)} f(a_n)$
- if $\text{sign}(f(a_n)) = \text{sign}(f(x_n))$ then
- $a_{n+1} = x_n, b_{n+1} = b_n$

Regula-Falsi method (cont.)

Thus we have the following iterative method:

- For $n = 1, 2, \dots$ do
- $x_n = a_n - \frac{(a_n - b_n)}{f(a_n) - f(b_n)} f(a_n)$
- if $\text{sign}(f(a_n)) = \text{sign}(f(x_n))$ then
- $a_{n+1} = x_n, b_{n+1} = b_n$
- else $a_{n+1} = a_n, b_{n+1} = x_n$.

Regula-Falsi method (cont.)

Thus we have the following iterative method:

- For $n = 1, 2, \dots$ do
- $x_n = a_n - \frac{(a_n - b_n)}{f(a_n) - f(b_n)} f(a_n)$
- if $\text{sign}(f(a_n)) = \text{sign}(f(x_n))$ then
- $a_{n+1} = x_n, b_{n+1} = b_n$
- else $a_{n+1} = a_n, b_{n+1} = x_n$.

Algorithm:

Input: $a, b \in \mathbb{R}$ such that $f(a)f(b) < 0$.

Output Approximate solution m of $f(x) = 0$.

Regula-Falsi method (cont.)

Thus we have the following iterative method:

- For $n = 1, 2, \dots$ do
- $x_n = a_n - \frac{(a_n - b_n)}{f(a_n) - f(b_n)} f(a_n)$
- if $\text{sign}(f(a_n)) = \text{sign}(f(x_n))$ then
- $a_{n+1} = x_n, b_{n+1} = b_n$
- else $a_{n+1} = a_n, b_{n+1} = x_n$.

Algorithm:

Input: $a, b \in \mathbb{R}$ such that $f(a)f(b) < 0$.

Output Approximate solution m of $f(x) = 0$.

- **while** $(b - a > \text{tol})$ **do**
- $m = a - (b - a)/(f(b)/f(a) - 1)$

Regula-Falsi method (cont.)

Thus we have the following iterative method:

- For $n = 1, 2, \dots$ do
- $x_n = a_n - \frac{(a_n - b_n)}{f(a_n) - f(b_n)} f(a_n)$
- if $\text{sign}(f(a_n)) = \text{sign}(f(x_n))$ then
- $a_{n+1} = x_n, b_{n+1} = b_n$
- else $a_{n+1} = a_n, b_{n+1} = x_n$.

Algorithm:

Input: $a, b \in \mathbb{R}$ such that $f(a)f(b) < 0$.

Output Approximate solution m of $f(x) = 0$.

- **while** $(b - a > \text{tol})$ **do**
- $m = a - (b - a)/(f(b)/f(a) - 1)$
- **if** $\text{sign}(f(a)) = \text{sign}(f(m))$ **then**

Regula-Falsi method (cont.)

Thus we have the following iterative method:

- For $n = 1, 2, \dots$ do
- $x_n = a_n - \frac{(a_n - b_n)}{f(a_n) - f(b_n)} f(a_n)$
- if $\text{sign}(f(a_n)) = \text{sign}(f(x_n))$ then
- $a_{n+1} = x_n, b_{n+1} = b_n$
- else $a_{n+1} = a_n, b_{n+1} = x_n$.

Algorithm:

Input: $a, b \in \mathbb{R}$ such that $f(a)f(b) < 0$.

Output Approximate solution m of $f(x) = 0$.

- **while** $(b - a > \text{tol})$ **do**
- $m = a - (b - a)/(f(b)/f(a) - 1)$
- **if** $\text{sign}(f(a)) = \text{sign}(f(m))$ **then**
- $a = m$ **else** $b = m$

Convergence of Regula-Falsi method

Under additional assumptions the Regula-Falsi Method converges linearly.
The rate of convergence may be faster or slower than Bisection Method.

Convergence of Regula-Falsi method

Under additional assumptions the Regula-Falsi Method converges linearly. The rate of convergence may be faster or slower than Bisection Method.

Convergence analysis is easy when f is convex or concave.

Convergence of Regula-Falsi method

Under additional assumptions the Regula-Falsi Method converges linearly. The rate of convergence may be faster or slower than Bisection Method.

Convergence analysis is easy when f is convex or concave. So assume that f is convex, say, $f''(x) > 0$ on $[a, b]$ with $f(a) < 0$ and $f(b) > 0$.

Convergence of Regula-Falsi method

Under additional assumptions the Regula-Falsi Method converges linearly. The rate of convergence may be faster or slower than Bisection Method.

Convergence analysis is easy when f is convex or concave. So assume that f is convex, say, $f''(x) > 0$ on $[a, b]$ with $f(a) < 0$ and $f(b) > 0$.

Then f has exactly one zero α in $[a, b]$. The secant joining $(a, f(a))$ and $(b, f(b))$ lies above the graph $y = f(x)$ and hence intersects x -axis to the left of α .

Convergence of Regula-Falsi method

Under additional assumptions the Regula-Falsi Method converges linearly. The rate of convergence may be faster or slower than Bisection Method.

Convergence analysis is easy when f is convex or concave. So assume that f is convex, say, $f''(x) > 0$ on $[a, b]$ with $f(a) < 0$ and $f(b) > 0$.

Then f has exactly one zero α in $[a, b]$. The secant joining $(a, f(a))$ and $(b, f(b))$ lies above the graph $y = f(x)$ and hence intersects x -axis to the left of α . This will be case for all subsequent secants.

Convergence of Regula-Falsi method

Under additional assumptions the Regula-Falsi Method converges linearly. The rate of convergence may be faster or slower than Bisection Method.

Convergence analysis is easy when f is convex or concave. So assume that f is convex, say, $f''(x) > 0$ on $[a, b]$ with $f(a) < 0$ and $f(b) > 0$.

Then f has exactly one zero α in $[a, b]$. The secant joining $(a, f(a))$ and $(b, f(b))$ lies above the graph $y = f(x)$ and hence intersects x -axis to the left of α . **This will be case for all subsequent secants.**

Hence b remains fixed while a gets updated, producing increasing sequence with $x_1 = a$ and

$$x_{n+1} = x_n - \frac{x_n - b}{f(x_n) - f(b)} f(x_n), \quad n = 1, 2, \dots$$

Convergence of Regula-Falsi method

Under additional assumptions the Regula-Falsi Method converges linearly. The rate of convergence may be faster or slower than Bisection Method.

Convergence analysis is easy when f is convex or concave. So assume that f is convex, say, $f''(x) > 0$ on $[a, b]$ with $f(a) < 0$ and $f(b) > 0$.

Then f has exactly one zero α in $[a, b]$. The secant joining $(a, f(a))$ and $(b, f(b))$ lies above the graph $y = f(x)$ and hence intersects x -axis to the left of α . **This will be case for all subsequent secants.**

Hence b remains fixed while a gets updated, producing increasing sequence with $x_1 = a$ and

$$x_{n+1} = x_n - \frac{x_n - b}{f(x_n) - f(b)} f(x_n), \quad n = 1, 2, \dots$$

Since (x_n) is increasing and bounded above by α and hence convergent, letting $n \rightarrow \infty$, we have $f(x_n) \rightarrow 0$ and hence $x_n \rightarrow \alpha$. (Why?)

Secant method




Figure : Secant method for finding the zero of $x^3 + x - 1$.

The secant method is a simple variant of regular-falsi method in which it is no longer required that f has opposite signs at the end points of each intervals generated.

Secant method




Figure : Secant method for finding the zero of $x^3 + x - 1$.

The secant method is a simple variant of regular-falsi method in which it is no longer required that f has opposite signs at the end points of each intervals generated.

One starts with two arbitrary $x_0 \neq x_1$ generates the iterates

$$x_{n+1} = x_n - \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})} f(x_n), \quad n = 1, 2, 3, \dots$$

Convergence secant method

The secant method converges only locally, that is, only if x_0 and x_1 are sufficiently close to a zero of f . Suppose that $f(\alpha) = 0$. Then

Convergence secant method

The secant method converges only locally, that is, only if x_0 and x_1 are sufficiently close to a zero of f . Suppose that $f(\alpha) = 0$. Then

$$x_{n+1} - \alpha = x_n - \alpha - \frac{f(x_n)}{f[x_{n-1}, x_n]}$$

Convergence secant method

The secant method converges only locally, that is, only if x_0 and x_1 are sufficiently close to a zero of f . Suppose that $f(\alpha) = 0$. Then

$$x_{n+1} - \alpha = x_n - \alpha - \frac{f(x_n)}{f[x_{n-1}, x_n]} = (x_n - \alpha) \left(1 - \frac{f(x_n) - f(\alpha)}{(x_n - \alpha)f[x_{n-1}, x_n]} \right)$$

Convergence secant method

The secant method converges only locally, that is, only if x_0 and x_1 are sufficiently close to a zero of f . Suppose that $f(\alpha) = 0$. Then

$$\begin{aligned}x_{n+1} - \alpha &= x_n - \alpha - \frac{f(x_n)}{f[x_{n-1}, x_n]} = (x_n - \alpha) \left(1 - \frac{f(x_n) - f(\alpha)}{(x_n - \alpha)f[x_{n-1}, x_n]} \right) \\&= (x_n - \alpha) \left(1 - \frac{f[x_n, \alpha]}{f[x_{n-1}, x_n]} \right)\end{aligned}$$

Convergence secant method

The secant method converges only locally, that is, only if x_0 and x_1 are sufficiently close to a zero of f . Suppose that $f(\alpha) = 0$. Then

$$\begin{aligned}x_{n+1} - \alpha &= x_n - \alpha - \frac{f(x_n)}{f[x_{n-1}, x_n]} = (x_n - \alpha) \left(1 - \frac{f(x_n) - f(\alpha)}{(x_n - \alpha)f[x_{n-1}, x_n]} \right) \\&= (x_n - \alpha) \left(1 - \frac{f[x_n, \alpha]}{f[x_{n-1}, x_n]} \right) = (x_n - \alpha) \left(\frac{f[x_{n-1}, x_n] - f[x_n, \alpha]}{f[x_{n-1}, x_n]} \right)\end{aligned}$$

Convergence secant method

The secant method converges only locally, that is, only if x_0 and x_1 are sufficiently close to a zero of f . Suppose that $f(\alpha) = 0$. Then

$$\begin{aligned}x_{n+1} - \alpha &= x_n - \alpha - \frac{f(x_n)}{f[x_{n-1}, x_n]} = (x_n - \alpha) \left(1 - \frac{f(x_n) - f(\alpha)}{(x_n - \alpha)f[x_{n-1}, x_n]} \right) \\&= (x_n - \alpha) \left(1 - \frac{f[x_n, \alpha]}{f[x_{n-1}, x_n]} \right) = (x_n - \alpha) \left(\frac{f[x_{n-1}, x_n] - f[x_n, \alpha]}{f[x_{n-1}, x_n]} \right) \\&= (x_n - \alpha)(x_{n-1} - \alpha) \frac{f[x_{n-1}, x_n, \alpha]}{f[x_{n-1}, x_n]}.\end{aligned}$$

If α is a simple zero, that is, $f'(\alpha) \neq 0$, and if $x_n \rightarrow \alpha$ then

$$\frac{f[x_{n-1}, x_n, \alpha]}{f[x_{n-1}, x_n]} \longrightarrow \frac{f''(\alpha)}{2f'(\alpha)} \text{ as } n \rightarrow \infty$$

when f is twice continuously differentiable near α .

Convergence secant method

The secant method converges only locally, that is, only if x_0 and x_1 are sufficiently close to a zero of f . Suppose that $f(\alpha) = 0$. Then

$$\begin{aligned}x_{n+1} - \alpha &= x_n - \alpha - \frac{f(x_n)}{f[x_{n-1}, x_n]} = (x_n - \alpha) \left(1 - \frac{f(x_n) - f(\alpha)}{(x_n - \alpha)f[x_{n-1}, x_n]} \right) \\&= (x_n - \alpha) \left(1 - \frac{f[x_n, \alpha]}{f[x_{n-1}, x_n]} \right) = (x_n - \alpha) \left(\frac{f[x_{n-1}, x_n] - f[x_n, \alpha]}{f[x_{n-1}, x_n]} \right) \\&= (x_n - \alpha)(x_{n-1} - \alpha) \frac{f[x_{n-1}, x_n, \alpha]}{f[x_{n-1}, x_n]}.\end{aligned}$$

If α is a simple zero, that is, $f'(\alpha) \neq 0$, and if $x_n \rightarrow \alpha$ then

$$\frac{f[x_{n-1}, x_n, \alpha]}{f[x_{n-1}, x_n]} \longrightarrow \frac{f''(\alpha)}{2f'(\alpha)} \text{ as } n \rightarrow \infty$$

when f is twice continuously differentiable near α .

Then we have $\lim_{n \rightarrow \infty} \frac{x_{n+1} - \alpha}{x_n - \alpha} = 0$ showing that the convergence is better

than linear.

Convergence secant method

The secant method converges only locally, that is, only if x_0 and x_1 are sufficiently close to a zero of f . Suppose that $f(\alpha) = 0$. Then

$$\begin{aligned}x_{n+1} - \alpha &= x_n - \alpha - \frac{f(x_n)}{f[x_{n-1}, x_n]} = (x_n - \alpha) \left(1 - \frac{f(x_n) - f(\alpha)}{(x_n - \alpha)f[x_{n-1}, x_n]} \right) \\&= (x_n - \alpha) \left(1 - \frac{f[x_n, \alpha]}{f[x_{n-1}, x_n]} \right) = (x_n - \alpha) \left(\frac{f[x_{n-1}, x_n] - f[x_n, \alpha]}{f[x_{n-1}, x_n]} \right) \\&= (x_n - \alpha)(x_{n-1} - \alpha) \frac{f[x_{n-1}, x_n, \alpha]}{f[x_{n-1}, x_n]}.\end{aligned}$$

If α is a simple zero, that is, $f'(\alpha) \neq 0$, and if $x_n \rightarrow \alpha$ then

$$\frac{f[x_{n-1}, x_n, \alpha]}{f[x_{n-1}, x_n]} \longrightarrow \frac{f''(\alpha)}{2f'(\alpha)} \text{ as } n \rightarrow \infty$$

when f is twice continuously differentiable near α .

Then we have $\lim_{n \rightarrow \infty} \frac{x_{n+1} - \alpha}{x_n - \alpha} = 0$ showing that the convergence is better

than linear. The order of convergence is $p := \frac{1 + \sqrt{5}}{2}$.

Example

Consider $f(x) = \cos(x) \cosh(x) - 1$ with $a = 3\pi/2$ and $b = 2\pi$. Then f is convex on $[a, b]$ as $f''(x) = -2 \sin(x) \sinh(x) > 0$ for $x \in [a, b]$.

Example

Consider $f(x) = \cos(x) \cosh(x) - 1$ with $a = 3\pi/2$ and $b = 2\pi$. Then f is convex on $[a, b]$ as $f''(x) = -2 \sin(x) \sinh(x) > 0$ for $x \in [a, b]$.

Note that $f(a) = -1$ and $f(b) = \cosh(2\pi) - 1 > 0$. The fact that one of the false positions (in this case b) is fixed may go against Regula-Falsi method.

Example

Consider $f(x) = \cos(x) \cosh(x) - 1$ with $a = 3\pi/2$ and $b = 2\pi$. Then f is convex on $[a, b]$ as $f''(x) = -2 \sin(x) \sinh(x) > 0$ for $x \in [a, b]$.

Note that $f(a) = -1$ and $f(b) = \cosh(2\pi) - 1 > 0$. The fact that one of the false positions (in this case b) is fixed may go against Regula-Falsi method.

For $\text{tol} = 0.5 \times 10^{-7}, 0.5 \times 10^{-15}$, Regular-Falsi takes 42 and 87 iterations as compared to 25 and 52 by bisection method.

Example

Consider $f(x) = \cos(x) \cosh(x) - 1$ with $a = 3\pi/2$ and $b = 2\pi$. Then f is convex on $[a, b]$ as $f''(x) = -2\sin(x) \sinh(x) > 0$ for $x \in [a, b]$.

Note that $f(a) = -1$ and $f(b) = \cosh(2\pi) - 1 > 0$. The fact that one of the false positions (in this case b) is fixed may go against Regula-Falsi method.

For $\text{tol} = 0.5 \times 10^{-7}, 0.5 \times 10^{-15}$, Regular-Falsi takes 42 and 87 iterations as compared to 25 and 52 by bisection method.

By contrast, secant method converges faster than both Bisection and Regula-Falsi methods. The rate of convergence of Regula-Falsi method is the slowest among all three methods.

Convergence of Regula-Falsi and Secant Methods

Both Secant and Regula-Falsi Methods can converge very slowly.




Figure : Slow convergence of (a) Secant Method and (b) Regula-Falsi Method to the root 0 of $x^3 - 2x^2 + \frac{3}{2}x$.