

MA579H Scientific Computing

Numerical Integration I

Rafikul Alam
Department of Mathematics
IIT Guwahati

Lecture outline

- Quadrature formula
- Newton-Cotes formula
- Midpoint and composite midpoint rules
- Trapezoid and composite trapezoid rules
- Simpson's and composite Simpson's rule

Quadrature

Integrals such as

$$\int_0^2 e^{-x^2} dx, \int_0^\pi \cos(3 \sin(\log(1+x))) dx, \int_0^1 x^2 e^{\tan(\sin(x))} dx$$

are not amenable to exact evaluation. The only option is to approximate such integrals numerically.

Quadrature

Integrals such as

$$\int_0^2 e^{-x^2} dx, \quad \int_0^\pi \cos(3 \sin(\log(1+x))) dx, \quad \int_0^1 x^2 e^{\tan(\sin(x))} dx$$

are not amenable to exact evaluation. The only option is to approximate such integrals numerically.

Problem: Let $f \in C[a, b]$. Compute the integral $I(f) := \int_a^b f(x) dx$.

Quadrature

Integrals such as

$$\int_0^2 e^{-x^2} dx, \int_0^\pi \cos(3 \sin(\log(1+x))) dx, \int_0^1 x^2 e^{\tan(\sin(x))} dx$$

are not amenable to exact evaluation. The only option is to approximate such integrals numerically.

Problem: Let $f \in C[a, b]$. Compute the integral $I(f) := \int_a^b f(x) dx$.

Strategy: Integration via polynomial interpolation.

- Approximate f using an **interpolating polynomial** $p_n(x)$.
- Evaluate $Q_n(f) := \int_a^b p_n(x) dx$, since it is easy to evaluate integral of a polynomial. We expect that $Q_n(f) \approx I(f)$.

Quadrature

Consider the nodes $[x_0, \dots, x_n]$ in $[a, b]$ and the Lagrange interpolating polynomial $p_n(x)$ given by $p_n(x) := f(x_0)\ell_0(x) + \dots + f(x_n)\ell_n(x)$, where

Quadrature

Consider the nodes $[x_0, \dots, x_n]$ in $[a, b]$ and the Lagrange interpolating polynomial $p_n(x)$ given by $p_n(x) := f(x_0)\ell_0(x) + \dots + f(x_n)\ell_n(x)$, where

$$\ell_j(x) := \frac{w(x)}{(x - x_j)w'(x_j)} = \prod_{i \neq j} \frac{(x - x_i)}{(x_j - x_i)}, \quad j = 0 : n,$$

and $w(x) := (x - x_0) \cdots (x - x_n)$. Then

Quadrature

Consider the nodes $[x_0, \dots, x_n]$ in $[a, b]$ and the Lagrange interpolating polynomial $p_n(x)$ given by $p_n(x) := f(x_0)\ell_0(x) + \dots + f(x_n)\ell_n(x)$, where

$$\ell_j(x) := \frac{w(x)}{(x - x_j)w'(x_j)} = \prod_{i \neq j} \frac{(x - x_i)}{(x_j - x_i)}, \quad j = 0 : n,$$

and $w(x) := (x - x_0) \cdots (x - x_n)$. Then

$$Q_n(f) = \int_a^b p_n(x) dx = \sum_{j=0}^n f(x_j) \int_a^b \ell_j(x) dx = \sum_{j=0}^n w_j f(x_j),$$

where $w_j := \int_a^b \ell_j(x) dx$ is called the j -th quadrature weight and

$$Q_n(f) := \sum_{j=0}^n w_j f(x_j)$$

is called a quadrature formula or a quadrature rule.

Quadrature

Consider the nodes $[x_0, \dots, x_n]$ in $[a, b]$ and the Lagrange interpolating polynomial $p_n(x)$ given by $p_n(x) := f(x_0)\ell_0(x) + \dots + f(x_n)\ell_n(x)$, where

$$\ell_j(x) := \frac{w(x)}{(x - x_j)w'(x_j)} = \prod_{i \neq j} \frac{(x - x_i)}{(x_j - x_i)}, \quad j = 0 : n,$$

and $w(x) := (x - x_0) \cdots (x - x_n)$. Then

$$Q_n(f) = \int_a^b p_n(x) dx = \sum_{j=0}^n f(x_j) \int_a^b \ell_j(x) dx = \sum_{j=0}^n w_j f(x_j),$$

where $w_j := \int_a^b \ell_j(x) dx$ is called the j -th quadrature weight and

$$Q_n(f) := \sum_{j=0}^n w_j f(x_j)$$

is called a quadrature formula or a quadrature rule.

Note that the quadrature weights do not depend on the function $f(x)$ and hence can be computed independently of $f(x)$ and stored.

Error estimate

Set $E_n(f) := I(f) - Q_n(f)$. If $f \in C^{n+1}[a, b]$ then

$$\begin{aligned}|E_n(f)| &\leq \int_a^b |f(x) - p_n(x)| dx = \frac{1}{(n+1)!} \int_a^b |f^{(n+1)}(\xi_x) w(x)| dx \\ &\leq \frac{\|f^{(n+1)}\|_\infty}{(n+1)!} \int_a^b |(x - x_0)(x - x_1) \cdots (x - x_n)| dx.\end{aligned}$$

Error estimate

Set $E_n(f) := I(f) - Q_n(f)$. If $f \in C^{n+1}[a, b]$ then

$$\begin{aligned}|E_n(f)| &\leq \int_a^b |f(x) - p_n(x)| dx = \frac{1}{(n+1)!} \int_a^b |f^{(n+1)}(\xi_x) w(x)| dx \\ &\leq \frac{\|f^{(n+1)}\|_\infty}{(n+1)!} \int_a^b |(x - x_0)(x - x_1) \cdots (x - x_n)| dx.\end{aligned}$$

This shows that for Chebyshev nodes $E_n(f) \leq \left(\frac{b-a}{2}\right)^{n+2} \frac{\|f^{(n+1)}\|_\infty}{2^{n-1}(n+1)!}$.

Error estimate

Set $E_n(f) := I(f) - Q_n(f)$. If $f \in C^{n+1}[a, b]$ then

$$\begin{aligned}|E_n(f)| &\leq \int_a^b |f(x) - p_n(x)| dx = \frac{1}{(n+1)!} \int_a^b |f^{(n+1)}(\xi_x) w(x)| dx \\ &\leq \frac{\|f^{(n+1)}\|_\infty}{(n+1)!} \int_a^b |(x - x_0)(x - x_1) \cdots (x - x_n)| dx.\end{aligned}$$

This shows that for Chebyshev nodes $E_n(f) \leq \left(\frac{b-a}{2}\right)^{n+2} \frac{\|f^{(n+1)}\|_\infty}{2^{n-1}(n+1)!}$.

Note that $E_n(f) = 0$, that is, $Q_n(f) = I(f)$ when $f \in \mathcal{P}_n$.

Error estimate

Set $E_n(f) := I(f) - Q_n(f)$. If $f \in C^{n+1}[a, b]$ then

$$\begin{aligned}|E_n(f)| &\leq \int_a^b |f(x) - p_n(x)| dx = \frac{1}{(n+1)!} \int_a^b |f^{(n+1)}(\xi_x) w(x)| dx \\ &\leq \frac{\|f^{(n+1)}\|_\infty}{(n+1)!} \int_a^b |(x - x_0)(x - x_1) \cdots (x - x_n)| dx.\end{aligned}$$

This shows that for Chebyshev nodes $E_n(f) \leq \left(\frac{b-a}{2}\right)^{n+2} \frac{\|f^{(n+1)}\|_\infty}{2^{n-1}(n+1)!}$.

Note that $E_n(f) = 0$, that is, $Q_n(f) = I(f)$ when $f \in \mathcal{P}_n$.

Exactness: A quadrature rule $Q_n(f)$ is said to have **degree of exactness m** if $Q_n(f) = I(f)$ for all $f \in \mathcal{P}_m$.

Error estimate

Set $E_n(f) := I(f) - Q_n(f)$. If $f \in C^{n+1}[a, b]$ then

$$\begin{aligned}|E_n(f)| &\leq \int_a^b |f(x) - p_n(x)| dx = \frac{1}{(n+1)!} \int_a^b |f^{(n+1)}(\xi_x) w(x)| dx \\ &\leq \frac{\|f^{(n+1)}\|_\infty}{(n+1)!} \int_a^b |(x - x_0)(x - x_1) \cdots (x - x_n)| dx.\end{aligned}$$

This shows that for Chebyshev nodes $E_n(f) \leq \left(\frac{b-a}{2}\right)^{n+2} \frac{\|f^{(n+1)}\|_\infty}{2^{n-1}(n+1)!}$.

Note that $E_n(f) = 0$, that is, $Q_n(f) = I(f)$ when $f \in \mathcal{P}_n$.

Exactness: A quadrature rule $Q_n(f)$ is said to have **degree of exactness m** if $Q_n(f) = I(f)$ for all $f \in \mathcal{P}_m$.

The quadrature rule $Q_n(f) = \int_a^b p_n(x) dx$ has degree of exactness $\geq n$.

Newton-Cotes quadrature

A quadrature rule based on an interpolating polynomial $p_n(x)$ at $n + 1$ equally spaced nodes in $[a, b]$ is called **Newton-Cotes formula** of order n .

Newton-Cotes quadrature

A quadrature rule based on an interpolating polynomial $p_n(x)$ at $n + 1$ equally spaced nodes in $[a, b]$ is called **Newton-Cotes formula** of order n .

Consider the equally spaced nodes $[x_0, \dots, x_n]$ and the Lagrange interpolant $p_n(x)$, where $x_j = a + jh$, $j = 0 : n$, and $h := (b - a)/n$.

Newton-Cotes quadrature

A quadrature rule based on an interpolating polynomial $p_n(x)$ at $n + 1$ equally spaced nodes in $[a, b]$ is called **Newton-Cotes formula** of order n .

Consider the equally spaced nodes $[x_0, \dots, x_n]$ and the Lagrange interpolant $p_n(x)$, where $x_j = a + jh$, $j = 0 : n$, and $h := (b - a)/n$.

Since for equally spaced nodes $w(x) := (x - x_0)(x - x_1) \cdots (x - x_n)$ oscillates near the end nodes, it follows from

Newton-Cotes quadrature

A quadrature rule based on an interpolating polynomial $p_n(x)$ at $n + 1$ equally spaced nodes in $[a, b]$ is called **Newton-Cotes formula** of order n .

Consider the equally spaced nodes $[x_0, \dots, x_n]$ and the Lagrange interpolant $p_n(x)$, where $x_j = a + jh$, $j = 0 : n$, and $h := (b - a)/n$.

Since for equally spaced nodes $w(x) := (x - x_0)(x - x_1) \cdots (x - x_n)$ oscillates near the end nodes, it follows from

$$|E_n(f)| \leq \frac{\|f^{(n+1)}\|_\infty}{(n+1)!} \int_a^b |(x - x_0)(x - x_1) \cdots (x - x_n)| dx$$

that $Q_n(f)$ is susceptible to **Runge's phenomenon**.

Newton-Cotes quadrature

A quadrature rule based on an interpolating polynomial $p_n(x)$ at $n + 1$ equally spaced nodes in $[a, b]$ is called **Newton-Cotes formula** of order n .

Consider the equally spaced nodes $[x_0, \dots, x_n]$ and the Lagrange interpolant $p_n(x)$, where $x_j = a + jh$, $j = 0 : n$, and $h := (b - a)/n$.

Since for equally spaced nodes $w(x) := (x - x_0)(x - x_1) \cdots (x - x_n)$ oscillates near the end nodes, it follows from

$$|E_n(f)| \leq \frac{\|f^{(n+1)}\|_\infty}{(n+1)!} \int_a^b |(x - x_0)(x - x_1) \cdots (x - x_n)| dx$$

that $Q_n(f)$ is susceptible to **Runge's phenomenon**.

Newton-Cotes is not useful for large n . There are two natural ways to ensure $Q_n(f) \rightarrow I(f)$ as $n \rightarrow \infty$:

- Don't use equally spaced nodes.
- Integrate piecewise polynomial interpolant.

Midpoint rule

The quadrature corresponding to $p_0(x) = f((a+b)/2)$ is called the **midpoint rule** and is given by

$$M(f) = (b-a)f\left(\frac{a+b}{2}\right).$$

Midpoint rule

The quadrature corresponding to $p_0(x) = f((a+b)/2)$ is called the **midpoint rule** and is given by

$$M(f) = (b-a)f\left(\frac{a+b}{2}\right).$$

This corresponds to a Newton-Cotes formula of order $n = 0$ and hence is expected to be exact for polynomials of degree 0.

Midpoint rule

The quadrature corresponding to $p_0(x) = f((a+b)/2)$ is called the **midpoint rule** and is given by

$$M(f) = (b-a)f\left(\frac{a+b}{2}\right).$$

This corresponds to a Newton-Cotes formula of order $n = 0$ and hence is expected to be exact for polynomials of degree 0.

Note, however, that

$$M(x) = \frac{b^2 - a^2}{2} = \int_a^b x dx,$$

which shows that it is exact for polynomials of degree 1. Thus, the midpoint rule is **exact for polynomials of degree 1**.

Midpoint rule

Theorem: If $f \in C^2[a, b]$ then there exists $\theta \in [a, b]$ such that

$$\int_a^b f(x) dx = M(f) + \frac{f''(\theta)}{24}(b-a)^3.$$

Hence $E(f) := I(f) - M(f) = \frac{f''(\theta)}{24}(b-a)^3.$

Midpoint rule

Theorem: If $f \in C^2[a, b]$ then there exists $\theta \in [a, b]$ such that

$$\int_a^b f(x)dx = M(f) + \frac{f''(\theta)}{24}(b-a)^3.$$

Hence $E(f) := I(f) - M(f) = \frac{f''(\theta)}{24}(b-a)^3$.

Proof: Set $w := (a+b)/2$. Then by Taylor's theorem

$$f(x) = f(w) + f'(w)(x-w) + f''(\theta_x)(x-w)^2/2.$$

Upon integration, and using MVT for integral, we have

Midpoint rule

Theorem: If $f \in C^2[a, b]$ then there exists $\theta \in [a, b]$ such that

$$\int_a^b f(x)dx = M(f) + \frac{f''(\theta)}{24}(b-a)^3.$$

Hence $E(f) := I(f) - M(f) = \frac{f''(\theta)}{24}(b-a)^3$.

Proof: Set $w := (a+b)/2$. Then by Taylor's theorem

$$f(x) = f(w) + f'(w)(x-w) + f''(\theta_x)(x-w)^2/2.$$

Upon integration, and using MVT for integral, we have

$$\begin{aligned}\int_a^b f(x)dx &= (b-a)f(w) + 0 + \frac{f''(\theta)}{2} \int_a^b (x-w)^2 dx \\ &= M(f) + \frac{f''(\theta)}{24}(b-a)^3.\end{aligned}$$

Composite midpoint rule

Consider the equally spaced nodes $[x_0, \dots, x_n]$, where $x_j = a + jh$, $j = 0 : n$, and $h := (b - a)/n$. Set $w_j := (x_j + x_{j+1})/2$ for $j = 0 : n - 1$. Then

$$I(f) = \int_a^b f(x) dx = \sum_{j=1}^n \int_{x_{j-1}}^{x_j} f(x) dx.$$

Composite midpoint rule

Consider the equally spaced nodes $[x_0, \dots, x_n]$, where $x_j = a + jh$, $j = 0 : n$, and $h := (b - a)/n$. Set $w_j := (x_j + x_{j+1})/2$ for $j = 0 : n - 1$. Then

$$I(f) = \int_a^b f(x)dx = \sum_{j=1}^n \int_{x_{j-1}}^{x_j} f(x)dx.$$

By midpoint rule, $\int_{x_{j-1}}^{x_j} f(x)dx \approx hf(w_{j-1})$. Hence

$$I(f) = \int_a^b f(x)dx = \sum_{j=1}^n \int_{x_{j-1}}^{x_j} f(x)dx \approx \sum_{j=1}^n hf(w_{j-1}).$$

Composite midpoint rule

Consider the equally spaced nodes $[x_0, \dots, x_n]$, where $x_j = a + jh$, $j = 0 : n$, and $h := (b - a)/n$. Set $w_j := (x_j + x_{j+1})/2$ for $j = 0 : n - 1$. Then

$$I(f) = \int_a^b f(x)dx = \sum_{j=1}^n \int_{x_{j-1}}^{x_j} f(x)dx.$$

By midpoint rule, $\int_{x_{j-1}}^{x_j} f(x)dx \approx hf(w_{j-1})$. Hence

$$I(f) = \int_a^b f(x)dx = \sum_{j=1}^n \int_{x_{j-1}}^{x_j} f(x)dx \approx \sum_{j=1}^n hf(w_{j-1}).$$

This yields the **composite midpoint rule**

$$M_n(f) := \sum_{j=1}^n hf(w_{j-1})$$

Composite midpoint rule

Consider the equally spaced nodes $[x_0, \dots, x_n]$, where $x_j = a + jh$, $j = 0 : n$, and $h := (b - a)/n$. Set $w_j := (x_j + x_{j+1})/2$ for $j = 0 : n - 1$. Then

$$I(f) = \int_a^b f(x) dx = \sum_{j=1}^n \int_{x_{j-1}}^{x_j} f(x) dx.$$

By midpoint rule, $\int_{x_{j-1}}^{x_j} f(x) dx \approx hf(w_{j-1})$. Hence

$$I(f) = \int_a^b f(x) dx = \sum_{j=1}^n \int_{x_{j-1}}^{x_j} f(x) dx \approx \sum_{j=1}^n hf(w_{j-1}).$$

This yields the **composite midpoint rule**

$$\begin{aligned} M_n(f) &:= \sum_{j=1}^n hf(w_{j-1}) \\ &= h[f(w_0) + f(w_2) + \dots + f(w_{n-1})]. \end{aligned}$$

Error in composite midpoint rule

By midpoint rule, $\int_{x_{j-1}}^{x_j} f(x) dx - hf(w_{j-1}) = \frac{f''(\theta_j)}{24} h^3.$

Error in composite midpoint rule

By midpoint rule, $\int_{x_{j-1}}^{x_j} f(x) dx - hf(w_{j-1}) = \frac{f''(\theta_j)}{24} h^3$.

Consequently, there exists $\theta \in [a, b]$ such that

$$\begin{aligned} E_n(f) &:= I(f) - M_n(f) = \sum_{j=1}^n \frac{f''(\theta_j)}{24} h^3 \\ &= n \frac{f''(\theta)}{24} h^3 = \frac{h^2}{24} (b-a) f''(\theta). \end{aligned}$$

Error in composite midpoint rule

By midpoint rule, $\int_{x_{j-1}}^{x_j} f(x)dx - hf(w_{j-1}) = \frac{f''(\theta_j)}{24}h^3$.

Consequently, there exists $\theta \in [a, b]$ such that

$$\begin{aligned} E_n(f) &:= I(f) - M_n(f) = \sum_{j=1}^n \frac{f''(\theta_j)}{24}h^3 \\ &= n \frac{f''(\theta)}{24}h^3 = \frac{h^2}{24}(b-a)f''(\theta). \end{aligned}$$

Hence we have $|E_n(f)| \leq \frac{h^2}{24}(b-a)\|f''\|_\infty$.

Trapezoid rule

The quadrature $Q_n(f)$ is called Trapezoid rule when $n = 1$. Then nodes are $x_0 = a$ and $x_1 = b$, and $\ell_0(x) = \frac{x-b}{a-b}$ and $\ell_1(x) = \frac{x-a}{b-a}$.

Trapezoid rule

The quadrature $Q_n(f)$ is called Trapezoid rule when $n = 1$. Then nodes are $x_0 = a$ and $x_1 = b$, and $\ell_0(x) = \frac{x-b}{a-b}$ and $\ell_1(x) = \frac{x-a}{b-a}$.

The weights are given by

$$w_0 = \int_a^b \ell_0(x) dx = \frac{b-a}{2} = \int_a^b \ell_1(x) dx = w_1.$$

Trapezoid rule

The quadrature $Q_n(f)$ is called Trapezoid rule when $n = 1$. Then nodes are $x_0 = a$ and $x_1 = b$, and $\ell_0(x) = \frac{x-b}{a-b}$ and $\ell_1(x) = \frac{x-a}{b-a}$.

The weights are given by

$$w_0 = \int_a^b \ell_0(x) dx = \frac{b-a}{2} = \int_a^b \ell_1(x) dx = w_1.$$

Hence the Trapezoid rule is given by

$$T(f) = \frac{(b-a)}{2} [f(a) + f(b)].$$

Trapezoid rule

Theorem: If $f \in C^2[a, b]$ then there exists $\theta \in [a, b]$ such that

$$\int_a^b f(x)dx = T(f) - \frac{f''(\theta)}{12}(b-a)^3.$$

Hence $E(f) := I(f) - T(f) = -\frac{f''(\theta)}{12}(b-a)^3.$

Trapezoid rule

Theorem: If $f \in C^2[a, b]$ then there exists $\theta \in [a, b]$ such that

$$\int_a^b f(x)dx = T(f) - \frac{f''(\theta)}{12}(b-a)^3.$$

Hence $E(f) := I(f) - T(f) = -\frac{f''(\theta)}{12}(b-a)^3$.

Proof: By mean value theorem (MVT) of integral, we have

$$\begin{aligned} E(f) &= \int_a^b (f(x) - p(x))dx \\ &= \int_a^b \frac{f''(\xi_x)}{2}(x-a)(x-b)dx \\ &= \frac{f''(\theta)}{2} \int_a^b (x-a)(x-b)dx \quad \text{for some } \theta \in (a, b) \\ &= -\frac{f''(\theta)}{12}(b-a)^3. \end{aligned}$$

Composite trapezoid rule

Consider the equally spaced nodes $[x_0, \dots, x_n]$, where $x_j = a + jh$, $j = 0 : n$, and $h := (b - a)/n$.

Composite trapezoid rule

Consider the equally spaced nodes $[x_0, \dots, x_n]$, where $x_j = a + jh$, $j = 0 : n$, and $h := (b - a)/n$. Then

$$I(f) = \int_a^b f(x) dx = \sum_{j=1}^n \int_{x_{j-1}}^{x_j} f(x) dx.$$

Composite trapezoid rule

Consider the equally spaced nodes $[x_0, \dots, x_n]$, where $x_j = a + jh$, $j = 0 : n$, and $h := (b - a)/n$. Then

$$I(f) = \int_a^b f(x) dx = \sum_{j=1}^n \int_{x_{j-1}}^{x_j} f(x) dx.$$

By trapezoid rule, $\int_{x_{j-1}}^{x_j} f(x) dx \approx \frac{h}{2}[f(x_{j-1}) + f(x_j)]$. Hence

$$I(f) = \int_a^b f(x) dx = \sum_{j=1}^n \int_{x_{j-1}}^{x_j} f(x) dx \approx \sum_{j=1}^n \frac{h}{2}[f(x_{j-1}) + f(x_j)].$$

Composite trapezoid rule

Consider the equally spaced nodes $[x_0, \dots, x_n]$, where $x_j = a + jh$, $j = 0 : n$, and $h := (b - a)/n$. Then

$$I(f) = \int_a^b f(x) dx = \sum_{j=1}^n \int_{x_{j-1}}^{x_j} f(x) dx.$$

By trapezoid rule, $\int_{x_{j-1}}^{x_j} f(x) dx \approx \frac{h}{2}[f(x_{j-1}) + f(x_j)]$. Hence

$$I(f) = \int_a^b f(x) dx = \sum_{j=1}^n \int_{x_{j-1}}^{x_j} f(x) dx \approx \sum_{j=1}^n \frac{h}{2}[f(x_{j-1}) + f(x_j)].$$

This yields the **composite trapezoid rule**

$$\begin{aligned} T_n(f) &:= \sum_{j=1}^n \frac{h}{2}[f(x_{j-1}) + f(x_j)] \\ &= h \left[\frac{f(x_0)}{2} + f(x_1) + \dots + f(x_{n-1}) + \frac{f(x_n)}{2} \right]. \end{aligned}$$

Error in composite trapezoid rule

By trapezoid rule,
$$\int_{x_{j-1}}^{x_j} f(x) dx - \frac{h}{2}[f(x_{j-1}) + f(x_j)] = -\frac{f''(\theta_j)}{12} h^3.$$

Error in composite trapezoid rule

By trapezoid rule, $\int_{x_{j-1}}^{x_j} f(x)dx - \frac{h}{2}[f(x_{j-1}) + f(x_j)] = -\frac{f''(\theta_j)}{12}h^3.$

By intermediate value theorem (IVT), there exists $\theta \in [a, b]$ such that

$$\sum_{j=1}^n f''(\theta_j) = nf''(\theta).$$

Error in composite trapezoid rule

By trapezoid rule, $\int_{x_{j-1}}^{x_j} f(x)dx - \frac{h}{2}[f(x_{j-1}) + f(x_j)] = -\frac{f''(\theta_j)}{12}h^3$.

By intermediate value theorem (IVT), there exists $\theta \in [a, b]$ such that

$$\sum_{j=1}^n f''(\theta_j) = nf''(\theta).$$

Consequently, we have

$$\begin{aligned} E_n(f) &:= I(f) - T_n(f) = -\sum_{j=1}^n \frac{f''(\theta_j)}{12}h^3 \\ &= -n\frac{f''(\theta)}{12}h^3 = -\frac{h^2}{12}(b-a)f''(\theta). \end{aligned}$$

Error in composite trapezoid rule

By trapezoid rule, $\int_{x_{j-1}}^{x_j} f(x)dx - \frac{h}{2}[f(x_{j-1}) + f(x_j)] = -\frac{f''(\theta_j)}{12}h^3$.

By intermediate value theorem (IVT), there exists $\theta \in [a, b]$ such that

$$\sum_{j=1}^n f''(\theta_j) = nf''(\theta).$$

Consequently, we have

$$\begin{aligned} E_n(f) &:= I(f) - T_n(f) = -\sum_{j=1}^n \frac{f''(\theta_j)}{12}h^3 \\ &= -n\frac{f''(\theta)}{12}h^3 = -\frac{h^2}{12}(b-a)f''(\theta). \end{aligned}$$

Hence $|E_n(f)| \leq \frac{h^2}{12}(b-a)\|f''\|_{\infty}$.

Example

Consider the integral

$$I(f) := \int_0^1 e^{-x^2} dx.$$

The two Newton-Cotes (midpoint and trapezoid) formula yield

$$M(f) = (1 - 0)e^{-0.25} \approx 0.778801$$

$$T(f) = \frac{1}{2}[e^0 + e^{-1}] \approx 0.683940.$$

The correctly rounded value is $I = 0.746824$.

Example

Consider the integral

$$I(f) := \int_0^1 e^{-x^2} dx.$$

The two Newton-Cotes (midpoint and trapezoid) formula yield

$$M(f) = (1 - 0)e^{-0.25} \approx 0.778801$$

$$T(f) = \frac{1}{2}[e^0 + e^{-1}] \approx 0.683940.$$

The correctly rounded value is $I = 0.746824$. Note that

$$|I(f) - M(f)| = 0.031977 \text{ and } |I(f) - T(f)| = 0.062884.$$

Example

Consider the integral

$$I(f) := \int_0^1 e^{-x^2} dx.$$

The two Newton-Cotes (midpoint and trapezoid) formula yield

$$M(f) = (1 - 0)e^{-0.25} \approx 0.778801$$

$$T(f) = \frac{1}{2}[e^0 + e^{-1}] \approx 0.683940.$$

The correctly rounded value is $I = 0.746824$. Note that

$$|I(f) - M(f)| = 0.031977 \text{ and } |I(f) - T(f)| = 0.062884.$$

The error from the trapezoid rule is about twice that of the midpoint rule, which validates the error estimate.

Simpson's rule

The Newton-Cotes quadrature $Q_n(f)$ is called **Simpson's rule** when $n = 2$. The nodes are $x_0 = a$, $x_1 = (a + b)/2$ and $x_2 = b$.

Simpson's rule

The Newton-Cotes quadrature $Q_n(f)$ is called **Simpson's rule** when $n = 2$. The nodes are $x_0 = a$, $x_1 = (a + b)/2$ and $x_2 = b$. Then the **Simpson's rule** is given by

$$S(f) = \left[w_0 f(a) + w_1 f\left(\frac{a+b}{2}\right) + w_2 f(b) \right] \approx \int_a^b f(x) dx,$$

where $w_i := \int_a^b \ell_i(x) dx$, $i = 0, 1, 2$.

Simpson's rule

The Newton-Cotes quadrature $Q_n(f)$ is called **Simpson's rule** when $n = 2$. The nodes are $x_0 = a$, $x_1 = (a + b)/2$ and $x_2 = b$. Then the **Simpson's rule** is given by

$$S(f) = \left[w_0 f(a) + w_1 f\left(\frac{a+b}{2}\right) + w_2 f(b) \right] \approx \int_a^b f(x) dx,$$

where $w_i := \int_a^b \ell_i(x) dx$, $i = 0, 1, 2$.

A messy calculation shows that $w_0 = w_2 = (b - a)/6$ and $w_1 = 2(b - a)/3$. Hence

$$S(f) = \frac{(b - a)}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right].$$

Simpson's rule

The Newton-Cotes quadrature $Q_n(f)$ is called **Simpson's rule** when $n = 2$. The nodes are $x_0 = a$, $x_1 = (a + b)/2$ and $x_2 = b$. Then the **Simpson's rule** is given by

$$S(f) = \left[w_0 f(a) + w_1 f\left(\frac{a+b}{2}\right) + w_2 f(b) \right] \approx \int_a^b f(x) dx,$$

where $w_i := \int_a^b \ell_i(x) dx$, $i = 0, 1, 2$.

A messy calculation shows that $w_0 = w_2 = (b - a)/6$ and $w_1 = 2(b - a)/3$. Hence

$$S(f) = \frac{(b-a)}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right].$$

Observe that $S(f) = \frac{1}{3}T(f) + \frac{2}{3}M(f)$, where $T(f)$ is the trapezoid rule and $M(f) = (b - a)f\left(\frac{a+b}{2}\right)$ is the midpoint rule.

Simpson's rule

Example: Consider $\int_0^1 f(x)dx \approx w_0f(0) + w_1f(1/2) + w_2f(1)$. Then nodes are 0, 1/2, and 1.

Simpson's rule

Example: Consider $\int_0^1 f(x)dx \approx w_0f(0) + w_1f(1/2) + w_2f(1)$. Then nodes are 0, 1/2, and 1. The Lagrange basis is given by

$$\ell_0(x) = 2 \left(x - \frac{1}{2} \right) (x - 1), \quad \ell_1(x) = -4x(x - 1), \quad \ell_2(x) = 2x \left(x - \frac{1}{2} \right).$$

Simpson's rule

Example: Consider $\int_0^1 f(x)dx \approx w_0f(0) + w_1f(1/2) + w_2f(1)$. Then nodes are 0, 1/2, and 1. The Lagrange basis is given by

$$\ell_0(x) = 2 \left(x - \frac{1}{2} \right) (x - 1), \quad \ell_1(x) = -4x(x - 1), \quad \ell_2(x) = 2x \left(x - \frac{1}{2} \right).$$

$$\text{Then } w_0 = \int_0^1 \ell_0(x) = \frac{1}{6} = \int_0^1 \ell_2(x) = w_2 \quad \text{and} \quad w_1 = \int_0^1 \ell_1(x) = \frac{2}{3}.$$

Hence

$$S(f) = \frac{1}{6} \left[f(0) + 4f\left(\frac{1}{2}\right) + f(1) \right].$$

Simpson's rule

Example: Consider $\int_0^1 f(x)dx \approx w_0f(0) + w_1f(1/2) + w_2f(1)$. Then nodes are 0, 1/2, and 1. The Lagrange basis is given by

$$\ell_0(x) = 2 \left(x - \frac{1}{2} \right) (x - 1), \quad \ell_1(x) = -4x(x - 1), \quad \ell_2(x) = 2x \left(x - \frac{1}{2} \right).$$

$$\text{Then } w_0 = \int_0^1 \ell_0(x) = \frac{1}{6} = \int_0^1 \ell_2(x) = w_2 \quad \text{and} \quad w_1 = \int_0^1 \ell_1(x) = \frac{2}{3}.$$

Hence

$$S(f) = \frac{1}{6} \left[f(0) + 4f\left(\frac{1}{2}\right) + f(1) \right].$$

Observe that $S(x^j) = \int_0^1 x^j dx$ for $j = 0, 1, 2, 3$.

Simpson's rule

Example: Consider $\int_0^1 f(x)dx \approx w_0 f(0) + w_1 f(1/2) + w_2 f(1)$. Then nodes are 0, 1/2, and 1. The Lagrange basis is given by

$$\ell_0(x) = 2 \left(x - \frac{1}{2} \right) (x - 1), \quad \ell_1(x) = -4x(x - 1), \quad \ell_2(x) = 2x \left(x - \frac{1}{2} \right).$$

$$\text{Then } w_0 = \int_0^1 \ell_0(x) = \frac{1}{6} = \int_0^1 \ell_2(x) = w_2 \quad \text{and} \quad w_1 = \int_0^1 \ell_1(x) = \frac{2}{3}.$$

Hence

$$S(f) = \frac{1}{6} \left[f(0) + 4f\left(\frac{1}{2}\right) + f(1) \right].$$

Observe that $S(x^j) = \int_0^1 x^j dx$ for $j = 0, 1, 2, 3$. Hence $S(f)$ is **exact for polynomial of degree ≤ 3** .

Simpson's rule

Example: Consider $\int_0^1 f(x)dx \approx w_0 f(0) + w_1 f(1/2) + w_2 f(1)$. Then nodes are 0, 1/2, and 1. The Lagrange basis is given by

$$\ell_0(x) = 2 \left(x - \frac{1}{2} \right) (x - 1), \quad \ell_1(x) = -4x(x - 1), \quad \ell_2(x) = 2x \left(x - \frac{1}{2} \right).$$

$$\text{Then } w_0 = \int_0^1 \ell_0(x) = \frac{1}{6} = \int_0^1 \ell_2(x) = w_2 \quad \text{and} \quad w_1 = \int_0^1 \ell_1(x) = \frac{2}{3}.$$

Hence

$$S(f) = \frac{1}{6} \left[f(0) + 4f\left(\frac{1}{2}\right) + f(1) \right].$$

Observe that $S(x^j) = \int_0^1 x^j dx$ for $j = 0, 1, 2, 3$. Hence $S(f)$ is **exact for polynomial of degree ≤ 3** . As per the error analysis of quadrature rule, $S(f)$ is expected to be exact for polynomials of degree ≤ 2 .

Error estimation of Simpson's rule

Theorem: Suppose that $f \in C^4[a, b]$. Then

$$\int_a^b f(x) dx = \frac{(b-a)}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] + E_2(f),$$

where

Error estimation of Simpson's rule

Theorem: Suppose that $f \in C^4[a, b]$. Then

$$\int_a^b f(x) dx = \frac{(b-a)}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] + E_2(f),$$

where $E_2(f) = -\frac{1}{90} \left(\frac{b-a}{2} \right)^5 f^{(4)}(\theta)$ for some $\theta \in [a, b]$.

Error estimation of Simpson's rule

Theorem: Suppose that $f \in C^4[a, b]$. Then

$$\int_a^b f(x) dx = \frac{(b-a)}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] + E_2(f),$$

where $E_2(f) = -\frac{1}{90} \left(\frac{b-a}{2} \right)^5 f^{(4)}(\theta)$ for some $\theta \in [a, b]$.

Composite Simpson's rule

Suppose that n is even. Set $h := (b - a)/n$ and consider the nodes $x_j = a + jh$ for $j = 0 : n$. Then

$$\int_a^b f(x) dx = \int_{x_0}^{x_2} f(x) dx + \int_{x_2}^{x_4} f(x) dx + \cdots + \int_{x_{n-2}}^{x_n} f(x) dx$$

Composite Simpson's rule

Suppose that n is even. Set $h := (b - a)/n$ and consider the nodes $x_j = a + jh$ for $j = 0 : n$. Then

$$\begin{aligned}\int_a^b f(x) dx &= \int_{x_0}^{x_2} f(x) dx + \int_{x_2}^{x_4} f(x) dx + \cdots + \int_{x_{n-2}}^{x_n} f(x) dx \\ &= \sum_{j=1}^{n/2} \int_{x_{2j-2}}^{x_{2j}} f(x) dx\end{aligned}$$

Composite Simpson's rule

Suppose that n is even. Set $h := (b - a)/n$ and consider the nodes $x_j = a + jh$ for $j = 0 : n$. Then

$$\begin{aligned}\int_a^b f(x) dx &= \int_{x_0}^{x_2} f(x) dx + \int_{x_2}^{x_4} f(x) dx + \cdots + \int_{x_{n-2}}^{x_n} f(x) dx \\ &= \sum_{j=1}^{n/2} \int_{x_{2j-2}}^{x_{2j}} f(x) dx \approx \sum_{j=1}^{n/2} \frac{h}{3} [f(x_{2j-2}) + 4f(x_{2j-1}) + f(x_{2j})]\end{aligned}$$

Composite Simpson's rule

Suppose that n is even. Set $h := (b - a)/n$ and consider the nodes $x_j = a + jh$ for $j = 0 : n$. Then

$$\begin{aligned}\int_a^b f(x) dx &= \int_{x_0}^{x_2} f(x) dx + \int_{x_2}^{x_4} f(x) dx + \cdots + \int_{x_{n-2}}^{x_n} f(x) dx \\&= \sum_{j=1}^{n/2} \int_{x_{2j-2}}^{x_{2j}} f(x) dx \approx \sum_{j=1}^{n/2} \frac{h}{3} [f(x_{2j-2}) + 4f(x_{2j-1}) + f(x_{2j})] \\&\approx \frac{h}{3} [f(x_0) + 2 \sum_{j=2}^{n/2} f(x_{2j-2}) + 4 \sum_{j=1}^{n/2} f(x_{2j-1}) + f(x_n)].\end{aligned}$$

Composite Simpson's rule

Suppose that n is even. Set $h := (b - a)/n$ and consider the nodes $x_j = a + jh$ for $j = 0 : n$. Then

$$\begin{aligned}\int_a^b f(x) dx &= \int_{x_0}^{x_2} f(x) dx + \int_{x_2}^{x_4} f(x) dx + \cdots + \int_{x_{n-2}}^{x_n} f(x) dx \\&= \sum_{j=1}^{n/2} \int_{x_{2j-2}}^{x_{2j}} f(x) dx \approx \sum_{j=1}^{n/2} \frac{h}{3} [f(x_{2j-2}) + 4f(x_{2j-1}) + f(x_{2j})] \\&\approx \frac{h}{3} [f(x_0) + 2 \sum_{j=2}^{n/2} f(x_{2j-2}) + 4 \sum_{j=1}^{n/2} f(x_{2j-1}) + f(x_n)].\end{aligned}$$

Thus the composite Simpson's rule is given by

$$S_n(f) := \frac{h}{3} [f(x_0) + 2 \sum_{j=2}^{n/2} f(x_{2j-2}) + 4 \sum_{j=1}^{n/2} f(x_{2j-1}) + f(x_n)].$$

Error in composite Simpson's rule

Thus the composite Simpson's rule is given by

$$S_n(f) := \frac{h}{3} \left[f(x_0) + 2 \sum_{j=2}^{n/2} f(x_{2j-2}) + 4 \sum_{j=1}^{n/2} f(x_{2j-1}) + f(x_n) \right]$$

Error in composite Simpson's rule

Thus the composite Simpson's rule is given by

$$S_n(f) := \frac{h}{3} \left[f(x_0) + 2 \sum_{j=2}^{n/2} f(x_{2j-2}) + 4 \sum_{j=1}^{n/2} f(x_{2j-1}) + f(x_n) \right]$$

with error $E_n^S(f) = -\frac{1}{90} h^5 \sum_{j=1}^{n/2} f^{(4)}(\theta_j)$ for some $\theta_j \in (x_{2j-2}, x_{2j})$, $j = 1 : (n/2)$.

Error in composite Simpson's rule

Thus the composite Simpson's rule is given by

$$S_n(f) := \frac{h}{3} \left[f(x_0) + 2 \sum_{j=2}^{n/2} f(x_{2j-2}) + 4 \sum_{j=1}^{n/2} f(x_{2j-1}) + f(x_n) \right]$$

with error $E_n^S(f) = -\frac{1}{90} h^5 \sum_{j=1}^{n/2} f^{(4)}(\theta_j)$ for some $\theta_j \in (x_{2j-2}, x_{2j})$, $j = 1 : (n/2)$.

By IVT (intermediate value theorem), there exists $\theta \in (a, b)$ such that

$$\sum_{j=1}^{n/2} f^{(4)}(\theta_j) = \frac{n}{2} f^{(4)}(\theta)$$

As $nh = b - a$,

$$E_n^S(f) = -\frac{b-a}{180} h^4 f^{(4)}(\theta).$$

Example

Consider the integral

$$I(f) := \int_0^1 e^{-x^2} dx.$$

The three Newton-Cotes (midpoint, trapezoid, and Simpson's rule) formula yield

$$M(f) = (1 - 0)e^{-0.25} \approx 0.778801$$

$$T(f) = \frac{1}{2}[e^0 + e^{-1}] \approx 0.683940$$

$$S(f) = \frac{1}{6}[e^0 + 4e^{-0.25} + e^{-1}] \approx 0.747180$$

The correctly rounded value is $I = 0.746824$.

Example

Consider the integral

$$I(f) := \int_0^1 e^{-x^2} dx.$$

The three Newton-Cotes (midpoint, trapezoid, and Simpson's rule) formula yield

$$M(f) = (1 - 0)e^{-0.25} \approx 0.778801$$

$$T(f) = \frac{1}{2}[e^0 + e^{-1}] \approx 0.683940$$

$$S(f) = \frac{1}{6}[e^0 + 4e^{-0.25} + e^{-1}] \approx 0.747180$$

The correctly rounded value is $I = 0.746824$.

$$|I(f) - M(f)| = 0.031977, |I(f) - T(f)| = 0.062884, |I(f) - S(f)| = 0.000356.$$

The Simpson's rule, with an error of only 0.000356, seems remarkably accurate considering the size of the interval over which it is applied.