

# MA579H Scientific Computing


## Solutions of Nonlinear equations II

Rafikul Alam  
Department of Mathematics  
IIT Guwahati

# Outline

- Newton method
- Fixed point iteration

# Newton's method



# Newton's method

Let

$f \longrightarrow$  continuously differentiable;

$\alpha \in \mathbb{R} \longrightarrow f(\alpha) = 0$ ;

$x_0 \in \mathbb{R} \longrightarrow$  initial guess.

# Newton's method

Let

$f \longrightarrow$  continuously differentiable;

$\alpha \in \mathbb{R} \longrightarrow f(\alpha) = 0$ ;

$x_0 \in \mathbb{R} \longrightarrow$  initial guess.

The tangent to  $y = f(x)$  at  $(x_0, f(x_0))$  is

$$y = f(x_0) + f'(x_0)(x - x_0).$$

Suppose the tangent is not parallel to the  $x$ -axis so that  $f'(x_0) \neq 0$ . Then it cuts the  $x$ -axis at

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}.$$

# Newton's method

Let

$f \longrightarrow$  continuously differentiable;

$\alpha \in \mathbb{R} \longrightarrow f(\alpha) = 0$ ;

$x_0 \in \mathbb{R} \longrightarrow$  initial guess.

The tangent to  $y = f(x)$  at  $(x_0, f(x_0))$  is

$$y = f(x_0) + f'(x_0)(x - x_0).$$

Suppose the tangent is not parallel to the  $x$ -axis so that  $f'(x_0) \neq 0$ . Then it cuts the  $x$ -axis at

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}.$$

Repeating the process results in Newton's method

$$x_{n+1} := x_n - \frac{f(x_n)}{f'(x_n)}, \quad n = 0, 1, \dots$$

# Newton's method

## Algorithm:

**Input:** A differentiable function  $f$

Initial guess  $c \in \mathbb{R}$  and a tolerance  $\text{tol}$

A limit  $N$  for maximum number of iteration.

**Output** Approximate solution  $c$  of  $f(x) = 0$  satisfying  $|f(c)| \leq \text{tol}$ .

# Newton's method

## Algorithm:

**Input:** A differentiable function  $f$

Initial guess  $c \in \mathbb{R}$  and a tolerance  $\text{tol}$

A limit  $N$  for maximum number of iteration.

**Output** Approximate solution  $c$  of  $f(x) = 0$  satisfying  $|f(c)| \leq \text{tol}$ .

- $\text{IT} = 0$
- **while** ( $|f(c)| > \text{tol}$ ) **and** ( $\text{IT} \leq N$ ) **do**
- $c = c - f(c)/f'(c)$
- $\text{IT} = \text{IT} + 1$



# Newton's method

## Algorithm:

**Input:** A differentiable function  $f$

Initial guess  $c \in \mathbb{R}$  and a tolerance `tol`

A limit `N` for maximum number of iteration.

**Output** Approximate solution  $c$  of  $f(x) = 0$  satisfying  $|f(c)| \leq \text{tol}$ .

- $\text{IT} = 0$
- **while** ( $|f(c)| > \text{tol}$ ) **and** ( $\text{IT} \leq N$ ) **do**
- $c = c - f(c)/f'(c)$
- $\text{IT} = \text{IT} + 1$

**Example:** Let  $a > 0$ . Then the square root  $\alpha = \sqrt{a}$  is the zero of  $f(x) = x^2 - a$  which gives

$$x_{n+1} = \frac{1}{2} \left( x_n + \frac{a}{x_n} \right), \quad n = 0, 1, 2, \dots$$

This scheme converges globally, that is,  $x_n \rightarrow \sqrt{a}$  for any  $x_0 \neq 0$ . **Why?**

# Newton's method from Secant method




Figure: Secant method for finding the zero of  $x^3 + x - 1$ .

Given two initial guesses  $x_0 \neq x_1$ , the secant method generates the iterates

$$x_{n+1} = x_n - \frac{f(x_n)}{f[x_{n-1}, x_n]}, \quad n = 1, 2, 3, \dots$$

# Newton's method from Secant method




Figure: Secant method for finding the zero of  $x^3 + x - 1$ .

Given two initial guesses  $x_0 \neq x_1$ , the secant method generates the iterates

$$x_{n+1} = x_n - \frac{f(x_n)}{f[x_{n-1}, x_n]}, \quad n = 1, 2, 3, \dots$$

Letting  $x_{n-1} \rightarrow x_n$ , the secant becomes the tangent to  $y = f(x)$  at  $x_n$  and yields Newton's method

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}, \quad n = 0, 1, 2, \dots$$

# Convergence of Newton's method

Suppose that  $f$  is twice continuously differentiable near  $\alpha$  and that  $\alpha$  is a simple zero, that is,  $f'(\alpha) \neq 0$ .

# Convergence of Newton's method

Suppose that  $f$  is twice continuously differentiable near  $\alpha$  and that  $\alpha$  is a simple zero, that is,  $f'(\alpha) \neq 0$ .

Then we have

$$x_{n+1} - \alpha = x_n - \alpha - \frac{f(x_n)}{f'(x_n)}$$

# Convergence of Newton's method

Suppose that  $f$  is twice continuously differentiable near  $\alpha$  and that  $\alpha$  is a simple zero, that is,  $f'(\alpha) \neq 0$ .

Then we have

$$\begin{aligned}x_{n+1} - \alpha &= x_n - \alpha - \frac{f(x_n)}{f'(x_n)} = (x_n - \alpha) \left( 1 - \frac{f(x_n) - f(\alpha)}{(x_n - \alpha)f'(x_n)} \right) \\&= (x_n - \alpha) \left( 1 - \frac{f[x_n, \alpha]}{f[x_n, x_n]} \right)\end{aligned}$$

# Convergence of Newton's method

Suppose that  $f$  is twice continuously differentiable near  $\alpha$  and that  $\alpha$  is a simple zero, that is,  $f'(\alpha) \neq 0$ .

Then we have

$$\begin{aligned}x_{n+1} - \alpha &= x_n - \alpha - \frac{f(x_n)}{f'(x_n)} = (x_n - \alpha) \left( 1 - \frac{f(x_n) - f(\alpha)}{(x_n - \alpha)f'(x_n)} \right) \\&= (x_n - \alpha) \left( 1 - \frac{f[x_n, \alpha]}{f[x_n, x_n]} \right) = (x_n - \alpha)^2 \frac{f[x_n, x_n, \alpha]}{f[x_n, x_n]}.\end{aligned}$$

# Convergence of Newton's method

Suppose that  $f$  is twice continuously differentiable near  $\alpha$  and that  $\alpha$  is a simple zero, that is,  $f'(\alpha) \neq 0$ .

Then we have

$$\begin{aligned}x_{n+1} - \alpha &= x_n - \alpha - \frac{f(x_n)}{f'(x_n)} = (x_n - \alpha) \left( 1 - \frac{f(x_n) - f(\alpha)}{(x_n - \alpha)f'(x_n)} \right) \\&= (x_n - \alpha) \left( 1 - \frac{f[x_n, \alpha]}{f[x_n, x_n]} \right) = (x_n - \alpha)^2 \frac{f[x_n, x_n, \alpha]}{f[x_n, x_n]}.\end{aligned}$$

Therefore, if  $x_n \rightarrow \alpha$  then

$$\lim_{n \rightarrow \infty} \frac{x_{n+1} - \alpha}{(x_n - \alpha)^2} = \frac{f''(\alpha)}{2f'(\alpha)}.$$



# Convergence of Newton's method

Suppose that  $f$  is twice continuously differentiable near  $\alpha$  and that  $\alpha$  is a simple zero, that is,  $f'(\alpha) \neq 0$ .

Then we have


$$\begin{aligned}x_{n+1} - \alpha &= x_n - \alpha - \frac{f(x_n)}{f'(x_n)} = (x_n - \alpha) \left( 1 - \frac{f(x_n) - f(\alpha)}{(x_n - \alpha)f'(x_n)} \right) \\&= (x_n - \alpha) \left( 1 - \frac{f[x_n, \alpha]}{f[x_n, x_n]} \right) = (x_n - \alpha)^2 \frac{f[x_n, x_n, \alpha]}{f[x_n, x_n]}.\end{aligned}$$

Therefore, if  $x_n \rightarrow \alpha$  then

$$\lim_{n \rightarrow \infty} \frac{x_{n+1} - \alpha}{(x_n - \alpha)^2} = \frac{f''(\alpha)}{2f'(\alpha)}.$$

This shows that the order of convergence of Newton's method is quadratic if  $f''(\alpha) \neq 0$ .

# Non-convergence of Newton's method



# Non-convergence Newton's method

**Example:** Now consider  $f(x) := \sin(x)$  for  $|x| < \pi/2$ . Then  $\alpha = 0$  is the only zero. Newton's method becomes  $x_{n+1} = x_n - \tan(x_n)$ ,  $n = 0, 1, \dots$

If  $\hat{x}$  is such that  $2\hat{x} = \tan(\hat{x})$  then with  $x_0 := \hat{x}$ , we have  $x_1 = -\hat{x}$  and  $x_2 = \hat{x}$ . Hence after two iterations Newton's method ends up with where we started. This is called cycle.

# Non-convergence Newton's method

**Example:** Now consider  $f(x) := \sin(x)$  for  $|x| < \pi/2$ . Then  $\alpha = 0$  is the only zero. Newton's method becomes  $x_{n+1} = x_n - \tan(x_n)$ ,  $n = 0, 1, \dots$

If  $\hat{x}$  is such that  $2\hat{x} = \tan(\hat{x})$  then with  $x_0 := \hat{x}$ , we have  $x_1 = -\hat{x}$  and  $x_2 = \hat{x}$ . Hence after two iterations Newton's method ends up where we started. This is called cycle.

This shows that Newton's method does not converge for  $x_0 = \hat{x}$ . However, Newton's method converges for any  $x_0$  such that  $|x_0| < |\hat{x}|$ . ■

# Non-convergence Newton's method

**Example:** Now consider  $f(x) := \sin(x)$  for  $|x| < \pi/2$ . Then  $\alpha = 0$  is the only zero. Newton's method becomes  $x_{n+1} = x_n - \tan(x_n)$ ,  $n = 0, 1, \dots$

If  $\hat{x}$  is such that  $2\hat{x} = \tan(\hat{x})$  then with  $x_0 := \hat{x}$ , we have  $x_1 = -\hat{x}$  and  $x_2 = \hat{x}$ . Hence after two iterations Newton's method ends up with where we started. This is called cycle.

This shows that Newton's method does not converge for  $x_0 = \hat{x}$ . However, Newton's method converges for any  $x_0$  such that  $|x_0| < |\hat{x}|$ . ■

**Observation:** If  $f \in C^2[a, b]$  is convex (or concave) and  $f(a)f(b) < 0$  then the tangents at the endpoints of  $[a, b]$  intersects  $x$ -axis within  $[a, b]$ . Hence on geometric ground Newton's method converges globally.

# Slow convergence of Newton's method

If  $\alpha$  is a zero of  $f$  with multiplicity more than 1 then  $f'(\alpha) = 0$  and the previous convergence analysis of Newton's Method does not apply.

# Slow convergence of Newton's method

If  $\alpha$  is a zero of  $f$  with multiplicity more than 1 then  $f'(\alpha) = 0$  and the previous convergence analysis of Newton's Method does not apply.

**Example:** For  $f(x) = x^m$ , the Newton iterations are

$$x_{n+1} = x_n - \frac{x_n^m}{mx_n^{m-1}} = \left(\frac{m-1}{m}\right)x_n.$$

Since  $\alpha := 0$  is the zero of  $f$ , we have  $e_n := |x_n - \alpha| = |x_n|$  and  $\frac{e_{n+1}}{e_n} = \frac{m-1}{m}$ . Hence the convergence is linear at the rate  $\frac{m-1}{m}$ .

# Slow convergence of Newton's method

If  $\alpha$  is a zero of  $f$  with multiplicity more than 1 then  $f'(\alpha) = 0$  and the previous convergence analysis of Newton's Method does not apply.

**Example:** For  $f(x) = x^m$ , the Newton iterations are

$$x_{n+1} = x_n - \frac{x_n^m}{mx_n^{m-1}} = \left( \frac{m-1}{m} \right) x_n.$$

Since  $\alpha := 0$  is the zero of  $f$ , we have  $e_n := |x_n - \alpha| = |x_n|$  and  $\frac{e_{n+1}}{e_n} = \frac{m-1}{m}$ . Hence the convergence is linear at the rate  $\frac{m-1}{m}$ .

**Theorem** Let  $f$  be an  $(m+1)$ -times continuously differentiable function on  $[a, b]$ . Let  $\alpha \in (a, b)$  be a zero of  $f$  of multiplicity  $m$ . Then Newton's Method is locally convergent to  $\alpha$  and the convergence is linear at the rate  $\frac{m-1}{m}$ .



# Modified Newton's Method for multiple roots

**Example:** The function  $f(x) = \sin x + x^2 \cos x - x^2 - x$  has a root of multiplicity 3 at 0. With starting guess  $x_0 = 1$ ,  $e_0 = 1$  and  $e_{n+1} \leq (2/3)^n$  and the number of steps say  $n$ , required to achieve accuracy up to 6 decimal digits

$$(2/3)^n < 0.5 * 10^{-6} \Rightarrow n > \frac{\log_{10}(0.5) - 6}{\log_{10}(2/3)} \approx 35.78.$$

# Modified Newton's Method for multiple roots

**Example:** The function  $f(x) = \sin x + x^2 \cos x - x^2 - x$  has a root of multiplicity 3 at 0. With starting guess  $x_0 = 1$ ,  $e_0 = 1$  and  $e_{n+1} \leq (2/3)^n$  and the number of steps say  $n$ , required to achieve accuracy up to 6 decimal digits

$$(2/3)^n < 0.5 * 10^{-6} \Rightarrow n > \frac{\log_{10}(0.5) - 6}{\log_{10}(2/3)} \approx 35.78.$$

**Theorem** Let  $f$  be an  $(m + 1)$ -times continuously differentiable function on  $[a, b]$ . Let  $\alpha \in (a, b)$  be a zero of  $f$  of multiplicity  $m$ . Then the modified Newton's Method

$$x_{n+1} = x_n - \frac{mf(x_n)}{f'(x_n)}$$

converges locally to  $\alpha$  and the rate of convergence is quadratic.

# Modified Newton's Method for multiple roots

**Example:** The function  $f(x) = \sin x + x^2 \cos x - x^2 - x$  has a root of multiplicity 3 at 0. With starting guess  $x_0 = 1$ ,  $e_0 = 1$  and  $e_{n+1} \leq (2/3)^n$  and the number of steps say  $n$ , required to achieve accuracy up to 6 decimal digits

$$(2/3)^n < 0.5 * 10^{-6} \Rightarrow n > \frac{\log_{10}(0.5) - 6}{\log_{10}(2/3)} \approx 35.78.$$

**Theorem** Let  $f$  be an  $(m + 1)$ -times continuously differentiable function on  $[a, b]$ . Let  $\alpha \in (a, b)$  be a zero of  $f$  of multiplicity  $m$ . Then the modified Newton's Method

$$x_{n+1} = x_n - \frac{mf(x_n)}{f'(x_n)}$$

converges locally to  $\alpha$  and the rate of convergence is quadratic.

# Modified Newton's Method for multiple roots

**Example:** The function  $f(x) = \sin x + x^2 \cos x - x^2 - x$  has a root of multiplicity 3 at 0. With starting guess  $x_0 = 1$ ,  $e_0 = 1$  and  $e_{n+1} \leq (2/3)^n$  and the number of steps say  $n$ , required to achieve accuracy up to 6 decimal digits

$$(2/3)^n < 0.5 * 10^{-6} \Rightarrow n > \frac{\log_{10}(0.5) - 6}{\log_{10}(2/3)} \approx 35.78.$$

**Theorem** Let  $f$  be an  $(m + 1)$ -times continuously differentiable function on  $[a, b]$ . Let  $\alpha \in (a, b)$  be a zero of  $f$  of multiplicity  $m$ . Then the modified Newton's Method

$$x_{n+1} = x_n - \frac{mf(x_n)}{f'(x_n)}$$

converges locally to  $\alpha$  and the rate of convergence is quadratic.

The Modified Newton's Method applied to  $f(x) = \sin x + x^2 \cos x - x^2 - x$  produces a solution correct up to 6 decimal digits in 5 iterations.

# Fixed Point

Consider a function  $\phi : [a, b] \rightarrow \mathbb{R}$ . A point  $c \in [a, b]$  is said to be a **fixed point** of  $\phi$  if  $\phi(c) = c$ .

# Fixed Point

Consider a function  $\phi : [a, b] \rightarrow \mathbb{R}$ . A point  $c \in [a, b]$  is said to be a **fixed point** of  $\phi$  if  $\phi(c) = c$ .

## Examples:

- (a)  $f(x) = x^2 + 3x + 1$  has a fixed point at  $-1$ .
- (b)  $f(x) = \sin x$  has a fixed point at  $0$ .
- (c) If  $\alpha$  is a zero of any function  $f : [a, b] \rightarrow \mathbb{R}$ , then  $\alpha$  is a fixed point of  $g(x) := x + f(x)$ .

# Fixed Point

Consider a function  $\phi : [a, b] \rightarrow \mathbb{R}$ . A point  $c \in [a, b]$  is said to be a **fixed point** of  $\phi$  if  $\phi(c) = c$ .

## Examples:

- (a)  $f(x) = x^2 + 3x + 1$  has a fixed point at  $-1$ .
- (b)  $f(x) = \sin x$  has a fixed point at  $0$ .
- (c) If  $\alpha$  is a zero of any function  $f : [a, b] \rightarrow \mathbb{R}$ , then  $\alpha$  is a fixed point of  $g(x) := x + f(x)$ .

*When does a function  $f$  have a unique fixed point?*

# Fixed Point

Consider a function  $\phi : [a, b] \rightarrow \mathbb{R}$ . A point  $c \in [a, b]$  is said to be a **fixed point** of  $\phi$  if  $\phi(c) = c$ .

## Examples:

- (a)  $f(x) = x^2 + 3x + 1$  has a fixed point at  $-1$ .
- (b)  $f(x) = \sin x$  has a fixed point at  $0$ .
- (c) If  $\alpha$  is a zero of any function  $f : [a, b] \rightarrow \mathbb{R}$ , then  $\alpha$  is a fixed point of  $g(x) := x + f(x)$ .

*When does a function  $f$  have a unique fixed point?* Ans: Contractive function.



# Fixed Point

Consider a function  $\phi : [a, b] \rightarrow \mathbb{R}$ . A point  $c \in [a, b]$  is said to be a **fixed point** of  $\phi$  if  $\phi(c) = c$ .

## Examples:

- (a)  $f(x) = x^2 + 3x + 1$  has a fixed point at  $-1$ .
- (b)  $f(x) = \sin x$  has a fixed point at  $0$ .
- (c) If  $\alpha$  is a zero of any function  $f : [a, b] \rightarrow \mathbb{R}$ , then  $\alpha$  is a fixed point of  $g(x) := x + f(x)$ .

*When does a function  $f$  have a unique fixed point?* Ans: Contractive function.

A function  $f : [a, b] \rightarrow \mathbb{R}$  is said to be a **contractive function or contraction on  $[a, b]$**  if there exists  $\lambda \in \mathbb{R}$  with  $0 < \lambda < 1$  such that  $|f(x) - f(y)| \leq \lambda|x - y|$  for all  $x, y \in [a, b]$ .

# Fixed Point

Consider a function  $\phi : [a, b] \rightarrow \mathbb{R}$ . A point  $c \in [a, b]$  is said to be a **fixed point** of  $\phi$  if  $\phi(c) = c$ .

## Examples:

- (a)  $f(x) = x^2 + 3x + 1$  has a fixed point at  $-1$ .
- (b)  $f(x) = \sin x$  has a fixed point at  $0$ .
- (c) If  $\alpha$  is a zero of any function  $f : [a, b] \rightarrow \mathbb{R}$ , then  $\alpha$  is a fixed point of  $g(x) := x + f(x)$ .

*When does a function  $f$  have a unique fixed point?* Ans: Contractive function.

A function  $f : [a, b] \rightarrow \mathbb{R}$  is said to be a **contractive function or contraction on  $[a, b]$**  if there exists  $\lambda \in \mathbb{R}$  with  $0 < \lambda < 1$  such that  $|f(x) - f(y)| \leq \lambda|x - y|$  for all  $x, y \in [a, b]$ .

## Examples:

- (a)  $f(x) = \frac{x^2 + 2x + 4}{4}$  is a contraction on  $(-2, 0.75)$ .
- (b)  $f(x) = \sin x$  is a contraction on  $[\frac{\pi}{6}, \frac{5\pi}{6}]$ .
- (c)  $f(x) = \cos x$  is a contraction on  $[0, \frac{\pi}{3}]$ .

# Fixed-point iteration

Let  $\phi : [a, b] \rightarrow \mathbb{R}$ . Suppose that  $\phi$  has a fixed-point in  $[a, b]$ . For  $x_0 \in [a, b]$ , consider the fixed-point iteration (FPI)

$$x_n = \phi(x_{n-1}), \quad n = 1, 2, \dots$$

If  $(x_n)$  converges to  $\alpha$  then  $\phi(\alpha) = \alpha$ . Thus  $\alpha$  is a fixed-point.

# Fixed-point iteration

Let  $\phi : [a, b] \rightarrow \mathbb{R}$ . Suppose that  $\phi$  has a fixed-point in  $[a, b]$ . For  $x_0 \in [a, b]$ , consider the fixed-point iteration (FPI)

$$x_n = \phi(x_{n-1}), \quad n = 1, 2, \dots$$

If  $(x_n)$  converges to  $\alpha$  then  $\phi(\alpha) = \alpha$ . Thus  $\alpha$  is a fixed-point.

**Remark:** The function  $\phi$  is called the **iteration function** for the fixed-point iteration.

# Fixed-point iteration

Let  $\phi : [a, b] \rightarrow \mathbb{R}$ . Suppose that  $\phi$  has a fixed-point in  $[a, b]$ . For  $x_0 \in [a, b]$ , consider the fixed-point iteration (FPI)

$$x_n = \phi(x_{n-1}), \quad n = 1, 2, \dots$$

If  $(x_n)$  converges to  $\alpha$  then  $\phi(\alpha) = \alpha$ . Thus  $\alpha$  is a fixed-point.


**Remark:** The function  $\phi$  is called the **iteration function** for the fixed-point iteration.

**Example:** Suppose we wish to solve  $\cos(e^x) = 0$ . Note that


$$\cos(e^c) = 0 \iff \cos(e^c) + c = c.$$

Setting  $\phi(x) := \cos(e^x) + x$ , we have  $\phi(c) = c$ . Hence the fixed-point iteration  $x_n = \phi(x_{n-1})$  can be used to solve  $\cos(e^x) = 0$ .

# Fixed-point iteration



# Fixed-point iteration



# Convergence of fixed-point iteration

**Question:** When does a function  $\phi$  have a fixed point?



# Convergence of fixed-point iteration

**Question:** When does a function  $\phi$  have a fixed point? When does a fixed-point iteration converge?

# Convergence of fixed-point iteration

**Question:** When does a function  $\phi$  have a fixed point? When does a fixed-point iteration converge?

A function  $\phi : [a, b] \rightarrow \mathbb{R}$  is said to be a contraction (or contractive) if there exists a constant  $0 < \lambda < 1$  such that

$$|\phi(x) - \phi(y)| \leq \lambda |x - y| \text{ for all } x, y \in [a, b].$$

# Convergence of fixed-point iteration

**Question:** When does a function  $\phi$  have a fixed point? When does a fixed-point iteration converge?

A function  $\phi : [a, b] \rightarrow \mathbb{R}$  is said to be a contraction (or contractive) if there exists a constant  $0 < \lambda < 1$  such that

$$|\phi(x) - \phi(y)| \leq \lambda|x - y| \text{ for all } x, y \in [a, b].$$

**Theorem:** If  $\phi : [a, b] \rightarrow \mathbb{R}$  is a contraction then it has a unique fixed point. Further, for any  $x_0 \in [a, b]$ , the iteration  $x_n = \phi(x_{n-1})$  converges to the unique fixed point.

**Proof:** Consider  $x_n = \phi(x_{n-1})$ ,  $n = 1, 2, \dots$ . Then  $|x_{n+1} - x_n| \leq \lambda|x_n - x_{n-1}| \Rightarrow (x_n)$  is a Cauchy sequence (Check).

# Convergence of fixed-point iteration

**Question:** When does a function  $\phi$  have a fixed point? When does a fixed-point iteration converge?

A function  $\phi : [a, b] \rightarrow \mathbb{R}$  is said to be a contraction (or contractive) if there exists a constant  $0 < \lambda < 1$  such that

$$|\phi(x) - \phi(y)| \leq \lambda|x - y| \text{ for all } x, y \in [a, b].$$

**Theorem:** If  $\phi : [a, b] \rightarrow \mathbb{R}$  is a contraction then it has a unique fixed point. Further, for any  $x_0 \in [a, b]$ , the iteration  $x_n = \phi(x_{n-1})$  converges to the unique fixed point.

**Proof:** Consider  $x_n = \phi(x_{n-1})$ ,  $n = 1, 2, \dots$ . Then  $|x_{n+1} - x_n| \leq \lambda|x_n - x_{n-1}| \Rightarrow (x_n)$  is a Cauchy sequence (Check).

Hence  $x_n \rightarrow c$  for some  $c \in [a, b]$ . Now  $x_n = \phi(x_{n-1}) \Rightarrow c = \phi(c)$ . The uniqueness of  $c$  is immediate. ■

**Observation:** The fixed point iteration converges linearly at the rate  $\lambda$ .

## Sufficient condition for contraction

**Fact:** Let  $\phi$  be differentiable on  $[a, b]$  such that  $\max_{x \in [a, b]} |\phi'(x)| < 1$ .  
Then  $\phi$  is a contraction on  $[a, b]$ .

## Sufficient condition for contraction

**Fact:** Let  $\phi$  be differentiable on  $[a, b]$  such that  $\max_{x \in [a, b]} |\phi'(x)| < 1$ . Then  $\phi$  is a contraction on  $[a, b]$ . Indeed, by mean value theorem  $|\phi(x) - \phi(y)| = |\phi'(c)| |x - y|$  shows that  $\phi(x)$  is a contraction on  $[a, b]$ .

## Sufficient condition for contraction

**Fact:** Let  $\phi$  be differentiable on  $[a, b]$  such that  $\max_{x \in [a, b]} |\phi'(x)| < 1$ . Then  $\phi$  is a contraction on  $[a, b]$ . Indeed, by mean value theorem  $|\phi(x) - \phi(y)| = |\phi'(c)| |x - y|$  shows that  $\phi(x)$  is a contraction on  $[a, b]$ .

**Example:** Consider  $\phi(x) = \cos(e^x) + x$ . Then

$$\phi'(x) = 1 - e^x \sin(e^x) \Rightarrow \lambda := \max_{x \in [0, 1]} |\phi'(x)| < 1.$$

Hence  $\phi(x)$  is a contraction on  $[0, 1]$ .

## Sufficient condition for contraction

**Fact:** Let  $\phi$  be differentiable on  $[a, b]$  such that  $\max_{x \in [a, b]} |\phi'(x)| < 1$ . Then  $\phi$  is a contraction on  $[a, b]$ . Indeed, by mean value theorem  $|\phi(x) - \phi(y)| = |\phi'(c)| |x - y|$  shows that  $\phi(x)$  is a contraction on  $[a, b]$ .

**Example:** Consider  $\phi(x) = \cos(e^x) + x$ . Then

$$\phi'(x) = 1 - e^x \sin(e^x) \Rightarrow \lambda := \max_{x \in [0, 1]} |\phi'(x)| < 1.$$

Hence  $\phi(x)$  is a contraction on  $[0, 1]$ .

**Newton method as a fixed-point iteration:** Consider  $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$ .

Then setting  $\phi(x) := x - \frac{f(x)}{f'(x)}$ , we have  $x_{n+1} = \phi(x_n)$ ,  $n = 0, 1, \dots$



## Sufficient condition for contraction

**Fact:** Let  $\phi$  be differentiable on  $[a, b]$  such that  $\max_{x \in [a, b]} |\phi'(x)| < 1$ . Then  $\phi$  is a contraction on  $[a, b]$ . Indeed, by mean value theorem  $|\phi(x) - \phi(y)| = |\phi'(c)| |x - y|$  shows that  $\phi(x)$  is a contraction on  $[a, b]$ .

**Example:** Consider  $\phi(x) = \cos(e^x) + x$ . Then

$$\phi'(x) = 1 - e^x \sin(e^x) \Rightarrow \lambda := \max_{x \in [0, 1]} |\phi'(x)| < 1.$$

Hence  $\phi(x)$  is a contraction on  $[0, 1]$ .

**Newton method as a fixed-point iteration:** Consider  $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$ .

Then setting  $\phi(x) := x - \frac{f(x)}{f'(x)}$ , we have  $x_{n+1} = \phi(x_n)$ ,  $n = 0, 1, \dots$

This shows that if  $\phi$  is a contraction on  $[a, b]$  then for any  $x_0 \in [a, b]$ , the Newton method will converge to a solution of  $f(x) = 0$ .

## Sufficient condition for contraction

**Fact:** Let  $\phi$  be differentiable on  $[a, b]$  such that  $\max_{x \in [a, b]} |\phi'(x)| < 1$ . Then  $\phi$  is a contraction on  $[a, b]$ . Indeed, by mean value theorem  $|\phi(x) - \phi(y)| = |\phi'(c)| |x - y|$  shows that  $\phi(x)$  is a contraction on  $[a, b]$ .

**Example:** Consider  $\phi(x) = \cos(e^x) + x$ . Then

$$\phi'(x) = 1 - e^x \sin(e^x) \Rightarrow \lambda := \max_{x \in [0, 1]} |\phi'(x)| < 1.$$

Hence  $\phi(x)$  is a contraction on  $[0, 1]$ .

**Newton method as a fixed-point iteration:** Consider  $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$ .

Then setting  $\phi(x) := x - \frac{f(x)}{f'(x)}$ , we have  $x_{n+1} = \phi(x_n)$ ,  $n = 0, 1, \dots$

This shows that if  $\phi$  is a contraction on  $[a, b]$  then for any  $x_0 \in [a, b]$ , the Newton method will converge to a solution of  $f(x) = 0$ . Now

$$\phi'(x) = 1 - \frac{f'(x)^2 - f(x)f''(x)}{f'(x)^2} = \frac{f(x)f''(x)}{f'(x)^2}$$

yields a sufficient condition for convergence of Newton method.

# Choice of iteration function

Consider the equation  $f(x) := xe^x - 1 = 0$ . Then fixed-point iterations are obtained as follows:

- $xe^x - 1 = 0 \iff x = e^{-x} \Rightarrow x_n = e^{-x_{n-1}} = \phi(x_{n-1}), n = 1, 2, \dots$

With the initial guess  $x_0 = 0.5$ ,  $x_n$  converges to  $c = 0.5671432\dots$

## Choice of iteration function

Consider the equation  $f(x) := xe^x - 1 = 0$ . Then fixed-point iterations are obtained as follows:

- $xe^x - 1 = 0 \iff x = e^{-x} \Rightarrow x_n = e^{-x_{n-1}} = \phi(x_{n-1}), n = 1, 2, \dots$   
With the initial guess  $x_0 = 0.5$ ,  $x_n$  converges to  $c = 0.5671432\dots$
- $xe^x - 1 = 0 \iff xe^x + x = 1 + x \iff x = \frac{1+x}{1+e^x}$  which gives

$$x_n = \frac{1 + x_{n-1}}{1 + e^{x_{n-1}}} = \phi(x_{n-1}), n = 1, 2, \dots$$

This time the convergence is much faster - we need only three iterations to obtain a 10-digit approximation of  $c$ . Indeed,  $x_1 = 0.5663110032$ ,  $x_2 = 0.5671431650$  and  $x_3 = 0.5671432904$ .

## Choice of iteration function

Consider the equation  $f(x) := xe^x - 1 = 0$ . Then fixed-point iterations are obtained as follows:

- $xe^x - 1 = 0 \iff x = e^{-x} \Rightarrow x_n = e^{-x_{n-1}} = \phi(x_{n-1}), n = 1, 2, \dots$   
With the initial guess  $x_0 = 0.5$ ,  $x_n$  converges to  $c = 0.5671432\dots$
- $xe^x - 1 = 0 \iff xe^x + x = 1 + x \iff x = \frac{1+x}{1+e^x}$  which gives

$$x_n = \frac{1 + x_{n-1}}{1 + e^{x_{n-1}}} = \phi(x_{n-1}), n = 1, 2, \dots$$


This time the convergence is much faster - we need only three iterations to obtain a 10-digit approximation of  $c$ . Indeed,  $x_1 = 0.5663110032$ ,  $x_2 = 0.5671431650$  and  $x_3 = 0.5671432904$ .

- $xe^x - 1 = 0 \iff x = x + 1 - xe^x$  which yields the iteration

$$x_n = x_{n-1} + 1 - x_{n-1}e^{x_{n-1}} = \phi(x_{n-1}), n = 1, 2, \dots$$

However, this iteration function  $\phi$  does not generate a convergent sequence.

# Non-convergence of Fixed-point iteration



# Finding zeros via fixed-point iterations

Note that if  $\alpha$  is a solution  $\alpha$  of  $f(x) = 0$  then  $\alpha$  is a fixed point of  $g(x) := f(x) + x$ .

In fact, solutions of nonlinear equations may be viewed as fixed points of infinitely many functions.

# Finding zeros via fixed-point iterations

Note that if  $\alpha$  is a solution  $\alpha$  of  $f(x) = 0$  then  $\alpha$  is a fixed point of  $g(x) := f(x) + x$ .

In fact, solutions of nonlinear equations may be viewed as fixed points of infinitely many functions.

**Example:** A solution of  $x^3 + x - 1 = 0$  may be found as a fixed point of some  $g(x)$

$$(a) \ g(x) = 1 - x^3 \quad (b) \ g(x) = (1 - x)^{\frac{1}{3}} \quad (c) \ g(x) = \frac{1 + 2x^3}{1 + 3x^2}.$$



# Finding zeros via fixed-point iterations

Note that if  $\alpha$  is a solution  $\alpha$  of  $f(x) = 0$  then  $\alpha$  is a fixed point of  $g(x) := f(x) + x$ .

In fact, solutions of nonlinear equations may be viewed as fixed points of infinitely many functions.


**Example:** A solution of  $x^3 + x - 1 = 0$  may be found as a fixed point of some  $g(x)$

$$(a) \ g(x) = 1 - x^3 \quad (b) \ g(x) = (1 - x)^{\frac{1}{3}} \quad (c) \ g(x) = \frac{1 + 2x^3}{1 + 3x^2}.$$

With  $x_0 = 0.5$  the iterations  $x_{i+1} = g(x_i)$


- *do not converge* for (a),
- *converge very slowly* for (b).
- **converge in 7 iterations** for (c).

# Finding zeros via fixed-point iterations



**Figure:** Diagrams for FPIs to find a zero of  $x^3 + x - 1$  showing (a) no convergence with  $g(x) = 1 - x^3$  (b) slow convergence for  $g(x) = (1 - x)^{1/3}$  and (c) fast convergence for  $g(x) = \frac{1+2x^3}{1+3x^2}$ .

# Finding zeros via fixed-point iterations



**Figure:** Diagrams for FPIs to find a zero of  $x^3 + x - 1$  showing (a) no convergence with  $g(x) = 1 - x^3$  (b) slow convergence for  $g(x) = (1 - x)^{1/3}$  and (c) fast convergence for  $g(x) = \frac{1+2x^3}{1+3x^2}$ .

Here  $r \approx 0.6823$  is the only real solution of  $x^3 + x - 1 = 0$  and

- $|g'(r)| \approx 1.3966 > 1$  for  $g(x) = 1 - x^3$ ;
- $|g'(r)| \approx 0.716 < 1$  for  $g(x) = (1 - x)^{1/3}$ ;
- $|g'(r)| = 0$  for  $g(x) = \frac{1+2x^3}{1+3x^2}$ .