

CS410 HW1

Russell Miller

October 26, 2011

1.8 I choose a number uniformly at random from the range [1, 1,000,000]. Using the inclusion-exclusion principle, determine the probability that the number chosen is divisible by one or more of 4, 6, and 9.

I wasn't sure exactly how to get these sets of numbers, so I wrote a script in Python that generated them. I checked each set of numbers to get intersects, and produced a size of each set. Let each variable $divX$ represent how many numbers are divisible by X , with multiple X s being the intersects, and mil just being the number 1,000,000. Here is the final formula I used to get this probability.

$$\begin{aligned}P(div4 \cup div6 \cup div9) &= P(div4) + P(div6) + P(div9) - P(div4 \cap div6) - P(div4 \cap div9) - \\&\quad P(div6 \cap div9) + P(div4 \cap div6 \cap div9) \\&= \frac{div4}{mil} + \frac{div6}{mil} + \frac{div9}{mil} - \frac{div46}{mil} - \frac{div49}{mil} - \frac{div69}{mil} + \frac{div469}{mil} \\&= \frac{250000}{mil} + \frac{166666}{mil} + \frac{111111}{mil} - \frac{83333}{mil} - \frac{27777}{mil} - \frac{55555}{mil} + \frac{27777}{mil} \\&\quad = .3\bar{8}9\end{aligned}$$

See attached transcript of Python's script.

1.10 I have a fair coin and a two-headed coin. I choose one of the two coins randomly with equal probability and flip it. Given that the flip was heads, what is the probability that I flipped the two-headed coin?

Let A be the event of choosing the two-headed coin. Let B be the event of flipping a heads. We're told that $P(B) = \frac{1}{2}$. Getting a heads while flipping the two-headed coin is 1. So $P(B | A) = 1$. Since there are 4 total possible outcomes (Heads, Heads, Heads, Tails), getting Heads is $\frac{3}{4}$, which is $P(A)$.

$$\begin{aligned}P(A | B) &= \frac{P(A \cap B)}{P(B)} \\&= \frac{P(B | A)P(A)}{P(B)} \\&= \frac{1 \times \frac{3}{4}}{\frac{1}{2}} \\&\quad = \frac{3}{2}\end{aligned}$$