PARAMETRIZING A POPULATION BALANCE EQUATION

ROBERT A. MILTON

Department of Chemical, Materials and Biological Engineering, The University of Sheffield, Sheffield S1 3JD, United Kingdom

Abstract. Surely this cannot be that difficult.

1. Introduction

A plethora of quantities arise in this work, so we endeavour to consistently indicate their nature at a glance. All constants $\|\cdot\|, \langle\cdot\rangle, |\cdot| \in \overline{\mathbb{R}^+}$ are non-negative, and enclosed in $\|\cdot\|$ or $\langle\cdot\rangle$ without meaning (there are no norms or inner products here), or $|\cdot|$ only when taking the absolute value of $\cdot$. All *italic* quantities are dimensionless, to ease the unavoidable chore of tracking units. Undefined notation adheres to the Digital Library of Mathematical Functions [**?**].

A number $\mathrm{n}(\mathrm{t},\mathrm{L})$ (m^{-4}) of particles of length $\mathrm{L}\,(\mathrm{m})$ grow within each unit of reactor volume $\mathrm{V}\,(\mathrm{m}^3)$ as seconds $\mathrm{t}\,(\mathrm{s})$ pass. Conserving mass, the Population Balance Equation is

$$\begin{aligned} \partial_{\mathrm{t}}(\mathrm{Vn}(\mathrm{t},\mathrm{L})) &= -\mathrm{G}^+\partial_{\mathrm{L}}(\mathrm{Vn}(\mathrm{t},\mathrm{L})) \\ \mathrm{n}(0,\mathrm{L}) &= 0 \\ \mathrm{n}(\mathrm{t},0) &= \mathrm{B}^+/\mathrm{G}^+ \end{aligned} \tag{1.1}$$

where $\mathrm{B}^+\,(\mathrm{m}^{-3}\mathrm{s}^{-1})$ and $\mathrm{G}^+\,(\mathrm{m\,s}^{-1})$ are particle birth (nucleation) and growth rates. These relate empirically to supersaturation $S > 1$:

$$\begin{aligned} \text{birth} \quad \mathrm{B}^+ &:= \|\mathrm{B}\|\,(S-1)^{+|b|} \\ \text{growth} \quad \mathrm{G}^+ &:= \|\mathrm{G}\|\,(S-1)^{+|g|} \end{aligned} \tag{1.2}$$

where $(\#)^+ := \max(0,\#)$ denotes the positive part of $(\#)$. In experiments, critical supersaturation $S = 1$ might occur many times, but we cannot allow this. We must insist that

$$\text{supersaturation} \quad S \in (1, \sup S] \quad \text{at all times} \quad \mathrm{t} > \mathrm{t}^* \tag{1.3}$$

for finite supersaturation $\sup S \in \mathbb{R}$ and critical time $\mathrm{t}^* \in \mathbb{R}$ at which S first exceeds 1. This is not limiting per se, but it does mean that each critical instant in an experiment theoretically begins a new experiment. Throughout this work, an asterisk always denotes the critical instant $t = S - 1 = 0$

Given lengthscale $\|\mathrm{L}\|\,(\mathrm{m})$ and timescale $\|\mathrm{t}\|\,(\mathrm{s})$, dimensionless prefactors are defined as

$$\begin{aligned} \text{birth} \quad \|B\| &:= \|\mathrm{B}\|\,\|\mathrm{L}\|^3\,\|\mathrm{t}\| \\ \text{growth} \quad \|G\| &:= \|\mathrm{G}\|\,\|\mathrm{L}\|^{-1}\,\|\mathrm{t}\| \end{aligned} \tag{1.4}$$

together with dimensionless

$$\begin{aligned} \text{time} \quad t &:= \|\mathrm{t}\|^{-1}\,(\mathrm{t}-\mathrm{t}^*) \\ \text{volume} \quad V(t) &:= \|\mathrm{L}\|^{-3}\,\mathrm{V} \end{aligned} \tag{1.5}$$

This work aims to elucidate the dimensionless empirical birth and growth parameters $\|B\|, |b|, \|G\|, |g|$.

Email address: r.a.milton@sheffield.ac.uk.

Key words and phrases. Asymptotics.

This work was funded by Solomon f. Brown.

2. The Crystal Size Distribution (CSD)

The Population Balance Equation conveys a symmetry between synthetic 'time' T and 'length' L defined as

$$S(t) := \max(S, 1) \tag{2.1a}$$

$$T := \|\mathrm{L}\|^{-1} \int_{\mathrm{t}^*}^{\mathrm{t}} \mathrm{G}^+ \,\mathrm{dt} = \int_0^t \|G\| \left(S(\dot{t}) - 1\right)^{|g|} \mathrm{d}\dot{t} \tag{2.1b}$$

$$L := T - \|\mathrm{L}\|^{-1} \mathrm{L} \tag{2.1c}$$

These forms are chosen to render the PBE linear and invariant to exchanging T and L. This symmetry manifests in the particle count $N(T)f(T, L)$ defined as

$$N(\mathbb{R}^-) := \{0\} \tag{2.2a}$$

$$N(T) := \int_{\mathrm{t}^*}^{\mathrm{t}} \mathrm{B}^+ \mathrm{V} \,\mathrm{d\dot{t}} = \int_0^t \|B\| \left(S(\dot{t}) - 1\right)^{|b|} V(\dot{t}) \,\mathrm{d}\dot{t} \tag{2.2b}$$

$$f(T, L) := N(T)^{-1} \|\mathrm{L}\|^{-1} \mathrm{Vn(t, L)} \tag{2.2c}$$

Substituting these variables in Population Balance **??** immediately yields

$$\begin{aligned} \partial_T N(T) f(T, L) &= 0 \\ N(T) f(T, L) &= \partial_t N(T) / \partial_t T = \partial_T N(T) \end{aligned} \tag{2.3}$$

These two lines combine to establish

$$\begin{aligned} f(T, L) &= N(T)^{-1} \partial_L N(L) \\ N(T)^{-1} N(L) &= \int_0^L f(T, \dot{L}) \,\mathrm{d}\dot{L} \end{aligned} \tag{2.4}$$

The Crystal Size Distribution (CSD) at 'time' T is $N(T)^{-1}N(L)$, its probability density $f(T, L)$, and its moments

$$M_m(T) := \int_0^T f(T, L)(T - L)^m \,\mathrm{d}L = N(T)^{-1} m \int_0^T N(\dot{T}) M_{m-1}(\dot{T}) \,\mathrm{d}\dot{T} \tag{2.5}$$

The number of particles born by 'time' T is $N(T)$, and the number of particles whose 'length' does not exceed L is $N(L)$. This is the symmetry conveyed by the PBE: 'time' T and 'length' L are effectively the same thing. Like a growing child, a crystal equates how old it is with how big it is.

The CSD $N(T)^{-1}N(L)$ is the fraction of particles at 'time' T whose 'length' does not exceed L, which is the same as the fraction of particles at 'time' T born by 'time' L.

This Section determines the CSD from supersaturation $S(t)$ and volume $V(t)$, expanded about $t = \bar{t}$ as $V(t) \;= \pm \left\|\bar{V}\right\| \pm \left\|\bar{V}'\right\| (t - \bar{t})$

The partial derivatives in the second line are normally provided by an exogenous model of supersaturation, our sign convention dictating $\|\bar{\sigma}^{(n)}\| \geq 0$.

Completing the square

$$+O\left(\|\bar{\sigma}'''\| \frac{(t - \bar{t})^3}{6}\right) \tag{2.6}$$

so

$$V(t) \;= \pm \|\bar{v}\| \pm \|\bar{v}'\| \left(\pm \|\bar{\sigma}''\|\right) (t \pm \|\bar{t}\|) \pm \|\bar{v}''\| \left(\pm \|\bar{\sigma}''\|\right)^2 (t \pm \|\bar{t}\|)^2 + O\left(\left\|\bar{V}'''\right\| \frac{(t - \bar{t})^3}{6}\right)$$

$$\begin{aligned} \pm \|\bar{v}''\| &:= \pm \left\|\bar{V}''\right\| \left(\pm \|\bar{\sigma}''\|\right)^{-2} 2^{-1} \\ \pm \|\bar{v}'\| &:= \pm \left\|\bar{V}'\right\| \left(\pm \|\bar{\sigma}''\|\right)^{-1} \pm \|\bar{v}''\| \left(\pm \|\bar{\sigma}'\|\right) 2 \end{aligned} \tag{2.7}$$

Asymptotic 'time' T is determined as

$$\begin{aligned} T &= \|G\| \int_0^t (S(\dot{t}) - 1)^{|g|} \mathrm{d}\dot{t} = \|G\| \int_0^t \exp\left(|g| \ln(S(\dot{t}) - 1)\right) \mathrm{d}\dot{t} \\ &= \|G\| \int_0^t \left(1 + |g| \ln(S(\dot{t}) - 1) + O(|g|^2)\right) \mathrm{d}\dot{t} \end{aligned} \tag{2.8}$$

References

1. *NIST Digital Library of Mathematical Functions*, `https://dlmf.nist.gov/`, Release 1.2.6 of 2026-03-15, F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.