PARAMETRIZING A POPULATION BALANCE EQUATION

ROBERT A. MILTON

Department of Chemical, Materials and Biological Engineering, The University of Sheffield, Sheffield S1 3JD, United Kingdom

Abstract. Surely this cannot be that difficult.

1. Introduction

A plethora of quantities arise in this work, so we endeavour to consistently indicate their nature at a glance. All constants $\|\cdot\|, \langle\cdot\rangle, |\cdot| \in \overline{\mathbb{R}^+}$ are non-negative, and enclosed in $\|\cdot\|$ or $\langle\cdot\rangle$ without meaning (there are no norms or inner products here), or $|\cdot|$ only when taking the absolute value of $\cdot$. All italic quantities are dimensionless, to ease the unavoidable chore of tracking units. Undefined notation adheres to the Digital Library of Mathematical Functions [1].

A number $\mathrm{n}(\mathrm{t},\mathrm{L})$ (m^{-4}) of particles of length $\mathrm{L}\,(\mathrm{m})$ grow within each unit of reactor volume $\mathrm{V}\,(\mathrm{m}^3)$ as seconds $\mathrm{t}\,(\mathrm{s})$ pass. Conserving mass, the Population Balance Equation is

$$\begin{aligned} \partial_{\mathrm{t}}(\mathrm{Vn}(\mathrm{t},\mathrm{L})) &= -\mathrm{G}^+\partial_{\mathrm{L}}(\mathrm{Vn}(\mathrm{t},\mathrm{L})) \\ \mathrm{n}(0,\mathrm{L}) &= 0 \\ \mathrm{n}(\mathrm{t},0) &= \mathrm{B}^+/\mathrm{G}^+ \end{aligned} \tag{1.1}$$

where $\mathrm{B}^+\,(\mathrm{m}^{-3}\mathrm{s}^{-1})$ and $\mathrm{G}^+\,(\mathrm{m}\,\mathrm{s}^{-1})$ are particle birth (nucleation) and growth rates. These relate empirically to supersaturation $S > 1$:

$$\begin{aligned} \text{birth} \quad \mathrm{B}^+ &:= \|\mathrm{B}\|\,(S-1)^{+|b|} \\ \text{growth} \quad \mathrm{G}^+ &:= \|\mathrm{G}\|\,(S-1)^{+|g|} \end{aligned} \tag{1.2}$$

where $(\#)^+ := \max(0,\#)$ denotes the positive part of $(\#)$. In experiments, critical supersaturation $S = 1$ might occur many times, but we cannot allow this. We must insist that

$$\text{supersaturation} \quad S \in (1, \sup S] \quad \text{at all times} \quad \mathrm{t} > \mathrm{t}^* \tag{1.3}$$

for finite supersaturation $\sup S \in \mathbb{R}$ and critical time $\mathrm{t}^* \in \mathbb{R}$ at which S first exceeds 1. This is not limiting per se, but it does mean that each critical instant in an experiment theoretically begins a new experiment.

Given lengthscale $\|\mathrm{L}\|\,(\mathrm{m})$ and timescale $\|\mathrm{t}\|\,(\mathrm{s})$, dimensionless prefactors are defined as

$$\begin{aligned} \text{birth} \quad \|B\| &:= \|\mathrm{B}\|\,\|\mathrm{L}\|^3\,\|\mathrm{t}\| \\ \text{growth} \quad \|G\| &:= \|\mathrm{G}\|\,\|\mathrm{L}\|^{-1}\,\|\mathrm{t}\| \end{aligned} \tag{1.4}$$

together with dimensionless

$$\begin{aligned} \text{time} \quad t &:= \|\mathrm{t}\|^{-1}\,(\mathrm{t}-\mathrm{t}^*) \\ \text{volume} \quad V(t) &:= \|\mathrm{L}\|^{-3}\,\mathrm{V} \end{aligned} \tag{1.5}$$

This work aims to elucidate the dimensionless empirical birth and growth parameters $\|B\|, |b|, \|G\|, |g|$.

Email address: r.a.milton@sheffield.ac.uk.

Key words and phrases. Asymptotics.

This work was funded by Solomon f. Brown.

2. The Crystal Size Distribution (CSD)

The Population Balance Equation conveys a symmetry between synthetic 'time' T and 'length' L defined as

$$S(t) := \max(S, 1) \tag{2.1a}$$

$$T := \|\mathrm{L}\|^{-1} \int_0^{\mathrm{t}} \mathrm{G}^+ \,\mathrm{dt} = \int_0^t \|G\| \left(S(\dot{t}) - 1\right)^{|g|} \mathrm{d}\dot{t} \tag{2.1b}$$

$$L := T - \|\mathrm{L}\|^{-1} \mathrm{L} \tag{2.1c}$$

This symmetry manifests in the particle count $N(T)f(T, L)$ defined as

$$N(\mathbb{R}^-) := \{0\} \tag{2.2a}$$

$$N(T) := \int_0^{\mathrm{t}} \mathrm{B}^+\mathrm{V} \,\mathrm{d\dot{t}} = \int_0^t \|B\| \left(S(\dot{t}) - 1\right)^{|b|} V(\dot{t}) \,\mathrm{d}\dot{t} \tag{2.2b}$$

$$f(T, L) := N(T)^{-1} \|\mathrm{L}\|^{-1} \mathrm{Vn(t, L)} \tag{2.2c}$$

Substituting these variables in Population Balance Equation (1.1) immediately yields

$$\begin{aligned} \partial_T N(T) f(T, L) &= 0 \\ N(T) f(T, L) &= \partial_t N(T) / \partial_t T = \partial_T N(T) \end{aligned} \tag{2.3}$$

These two lines combine to establish

$$\begin{aligned} f(T, L) &= N(T)^{-1} \partial_L N(L) \\ N(T)^{-1} N(L) &= \int_0^L f(T, \dot{L}) \,\mathrm{d}\dot{L} \end{aligned} \tag{2.4}$$

The Crystal Size Distribution at 'time' T is $N(T)^{-1}N(L)$, its probability density $f(T, L)$, and its moments

$$M_m(T) := \int_0^T f(T, L)(T - L)^m \,\mathrm{d}L = N(T)^{-1} m \int_0^T N(\dot{T}) M_{m-1}(\dot{T}) \,\mathrm{d}\dot{T} \tag{2.5}$$

3. Asymptotics for small growth exponent

This Section assumes $|g| = o(1)$. Defining

$$s(t) = \ln S(t) \tag{3.1}$$

Equation (2.1b) integrates by parts to yield

$$\begin{aligned} T &= \|G\| \int_0^t (S(\dot{t}) - 1)^{|g|} \mathrm{d}\dot{t} = \|G\| \int_0^t \exp(|g| s(\dot{t})) \left(1 - \exp(-s(\dot{t}))\right)^{|g|} \mathrm{d}\dot{t} \\ &= \|G\| \sum_{k=0}^{\infty} \frac{(-t)^k t}{\Gamma(k+2)} \partial_t^k \left(\exp(|g| s(t)) \left(1 - \exp(-s(t))\right)^{|g|} \right) \\ &= \|G\| \left(S(t) - 1\right)^{|g|} t \left(1 + \sum_{j=1}^{\infty} |g|^j \sum_{k=j}^{\infty} \frac{(-t)^k \, \partial_t^{k-j} \left(s(t) + \ln\left(1 - \exp(-s(t))\right)\right)^j}{(k+1)\Gamma(k-j+1)\Gamma(j+1)} \right) \\ &= \|G\| \, t (S(t) - 1)^{|g|} \left(1 - |g| \sum_{k=1}^{\infty} \frac{(-t)^k \, \partial_t^{k-1} \left(s(t) + \ln\left(1 - \exp(-s(t))\right)\right)}{(k+1)\Gamma(k)} + O(|g|^2) \right) \end{aligned} \tag{3.2}$$

References

1. *NIST Digital Library of Mathematical Functions*, `https://dlmf.nist.gov/`, Release 1.2.6 of 2026-03-15, F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.