PARAMETRIZING A MODEL OF EFFICIENT CRYSTAL PRODUCTION

ROBERT A. MILTON

Department of Chemical, Materials and Biological Engineering, The University of Sheffield, Sheffield S1 3JD, United Kingdom

Abstract. Surely this cannot be that difficult.

1. Introduction

A plethora of quantities arise in this work, so we endeavour to consistently indicate their nature at a glance. All constants $\|\cdot\|, \langle\cdot\rangle, |\cdot| \in \overline{\mathbb{R}^+}$ are non-negative, including scale $\|\cdot\|$ and stoichiometry $\langle\cdot\rangle$ (to avoid confusion, please note that there are no norms or inner products in this work). Any other constant $\cdot$ be written as $|\cdot|$. All italic quantities are dimensionless, to ease the unavoidable chore of tracking units. Ions are written without charge superscripts (e.g. OH means OH^-) to lighten notation, as this is a work of mathematics and not chemistry. Undefined notation adheres to the Digital Library of Mathematical Functions [1].

Imagine a complexing agent A interacting with cations $\mathrm{C}_I \in \left\{\mathrm{C}_1, \ldots, \mathrm{C}_{\|I\|}\right\}$ in a reactor to form an ionic complex

$$\mathrm{P} \rightleftharpoons (\mathrm{C}_1)_{\langle 1\rangle}(\mathrm{C}_2)_{\langle 2\rangle}\cdots(\mathrm{C}_{\|I\|})_{\langle \|I\|\rangle}(\mathrm{OH})_{\langle OH\rangle} \quad \text{in stoichiometric ratios} \quad \sum_{I=1}^{\|I\|}\langle I\rangle = 1$$

Let $|\mathrm{K_P}|$ denote the association (equilibrium) constant of complex P, which crystallizes when its concentration exceeds its equilibrium solubility $|\mathrm{K_S}|$. Crystallization is indicated by the 3D supersaturation (ratio) exceeding unity

$$S^3 := |\mathrm{K_P}|\,|\mathrm{K_S}|^{-1}\,[\mathrm{OH}]^{\langle OH\rangle}\prod_{I=1}^{\|I\|}[\mathrm{C}_I]^{\langle I\rangle} > 1 \tag{1.1}$$

where $[\cdot]$ denotes the free ion concentration, as distinct from the total species concentration $\Sigma[\cdot]$ across all ions. The 1D supersaturation is S.

Crystals of P are manufactured from a steady supply of stoichiometric amounts of C_I, A and other ingredients. These flow into a cation-free initial volume $\|\mathrm{V}\|\,(\mathrm{dm}^3)$ of complexing agent at a specially chosen initial concentration $\|\mathrm{A}\|\,(\mathrm{mol\,dm}^{-3})$. Defining

$$\begin{aligned}
\text{supply rate:}\quad & |\mathrm{F}| && (\mathrm{dm^3 h^{-1}})\\
\text{complexing agent supply fraction:}\quad & |F_\mathrm{A}| < 1 &&\\
\text{complexing agent supply concentration:}\quad & \|\,\Sigma[\mathrm{A}]\| && (\mathrm{mol\,dm^{-3}})\\
\text{all cation supply fraction:}\quad & |F_\mathrm{C}| \le 1 - |F_\mathrm{A}| &&\\
\text{(individual) cation supply concentration:}\quad & \|\,\Sigma[\mathrm{C}_I]\| = \langle I\rangle\,\|\,\Sigma[\mathrm{C}]\| && (\mathrm{mol\,dm^{-3}})\\
\text{(variable) crystal yield:}\quad & \mathrm{P} && (\mathrm{mol})\\
\text{pH:}\quad & |H| &&
\end{aligned} \tag{1.2}$$

Email address: r.a.milton@sheffield.ac.uk.

Key words and phrases. Asymptotics.

This work was funded by Solomon f. Brown.

the manufacturing process amounts to

$$\begin{aligned}
\mathrm{V} &= \|\mathrm{V}\| + |\mathrm{F}|\,\mathrm{t} && (\mathrm{dm}^3)\\
\partial_\mathrm{t}\left(\mathrm{V}\,\Sigma[\mathrm{A}]\right) &= |\mathrm{F}|\,|F_\mathrm{A}|\,\|\,\Sigma[\mathrm{A}]\| && (\mathrm{mol\,h}^{-1})\\
\partial_\mathrm{t}\left(\mathrm{V}\,\Sigma[\mathrm{C}_I]\right) &= |\mathrm{F}|\,|F_\mathrm{C}|\,\langle I\rangle\,\|\,\Sigma[\mathrm{C}]\| - \langle I\rangle\partial_\mathrm{t}\mathrm{P} && (\mathrm{mol\,h}^{-1})\\
\Sigma[\mathrm{A}]_{\mathrm{t}=0} &= \|\mathrm{A}\| := |F_\mathrm{A}|\,\|\,\Sigma[\mathrm{A}]\| && (\mathrm{mol\,dm}^{-3})\\
\Sigma[\mathrm{C}_I]_{\mathrm{t}=0} &= 0 && (\mathrm{mol\,dm}^{-3})\\
\Sigma[\mathrm{P}]_{\mathrm{t}=0} &= 0 && (\mathrm{mol})
\end{aligned} \tag{1.3}$$

The specially chosen initial concentration is $\|\mathrm{A}\| := |F_\mathrm{A}|\,\|\,\Sigma[\mathrm{A}]\|\,(\mathrm{mol\,dm}^{-3})$ to keep $\Sigma[\mathrm{A}]$ constant. The all cation supply concentration is $\|\,\Sigma[\mathrm{C}]\| = \sum_{I=1}^{\|I\|}\langle I\rangle\,\|\,\Sigma[\mathrm{C}]\|\,(\mathrm{mol\,dm}^{-3})$, consisting of individual cations in stoichiometric ratios $\langle I\rangle$. The analysis which follows will admit changes of detail (changing ratios, outflow), but such considerations are best deferred: for clarity's sake alone we rigidly specify the system, relaxing unnecessary constraints only at the conclusion.

Crystal manufacture is modelled in three Sections: Supply, Saturation, and Crystallization.

2. Supply

Presumably cations are supplied in stoichiometry to avoid material waste. The resulting theoretical economy is delightful: every cation flows and precipitates identically. Non-dimensionalizing as

$$\begin{aligned}
t &:= \|\mathrm{V}\|^{-1}\,|\mathrm{F}|\,\mathrm{t}\\
V(t) &:= \|\mathrm{V}\|^{-1}\,\mathrm{V}\\
\Sigma A(t) &:= \|\mathrm{A}\|^{-1}\,\Sigma[\mathrm{A}]\\
\|\mathrm{C}\| &:= |F_\mathrm{C}|\,\|\,\Sigma[\mathrm{C}]\| && (\mathrm{mol\,dm}^{-3})\\
\Sigma C(t) &:= \langle I\rangle^{-1}\|\mathrm{C}\|^{-1}\,\Sigma[\mathrm{C}_I]\\
P(t) &:= \|\mathrm{C}\|^{-1}\,\|\mathrm{V}\|^{-1}\,\mathrm{P}
\end{aligned} \tag{2.1}$$

the dimensionless total species concentration $\Sigma C_I := \Sigma C$ is the same for all cations. Substituting Equation (2.1) in Equation (1.3) confirms the dimensionless supply solutions

$$\begin{aligned}
V(t) &= 1 + t\\
\Sigma A(t) &= 1\\
\Sigma C(t) &= \frac{t - P(t)}{1 + t}
\end{aligned} \tag{2.2}$$

In practice, mixing may be imperfect, delaying the spread of cations through the reactor volume. The spread is naturally monotonic, and we shall assume that is also linear and the same for all cations. Accordingly, the fraction of the reactor volume actually containing cations at time t is defined as

$$U(t) := \min\left(1, \frac{\|U\|\,t}{1 + t}\right) \quad \text{where} \quad \|U\| \geq |F_\mathrm{C}| \tag{2.3}$$

It is assumed that all other ingredients, including complexing agent A, mix perfectly and pervasively.

3. Saturation

Saturation is the result of aqueous complexing agent A bonding ionically with protons H and cations C_I to form

$$\begin{aligned}
&(\mathrm{A})(\mathrm{H}) && \text{with equilibrium constant } |\mathrm{K}_\mathrm{H}|.\\
&(\mathrm{C}_I)(\mathrm{A})_j && \text{with equilibrium constant } |\mathrm{K}_{Ij}|.
\end{aligned} \tag{3.1}$$

Saturation is governed by balancing masses of A and C_I across all ionic complexes

$$\begin{aligned}\Sigma[\mathrm{A}] &= [\mathrm{A}] + |\mathrm{K_H}|\,[\mathrm{A}][\mathrm{H}] + \sum_{I=1}^{\|I\|}[\mathrm{C}_I]\sum_{j=1} j\,|\mathrm{K}_{Ij}|\,[\mathrm{A}]^j \quad (\mathrm{mol\,dm^{-3}})\\ \Sigma[\mathrm{C}_I] &= [\mathrm{C}_I] + [\mathrm{C}_I]\sum_{j=1}|\mathrm{K}_{Ij}|\,[\mathrm{A}]^j \qquad\qquad (\mathrm{mol\,dm^{-3}})\end{aligned} \tag{3.2}$$

Ionic complex formation is reversibly instantaneous. This means that the saturation process has no intrinsic sense of time – the mass balance equations are algebraic, not differential – or so it appears. Time is exogenous to the saturation process, it felt only through the total species concentrations $\Sigma[\mathrm{A}]$ and $\Sigma[\mathrm{C}_I]$ resulting from the supply process.

Non-dimensionalization naturally introduces dimensionless equilibrium constants $|K_{Ij}|$ and initial free ammonia $\|a\|$ as

$$\begin{aligned}\|a\| &:= \|\mathrm{A}\|\left(1 + |\mathrm{K_H}|\ 10^{14-|H|}\right)^{-1} \qquad (\mathrm{mol\,dm^{-3}})\\ a(\tau) &:= \|a\|^{-1}\,[\mathrm{A}]\\ |K_{Ij}| &:= |\mathrm{K}_{Ij}|\,\|a\|^j = |\mathrm{K}_{Ij}|\,[\mathrm{A}]^j a(\tau)^{-j}\end{aligned} \tag{3.3}$$

The key variable linking supply to saturation S is the dimensionless free ammonia $a(\tau)$, varying with dimensionless time τ yet to be defined. From Section 2 $\Sigma[\mathrm{A}]$ is constant and $\Sigma[\mathrm{C}_I] = \langle I\rangle\,\Sigma[\mathrm{C}]$ is essentially a single function of time which drives everything relevant to saturation. So $\Sigma C\ (\mathrm{t})$ is a natural choice for the sense of time τ. But this single source of time of course affects supersaturation itself, and only really interests us in so far as it does. So the convenient measure of time is supersaturation itself

$$\begin{aligned}\tau := S - 1 &= \|S\|^{1/3}\prod_{I=1}^{\|I\|}\left(U(t)^{-1}\langle I\rangle^{-1}\,\|C\|^{-1}\,[\mathrm{C}_I]\right)^{\langle I\rangle/3} - 1\\ \text{where}\quad \|S\| &:= |\mathrm{K_P}|\,|\mathrm{K_S}|^{-1}\,[\mathrm{OH}]^{\langle OH\rangle}\,\|C\|\prod_{I=1}^{\|I\|}\langle I\rangle^{\langle I\rangle}\end{aligned} \tag{3.4}$$

The utility of this surprising choice is revealed through the dimensionless polynomials

$$\begin{aligned}\Xi_I(a(\tau)) &:= 1 + \sum_{j=1} K_{Ij}a(\tau)^j\\ \Xi(a(\tau)) &:= \prod_{I=1}^{\|I\|}\Xi_I(a(\tau))^{\langle I\rangle}\end{aligned} \tag{3.5}$$

Dimensionless mass balances translate Equation (3.2) into

$$\begin{aligned}(1-a(\tau))\,a(\tau)^{-1} &= \|I\|\,\|\mathrm{C}\|\,\|\mathrm{A}\|^{-1}\,\Sigma C(t)\,\partial_{a(\tau)}\ln\Xi(a(\tau))\\ (1+\tau)^3 &= \|S\|\ \Sigma C(t)\,U(t)^{-1}\Xi(a(\tau))^{-1}\end{aligned} \tag{3.6}$$

or

$$\begin{aligned}\ln\Xi(a(\tau)) &= \ln\|S\| + \ln\ \Sigma C(t) - \ln U(t) - 3\ln(1+\tau)\\ (1-a(\tau))\,\partial_\tau \ln a(\tau) &= \|I\|\,\|\mathrm{C}\|\,\|\mathrm{A}\|^{-1}\left(\partial_\tau\,\Sigma C(t) - \ \Sigma C(t)\partial_\tau \ln\left((1+\tau)^3U(t)\right)\right)\end{aligned} \tag{3.7}$$

References

1. *NIST Digital Library of Mathematical Functions*, `https://dlmf.nist.gov/`, Release 1.2.6 of 2026-03-15, F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.