

# 사다리꼴 방법에서의 오차


(Error in the Trapezoidal Rule)

# Error in the Trapezoidal Rule

# Error in the Trapezoidal Rule




# Error in the Trapezoidal Rule




## Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$




## Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$




## Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$




## Error in the Trapezoidal Rule






## Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$




## Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$




## Error in the Trapezoidal Rule


$$|f''(x)| \leq M, \quad x \in [a, b]$$



## Error in the Trapezoidal Rule





## Error in the Trapezoidal Rule





## Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$





## Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$



# Error in the Trapezoidal Rule



$$|f''(x)| \leq M, \quad x \in [a, b]$$







# Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$





# Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$





# Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$





# Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$





# Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$




# Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$




# Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$





$$\int_{x_i}^{x_{i+1}} \left\{ -\frac{M}{2} (x - x_i)(x - x_{i+1}) \right\} dx$$



# Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$





$$\int_{x_i}^{x_{i+1}} \left\{ -\frac{M}{2} (x - x_i)(x - x_{i+1}) \right\} dx = M \frac{(x_{i+1} - x_i)^3}{12}$$



# Error in the Trapezoidal Rule


$$|f''(x)| \leq M, \quad x \in [a, b]$$



$$\int_{x_i}^{x_{i+1}} \left\{ -\frac{M}{2} (x - x_i)(x - x_{i+1}) \right\} dx = M \frac{(x_{i+1} - x_i)^3}{12} = M \frac{(b - a)^3}{12n^3}$$


# Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$



$$T_n = \sum_{i=1}^n \left\{ \Delta x \times \frac{f(x_{i-1}) + f(x_i)}{2} \right\}$$

$$\int_{x_i}^{x_{i+1}} \left\{ -\frac{M}{2} (x - x_i)(x - x_{i+1}) \right\} dx = M \frac{(x_{i+1} - x_i)^3}{12} = M \frac{(b - a)^3}{12n^3}$$





$$\{m(x - x_i)(x - x_{i+1})\}'' = -M$$

$$2m = -M$$

$$m = -\frac{M}{2}$$

# Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$



$$\{m(x - x_i)(x - x_{i+1})\}'' = -M$$

$$2m = -M$$

$$m = -\frac{M}{2}$$


$$T_n = \sum_{i=1}^n \left\{ \Delta x \times \frac{f(x_{i-1}) + f(x_i)}{2} \right\}$$

$$\int_{x_i}^{x_{i+1}} \left\{ -\frac{M}{2} (x - x_i)(x - x_{i+1}) \right\} dx = M \frac{(x_{i+1} - x_i)^3}{12} = M \frac{(b - a)^3}{12n^3}$$

$$\therefore \left| \int_a^b f(x) dx - T_n \right| \leq M \frac{(b - a)^3}{12n^2}$$

# Error in the Trapezoidal Rule

$$|f''(x)| \leq M, \quad x \in [a, b]$$



$$T_n = \sum_{i=1}^n \left\{ \Delta x \times \frac{f(x_{i-1}) + f(x_i)}{2} \right\}$$

$$\int_{x_i}^{x_{i+1}} \left\{ -\frac{M}{2} (x - x_i)(x - x_{i+1}) \right\} dx = M \frac{(x_{i+1} - x_i)^3}{12} = M \frac{(b - a)^3}{12n^3}$$

$$\therefore \left| \int_a^b f(x) dx - T_n \right| \leq M \frac{(b - a)^3}{12n^2}$$

