

실수에서의 코시 슈바르츠 부등식

(Cauchy Schwarz inequality in \mathbb{R})

$$a_i, b_i \in \mathbb{R}$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right)$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) .$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right)$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

$a_i, b_i \in \mathbb{R}$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

$a_i, b_i \in \mathbb{R}$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$,

$a_i, b_i \in \mathbb{R}$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

$a_i, b_i \in \mathbb{R}$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$a_i, b_i \in \mathbb{R}$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$t \in \mathbb{R}$,

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0$$

$a_i, b_i \in \mathbb{R}$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right)$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \left(\sum_{i=1}^n b_i^2 \right) \leq 0$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \leq 0$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right)$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \leq$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \leq 0$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \leq 0 \quad (\because$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \leq 0 \quad \left(\because \sum_{i=1}^n a_i^2 > 0, \right)$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \leq 0 \quad \left(\because \sum_{i=1}^n a_i^2 > 0, D/4 \leq 0 \right)$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \leq 0 \quad \left(\because \sum_{i=1}^n a_i^2 > 0, D/4 \leq 0 \right)$$

\therefore

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \leq 0 \quad \left(\because \sum_{i=1}^n a_i^2 > 0, D/4 \leq 0 \right)$$

$$\therefore \left(\sum_{i=1}^n a_i^2 \right)$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \leq 0 \quad \left(\because \sum_{i=1}^n a_i^2 > 0, D/4 \leq 0 \right)$$

$$\therefore \left(\sum_{i=1}^n a_i^2 \right) \cdot$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \leq 0 \quad \left(\because \sum_{i=1}^n a_i^2 > 0, D/4 \leq 0 \right)$$

$$\therefore \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right)$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \leq 0 \quad \left(\because \sum_{i=1}^n a_i^2 > 0, D/4 \leq 0 \right)$$

$$\therefore \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \leq 0 \quad \left(\because \sum_{i=1}^n a_i^2 > 0, D/4 \leq 0 \right)$$

$$\therefore \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \leq 0 \quad \left(\because \sum_{i=1}^n a_i^2 > 0, D/4 \leq 0 \right)$$

$$\therefore \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

The two sides are equal if and only if (a_1, \dots, a_n) and (b_1, \dots, b_n) are linearly dependent. ($\because (1)$)

$$a_i, b_i \in \mathbb{R}$$

$$\left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

proof.

If $(a_1, \dots, a_n) = (0, \dots, 0)$, then it is trivial.

Assume $(a_1, \dots, a_n) \neq (0, \dots, 0)$

$$t \in \mathbb{R}, \sum_{i=1}^n (a_i t - b_i)^2 \geq 0 \quad (\because a_i t - b_i \in \mathbb{R}) \dots \dots (1)$$

$$\left(\sum_{i=1}^n a_i^2 \right) t^2 - 2 \left(\sum_{i=1}^n a_i b_i \right) t + \left(\sum_{i=1}^n b_i^2 \right) \geq 0$$

$$\left(\sum_{i=1}^n a_i b_i \right)^2 - \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \leq 0 \quad \left(\because \sum_{i=1}^n a_i^2 > 0, D/4 \leq 0 \right)$$

$$\therefore \left(\sum_{i=1}^n a_i^2 \right) \cdot \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

The two sides are equal if and only if (a_1, \dots, a_n) and (b_1, \dots, b_n) are linearly dependent. ($\because (1)$)

YouTube: https://youtu.be/U_TPopIZR8M

Click or paste URL into the URL search bar, and you can see a picture moving.