

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

양수에 대한 산술, 기하, 조화평균에 대한 부등식
(Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers)

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

► Start

► End

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Start ▶ End $x_1 > 0, \dots, x_n > 0$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Start

▶ End

$$x_1 > 0, \dots, x_n > 0$$

$$\frac{x_1 + \dots + x_n}{n}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Start ▶ End $x_1 > 0, \dots, x_n > 0$

$$\frac{x_1 + \dots + x_n}{n} \geq$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Start ▶ End $x_1 > 0, \dots, x_n > 0$

$$\frac{x_1 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \times \dots \times x_n}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Start ▶ End $x_1 > 0, \dots, x_n > 0$

$$\frac{x_1 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \times \dots \times x_n} \geq$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Start ▶ End $x_1 > 0, \dots, x_n > 0$

$$\frac{x_1 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \times \dots \times x_n} \geq \frac{n}{\frac{1}{x_1} + \dots + \frac{1}{x_n}}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Start ▶ End $x_1 > 0, \dots, x_n > 0$

$$\frac{x_1 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \times \dots \times x_n} \geq \frac{n}{\frac{1}{x_1} + \dots + \frac{1}{x_n}}$$

▶ Proof of first inequality

▶ Proof of second inequality

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

[▶ Home](#)[▶ Start](#)[▶ End](#)

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality
If $n = 2$.

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

If $n = 2$.

$$\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

If $n = 2$.

$$\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 = \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

If $n = 2$.

$$\begin{aligned}\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 &= \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2 \\ &= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

If $n = 2$.

$$\begin{aligned}\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 &= \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2 \\ &= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4} \\ &= \frac{(x_1 - x_2)^2}{4}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

If $n = 2$.

$$\begin{aligned}\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 &= \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2 \\ &= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4} \\ &= \frac{(x_1 - x_2)^2}{4} \geq\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

If $n = 2$.

$$\begin{aligned}\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 &= \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2 \\ &= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4} \\ &= \frac{(x_1 - x_2)^2}{4} \geq 0\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

If $n = 2$.

$$\begin{aligned}\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 &= \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2 \\ &= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4} \\ &= \frac{(x_1 - x_2)^2}{4} \geq 0\end{aligned}$$

$$\therefore \frac{x_1 + x_2}{2}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

If $n = 2$.

$$\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 = \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2$$

$$= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4}$$

$$= \frac{(x_1 - x_2)^2}{4} \geq 0$$

$$\therefore \frac{x_1 + x_2}{2} \geq$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

If $n = 2$.

$$\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 = \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2$$

$$= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4}$$

$$= \frac{(x_1 - x_2)^2}{4} \geq 0$$

$$\therefore \frac{x_1 + x_2}{2} \geq \sqrt{x_1 x_2}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

If $n = 2$.

$$\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 = \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2$$

$$= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4}$$

$$= \frac{(x_1 - x_2)^2}{4} \geq 0$$

$$\therefore \frac{x_1 + x_2}{2} \geq \sqrt{x_1 x_2}$$

$$n = 2$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

If $n = 2$.

$$\begin{aligned}\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 &= \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2 \\ &= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4} \\ &= \frac{(x_1 - x_2)^2}{4} \geq 0\end{aligned}$$

$$\therefore \frac{x_1 + x_2}{2} \geq \sqrt{x_1 x_2}$$

$n = 2$ is true.

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Assume

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Assume $n = 2^{k-1}$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Assume $n = 2^{k-1}$ is true.

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\frac{x_1 + \cdots + x_{2^k}}{2^k}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\frac{x_1 + \cdots + x_{2^k}}{2^k} =$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\frac{x_1 + \cdots + x_{2^k}}{2^k} = \frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\frac{x_1 + \cdots + x_{2^k}}{2^k} = \frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} +$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\frac{x_1 + \cdots + x_{2^k}}{2^k} = \frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\frac{x_1 + \cdots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\frac{x_1 + \cdots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2}$$
$$\geq$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\begin{aligned}\frac{x_1 + \cdots + x_{2^k}}{2^k} &= \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2} \\ &\geq \frac{2^{k-1} \sqrt{x_1 \cdots x_{2^{k-1}}}}{2}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\begin{aligned}\frac{x_1 + \cdots + x_{2^k}}{2^k} &= \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2} \\ &\geq 2^{k-1} \sqrt{x_1 \cdots x_{2^{k-1}}} +\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\begin{aligned}\frac{x_1 + \cdots + x_{2^k}}{2^k} &= \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2} \\ &\geq \frac{2^{k-1}\sqrt{x_1 \cdots x_{2^{k-1}}} + 2^{k-1}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\begin{aligned}\frac{x_1 + \cdots + x_{2^k}}{2^k} &= \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2} \\ &\geq \frac{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} + {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\begin{aligned}\frac{x_1 + \cdots + x_{2^k}}{2^k} &= \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2} \\ &\geq \frac{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} + {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2} \\ &\geq\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\begin{aligned}\frac{x_1 + \cdots + x_{2^k}}{2^k} &= \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2} \\ &\geq \frac{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} + {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2} \\ &\geq \sqrt{\phantom{x_1 + \cdots + x_{2^k}}}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\begin{aligned}\frac{x_1 + \cdots + x_{2^k}}{2^k} &= \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2} \\ &\geq \frac{2^{k-1}\sqrt{x_1 \cdots x_{2^{k-1}}} + 2^{k-1}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2} \\ &\geq \sqrt{2^{k-1}\sqrt{x_1 \cdots x_{2^{k-1}}}}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\begin{aligned}\frac{x_1 + \cdots + x_{2^k}}{2^k} &= \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2} \\ &\geq \frac{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} + {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2} \\ &\geq \sqrt{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} \cdot {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\begin{aligned}\frac{x_1 + \cdots + x_{2^k}}{2^k} &= \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2} \\ &\geq \frac{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} + {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2} \\ &\geq \sqrt{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} \cdot {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}} \\ &= \end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\begin{aligned}\frac{x_1 + \cdots + x_{2^k}}{2^k} &= \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2} \\ &\geq \frac{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} + {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2} \\ &\geq \sqrt{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} \cdot {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}} \\ &= \sqrt[2^k]{x_1 \cdots x_{2^k}}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\begin{aligned}\frac{x_1 + \cdots + x_{2^k}}{2^k} &= \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2} \\ &\geq \frac{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} + {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2} \\ &\geq \sqrt{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} \cdot {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}} \\ &= \sqrt[2^k]{x_1 \cdots x_{2^k}} \\ \therefore \frac{x_1 + \cdots + x_{2^k}}{2^k}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\begin{aligned}\frac{x_1 + \cdots + x_{2^k}}{2^k} &= \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2} \\ &\geq \frac{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} + {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2} \\ &\geq \sqrt{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} \cdot {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}} \\ &= \sqrt[2^k]{x_1 \cdots x_{2^k}} \\ \therefore \frac{x_1 + \cdots + x_{2^k}}{2^k} &\geq\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\begin{aligned}\frac{x_1 + \cdots + x_{2^k}}{2^k} &= \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2} \\ &\geq \frac{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} + {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2} \\ &\geq \sqrt{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} \cdot {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}} \\ &= \sqrt[2^k]{x_1 \cdots x_{2^k}} \\ \therefore \frac{x_1 + \cdots + x_{2^k}}{2^k} &\geq \sqrt[2^k]{x_1 \cdots x_{2^k}}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\frac{x_1 + \cdots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2}$$

$$\geq \frac{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} + {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2}$$

$$\geq \sqrt{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} \cdot {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}$$

$$= \sqrt[2^k]{x_1 \cdots x_{2^k}}$$

$$\therefore \frac{x_1 + \cdots + x_{2^k}}{2^k} \geq \sqrt[2^k]{x_1 \cdots x_{2^k}}$$

$$n = 2^k$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Assume $n = 2^{k-1}$ is true.

$$\frac{x_1 + \cdots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \cdots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \cdots + x_{2^k}}{2^{k-1}}}{2}$$

$$\geq \frac{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} + {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2}$$

$$\geq \sqrt{{}^{2^{k-1}}\sqrt{x_1 \cdots x_{2^{k-1}}} \cdot {}^{2^{k-1}}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}$$

$$= \sqrt[2^k]{x_1 \cdots x_{2^k}}$$

$$\therefore \frac{x_1 + \cdots + x_{2^k}}{2^k} \geq \sqrt[2^k]{x_1 \cdots x_{2^k}}$$

$n = 2^k$ is true.

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Let $m = 2^l$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Let $m = 2^l$ such that

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Let $m = 2^l$ such that $n < m$.

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Let $m = 2^l$ such that $n < m$.

α

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\alpha =$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\alpha = \frac{x_1 + \cdots + x_n}{n}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\alpha = \frac{x_1 + \cdots + x_n}{n} =$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\alpha = \frac{x_1 + \cdots + x_n}{n} = \frac{m}{n}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\alpha = \frac{x_1 + \cdots + x_n}{n} = \frac{m}{n}(x_1 + \cdots + x_n)$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\alpha = \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\alpha = \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m}$$

=

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n}{m}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + \underbrace{}_{\text{repeated } m-n \text{ times}}}{m}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + \frac{m-n}{n} \cdot 0}{m}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)} \\ &= \end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\ &= \end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\ &= x_1 + \cdots + x_n\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\ &= x_1 + \cdots + x_n + \end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\ &= x_1 + \cdots + x_n + (m-n)\alpha\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + (m-n)\alpha}{m}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + (m-n)\alpha}{m} \\ &= x_1 + \cdots + x_n\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + (m-n)\alpha}{m} \\ &= x_1 + \cdots + x_n + \end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + (m-n)\alpha}{m} \\ &= x_1 + \cdots + x_n + \underbrace{\alpha + \cdots + \alpha}_{m-n}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\ &= \frac{x_1 + \cdots + x_n + (m-n)\alpha}{m} \\ &= x_1 + \cdots + x_n + \underbrace{\alpha + \cdots + \alpha}_{m-n}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\&= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\&= \frac{x_1 + \cdots + x_n + (m-n)\alpha}{m} \\&= \frac{x_1 + \cdots + x_n + \overbrace{\alpha + \cdots + \alpha}^{m-n}}{m}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\&= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\&= \frac{x_1 + \cdots + x_n + (m-n)\alpha}{m} \\&= \frac{x_1 + \cdots + x_n + \overbrace{\alpha + \cdots + \alpha}^{m-n}}{m} \geq\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\&= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\&= \frac{x_1 + \cdots + x_n + (m-n)\alpha}{m} \\&= \frac{x_1 + \cdots + x_n + \overbrace{\alpha + \cdots + \alpha}^{m-n}}{m} \geq \sqrt[m]{x_1 \cdots x_n \alpha^{m-n}}\end{aligned}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\&= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\&= \frac{x_1 + \cdots + x_n + (m-n)\alpha}{m} \\&= \frac{x_1 + \cdots + x_n + \overbrace{\alpha + \cdots + \alpha}^{m-n}}{m} \geq \sqrt[m]{x_1 \cdots x_n \alpha^{m-n}}\end{aligned}$$

$$\alpha^m \geq x_1 \cdots x_n \alpha^{m-n},$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\&= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\&= \frac{x_1 + \cdots + x_n + (m-n)\alpha}{m} \\&= \frac{x_1 + \cdots + x_n + \overbrace{\alpha + \cdots + \alpha}^{m-n}}{m} \geq \sqrt[m]{x_1 \cdots x_n \alpha^{m-n}}\end{aligned}$$

$$\alpha^m \geq x_1 \cdots x_n \alpha^{m-n}, \quad \alpha^n \geq x_1 \cdots x_n,$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\&= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\&= \frac{x_1 + \cdots + x_n + (m-n)\alpha}{m} \\&= \frac{x_1 + \cdots + x_n + \overbrace{\alpha + \cdots + \alpha}^{m-n}}{m} \geq \sqrt[m]{x_1 \cdots x_n \alpha^{m-n}}\end{aligned}$$

$$\alpha^m \geq x_1 \cdots x_n \alpha^{m-n}, \quad \alpha^n \geq x_1 \cdots x_n, \quad \alpha \geq \sqrt[n]{x_1 \cdots x_n}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of first inequality

Let $m = 2^l$ such that $n < m$.

$$\begin{aligned}\alpha &= \frac{x_1 + \cdots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \cdots + x_n)}{m} \\&= \frac{x_1 + \cdots + x_n + \frac{m-n}{n}(x_1 + \cdots + x_n)}{m} \\&= \frac{x_1 + \cdots + x_n + (m-n)\alpha}{m} \\&= \frac{x_1 + \cdots + x_n + \overbrace{\alpha + \cdots + \alpha}^{m-n}}{m} \geq \sqrt[m]{x_1 \cdots x_n \alpha^{m-n}}\end{aligned}$$

$$\alpha^m \geq x_1 \cdots x_n \alpha^{m-n}, \quad \alpha^n \geq x_1 \cdots x_n, \quad \alpha \geq \sqrt[n]{x_1 \cdots x_n}$$

$$\therefore \frac{x_1 + \cdots + x_n}{n} \geq \sqrt[n]{x_1 \cdots x_n} \quad (n \text{ is true.})$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of second inequality

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of second inequality

$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of second inequality

$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n} \geq$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of second inequality

$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n} \geq \sqrt[n]{\frac{1}{x_1} \cdots \frac{1}{x_n}}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of second inequality

$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n} \geq \sqrt[n]{\frac{1}{x_1} \cdots \frac{1}{x_n}}$$

$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of second inequality

$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n} \geq \sqrt[n]{\frac{1}{x_1} \cdots \frac{1}{x_n}}$$

$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n} \geq$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of second inequality

$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n} \geq \sqrt[n]{\frac{1}{x_1} \cdots \frac{1}{x_n}}$$

$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n} \geq \frac{1}{\sqrt[n]{x_1 \cdots x_n}}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

- Proof of second inequality

$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n} \geq \sqrt[n]{\frac{1}{x_1} \cdots \frac{1}{x_n}}$$

$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n} \geq \frac{1}{\sqrt[n]{x_1 \cdots x_n}}$$

$$\therefore \sqrt[n]{x_1 \times \cdots \times x_n} \geq$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of second inequality

$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n} \geq \sqrt[n]{\frac{1}{x_1} \cdots \frac{1}{x_n}}$$

$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n} \geq \frac{1}{\sqrt[n]{x_1 \cdots x_n}}$$

$$\therefore \sqrt[n]{x_1 \times \cdots \times x_n} \geq \frac{n}{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

▶ Home

▶ Start

▶ End

• Proof of second inequality

$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n} \geq \sqrt[n]{\frac{1}{x_1} \cdots \frac{1}{x_n}}$$

$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n} \geq \frac{1}{\sqrt[n]{x_1 \cdots x_n}}$$

$$\therefore \sqrt[n]{x_1 \times \cdots \times x_n} \geq \frac{n}{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}$$

Inequality of Arithmetic and Geometric and Harmonic Means for n positive real numbers

Github:

<https://min7014.github.io/math20191104001.html>

Click or paste URL into the URL search bar,
and you can see a picture moving.