양수에 대한 산술, 기하, 조화평균에 대한 부등식 (Inequality of Arithmetic and Geometric and Harmonic Means for *n* positive numbers)



Start Ind 
$$x_1 > 0, \ldots, x_n > 0$$

Start Find 
$$x_1>0,\ldots,x_n>0$$
 
$$\frac{x_1+\cdots+x_n}{n}$$

Start Find 
$$x_1>0,\ldots,x_n>0$$
 
$$\frac{x_1+\cdots+x_n}{n}\geq$$

Start Find 
$$x_1>0,\ldots,x_n>0$$
 
$$\frac{x_1+\cdots+x_n}{n}\geq \sqrt[n]{x_1 imes\cdots imes x_n}$$

Start Find 
$$x_1>0,\ldots,x_n>0$$
 
$$\frac{x_1+\cdots+x_n}{n}\geq \sqrt[n]{x_1 imes\cdots imes x_n}\geq$$

Start Find 
$$x_1>0,\ldots,x_n>0$$
 
$$\frac{x_1+\cdots+x_n}{n}\geq \sqrt[n]{x_1 imes\cdots imes x_n}\geq \frac{n}{\dfrac{1}{x_1}+\cdots+\dfrac{1}{x_n}}$$

Start 
$$lacksymbol{\cdot}$$
 End  $x_1>0,\ldots,x_n>0$ 

$$\frac{x_1+\cdots+x_n}{n}\geq \sqrt[n]{x_1\times\cdots\times x_n}\geq \frac{n}{\frac{1}{x_1}+\cdots+\frac{1}{x_n}}$$

- ▶ Proof of first inequality
- ▶ Proof of second inequality







• Proof of first inequality If n = 2.



If 
$$n = 2$$
.  
 $\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2$ 



If 
$$n = 2$$
.  
 $\left(\frac{x_1 + x_2}{2}\right)^2 - \left(\sqrt{x_1 x_2}\right)^2 = \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2$ 



If 
$$n = 2$$
.  

$$\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 = \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2$$

$$= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4}$$

#### → Home → Start → End

If 
$$n = 2$$
.  

$$\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 = \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2$$

$$= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4}$$

$$= \frac{(x_1 - x_2)^2}{4}$$

#### → Home → Start → End

If 
$$n = 2$$
.  

$$\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 = \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2$$

$$= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4}$$

$$= \frac{(x_1 - x_2)^2}{4} \ge$$

#### → Home → Start → End

If 
$$n = 2$$
.  

$$\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 = \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2$$

$$= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4}$$

$$= \frac{(x_1 - x_2)^2}{4} \ge 0$$

#### → Home → Start → End

If 
$$n = 2$$
.
$$\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 = \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2$$

$$= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4}$$

$$= \frac{(x_1 - x_2)^2}{4} \ge 0$$

$$\therefore \frac{x_1 + x_2}{2}$$

#### → Home → Start → End

If 
$$n = 2$$
.
$$\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 = \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2$$

$$= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4}$$

$$= \frac{(x_1 - x_2)^2}{4} \ge 0$$

$$\therefore \frac{x_1 + x_2}{2} \ge$$

#### → Home → Start → End

If 
$$n = 2$$
.
$$\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 = \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2$$

$$= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4}$$

$$= \frac{(x_1 - x_2)^2}{4} \ge 0$$

$$\therefore \frac{x_1 + x_2}{2} \ge \sqrt{x_1 x_2}$$

#### → Home → Start → End

If 
$$n = 2$$
.
$$\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 = \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2$$

$$= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4}$$

$$= \frac{(x_1 - x_2)^2}{4} \ge 0$$

$$\therefore \frac{x_1 + x_2}{2} \ge \sqrt{x_1 x_2}$$

$$n = 2$$

n=2 is true.

#### → Home → Start → End

If 
$$n = 2$$
.
$$\left(\frac{x_1 + x_2}{2}\right)^2 - (\sqrt{x_1 x_2})^2 = \frac{x_1^2 + 2x_1 x_2 + x_2^2}{4} - x_1 x_2$$

$$= \frac{x_1^2 - 2x_1 x_2 + x_2^2}{4}$$

$$= \frac{(x_1 - x_2)^2}{4} \ge 0$$

$$\therefore \frac{x_1 + x_2}{2} \ge \sqrt{x_1 x_2}$$







• Proof of first inequality Assume



• Proof of first inequality Assume  $n = 2^{k-1}$ 



• Proof of first inequality Assume  $n = 2^{k-1}$  is true.



• Proof of first inequality Assume  $n = 2^{k-1}$  is true.

$$\frac{x_1+\cdots+x_{2^k}}{2^k}$$



• Proof of first inequality Assume  $n = 2^{k-1}$  is true.

$$\frac{x_1+\cdots+x_{2^k}}{2^k} =$$



Assume 
$$n = 2^{k-1}$$
 is true.  

$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}}$$



Assume 
$$n = 2^{k-1}$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \dots$$



Proof of first inequality

Assume 
$$n = 2^{k-1}$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}$$



Proof of first inequality

Assume 
$$n = 2^{k-1}$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}}{2}$$





Proof of first inequality

Assume 
$$n = 2^{k-1}$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}}{2}$$



Proof of first inequality

Assume 
$$n = 2^{k-1}$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}}{2}$$

$$\geq \frac{x_1 + \dots + x_{2^k}}{2^k}$$



Proof of first inequality

Assume 
$$n = 2^{k-1}$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}}{2}$$

$$\geq \frac{x_1 + \dots + x_{2^k}}{2^k}$$



Proof of first inequality

Assume 
$$h = 2$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}}{2}$$

$$\geq \frac{x_1 + \dots + x_{2^k}}{2^k}$$



• Proof of first inequality

Assume 
$$n = 2$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}}{2}$$

$$\geq \frac{\frac{z^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}} + \frac{z^{k-1}\sqrt{x_{2^{k-1}+1} \dots x_{2^k}}}{2}}{2}$$





Proof of first inequality

Assume 
$$n = 2$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}}{2}$$

$$\geq \frac{\frac{z^{k-1}\sqrt{x_1 \cdots x_{2^{k-1}}} + \frac{z^{k-1}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2}}{2}$$

$$\geq \frac{z^{k-1}\sqrt{x_1 \cdots x_{2^{k-1}}} + \frac{z^{k-1}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2}}{2}$$



• Proof of first inequality

Assume 
$$n = 2^{k-1}$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k}}}{2}$$

$$\geq \frac{2^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}} + 2^{k-1}\sqrt{x_{2^{k-1}+1} \dots x_{2^k}}}{2}$$

$$\geq \sqrt{}$$



Proof of first inequality

Assume 
$$n = 2$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}}{2}$$

$$\geq \frac{\frac{2^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}} + \frac{2^{k-1}\sqrt{x_{2^{k-1}+1} + \dots + x_{2^k}}}{2^k}}{2}$$

$$\geq \sqrt{\frac{2^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}}}{2^k}}$$



Proof of first inequality

Assume 
$$n = 2^{k-1}$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k}}}{2}$$

$$\geq \frac{2^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}} + 2^{k-1}\sqrt{x_{2^{k-1}+1} \dots x_{2^k}}}{2}$$

$$\geq \sqrt{2^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}}} 2^{k-1}\sqrt{x_{2^{k-1}+1} \dots x_{2^k}}$$



Proof of first inequality

Assume 
$$n = 2^{k-1}$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}}{2}$$

$$\geq \frac{\frac{z^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}} + \frac{z^{k-1}\sqrt{x_{2^{k-1}+1} \dots x_{2^k}}}{2}}{2}$$

$$\geq \sqrt{\frac{z^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}} + \frac{z^{k-1}\sqrt{x_{2^{k-1}+1} \dots x_{2^k}}}{2^k}}{2}}$$

$$= \frac{z^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}} + \frac{z^{k-1}\sqrt{x_{2^{k-1}+1} \dots x_{2^k}}}{2^k}}{2^k}$$



#### Proof of first inequality

Assume 
$$n = 2^{k-1}$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}}{2}$$

$$\geq \frac{\frac{2^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}} + \frac{2^{k-1}\sqrt{x_{2^{k-1}+1} \dots x_{2^k}}}{2}}{2}$$

$$\geq \sqrt{\frac{2^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}}}{2^{k-1}\sqrt{x_{2^{k-1}+1} \dots x_{2^k}}}}$$

$$= \frac{2^k\sqrt{x_1 \dots x_{2^k}}}{2^k}$$

#### → Start → End ▶ Home

Assume 
$$n = 2^{k-1}$$
 is true.

Assume 
$$n = 2^{n-1}$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}}{2}$$

$$\geq \frac{\frac{2^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}} + \frac{2^{k-1}\sqrt{x_{2^{k-1}+1} \dots x_{2^k}}}{2^k}}{2}$$

$$\geq \sqrt{\frac{2^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}}}{2^k}}$$

$$= \frac{2^k\sqrt{x_1 \dots x_{2^k}}}{2^k}$$

$$\therefore \frac{x_1 + \dots + x_{2^k}}{2^k}$$

#### ▶ Home → Start → End

Assume 
$$n = 2^{k-1}$$
 is true.

Assume 
$$n = 2^{n-1}$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}}{2}$$

$$\geq \frac{\frac{2^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}} + \frac{2^{k-1}\sqrt{x_{2^{k-1}+1} \dots x_{2^k}}}{2^k}}{2}$$

$$\geq \sqrt{\frac{2^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}}}{2^k}}$$

$$= \frac{2^k\sqrt{x_1 \dots x_{2^k}}}{2^k}$$

$$\therefore \frac{x_1 + \dots + x_{2^k}}{2^k} \geq$$

#### → Home → Start → End

Assume 
$$n = 2^{k-1}$$
 is true.

$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}}{2}$$

$$\geq \frac{\frac{z^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}} + \frac{z^{k-1}\sqrt{x_{2^{k-1}+1} \dots x_{2^k}}}{2^k}}{2}$$

$$\geq \sqrt{\frac{z^{k-1}\sqrt{x_1 \dots x_{2^{k-1}}}}{2^k}}$$

$$= \frac{z^k\sqrt{x_1 \dots x_{2^k}}}{2^k}$$

$$\therefore \frac{x_1 + \dots + x_{2^k}}{2^k} \geq \frac{z^k\sqrt{x_1 \dots x_{2^k}}}{2^k}$$

#### → Home → Start → End

Proof of first inequality

Assume 
$$n = 2$$
 is true.
$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}}{2}$$

$$\geq \frac{\frac{z^{k-1}\sqrt{x_1 \cdots x_{2^{k-1}}} + \frac{z^{k-1}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2^k}}{2}$$

$$\geq \sqrt{\frac{z^{k-1}\sqrt{x_1 \cdots x_{2^{k-1}}}}{2^k}}$$

$$= \frac{z^k\sqrt{x_1 \cdots x_{2^k}}}{2^k}$$

$$\therefore \frac{x_1 + \dots + x_{2^k}}{2^k} \geq \frac{z^k\sqrt{x_1 \cdots x_{2^k}}}{2^k}$$

$$\therefore n = 2^k$$

#### ► Home ► Start ► End

Proof of first inequality

$$\frac{x_1 + \dots + x_{2^k}}{2^k} = \frac{\frac{x_1 + \dots + x_{2^{k-1}}}{2^{k-1}} + \frac{x_{2^{k-1}+1} + \dots + x_{2^k}}{2^{k-1}}}{2}$$

$$\geq \frac{\frac{z^{k-1}\sqrt{x_1 \cdots x_{2^{k-1}}} + \frac{z^{k-1}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2^k}}{2}$$

$$\geq \sqrt{\frac{z^{k-1}\sqrt{x_1 \cdots x_{2^{k-1}}} + \frac{z^{k-1}\sqrt{x_{2^{k-1}+1} \cdots x_{2^k}}}{2^k}}}{2}$$

$$= \frac{z^k\sqrt{x_1 \cdots x_{2^k}}}{2^k}$$

$$\therefore \frac{x_1 + \dots + x_{2^k}}{2^k} \geq \frac{z^k\sqrt{x_1 \cdots x_{2^k}}}{2^k}$$

$$n = 2^k \text{ is true}$$







• Proof of first inequality Let  $m = 2^l$ 







• Proof of first inequality Let  $m = 2^l$  such that n < m.

 $\alpha$ 



```
\alpha =
```



$$\alpha = \frac{x_1 + \cdots + x_n}{n}$$



$$\alpha = \frac{x_1 + \cdots + x_n}{n} =$$



• Proof of first inequality Let  $m = 2^l$  such that n < m.  $\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{m}{n}$ 



• Proof of first inequality Let  $m = 2^l$  such that n < m.  $\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{m}{n} (x_1 + \dots + x_n)$ 



• Proof of first inequality Let  $m = 2^{l}$  such that n < m.  $\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$ 



=

Let 
$$m = 2^l$$
 such that  $n < m$ .  

$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$



Let 
$$m = 2^l$$
 such that  $n < m$ .  

$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n}{n}$$





Let 
$$m = 2^l$$
 such that  $n < m$ .  

$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}}{n}$$



Let 
$$m = 2^l$$
 such that  $n < m$ .  

$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{n}$$



Let 
$$m = 2^{l}$$
 such that  $n < m$ .
$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{m}$$



Let 
$$m = 2^{l}$$
 such that  $n < m$ .
$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{m}$$



Let 
$$m = 2^{l}$$
 such that  $n < m$ .
$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n}{m}$$



Let 
$$m = 2^{l}$$
 such that  $n < m$ .
$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{m}$$



Let 
$$m = 2^l$$
 such that  $n < m$ .  

$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{x_1 + \dots + x_n + (m - n)\alpha}$$



Let 
$$m = 2^l$$
 such that  $n < m$ .  

$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m}{n}(x_1 + \dots + x_n)}{m}$$



Let 
$$m = 2^l$$
 such that  $n < m$ .
$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{\frac{m}{n}}$$

$$= \frac{x_1 + \dots + x_n + (m - n)\alpha}{m}$$

$$= \frac{x_1 + \dots + x_n}{m}$$





Let 
$$m = 2^{l}$$
 such that  $n < m$ .

$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{\frac{m}{n}}$$

$$= \frac{x_1 + \dots + x_n + (m - n)\alpha}{m}$$

$$= x_1 + \dots + x_n + \overbrace{\alpha + \dots + \alpha}^{m - n}$$



Let 
$$m = 2^{l}$$
 such that  $n < m$ .

$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{\frac{m}{n}}$$

$$= \frac{x_1 + \dots + x_n + (m - n)\alpha}{m}$$

$$= x_1 + \dots + x_n + \overbrace{\alpha + \dots + \alpha}^{m - n}$$



Let 
$$m = 2^{l}$$
 such that  $n < m$ .
$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + (m - n)\alpha}{m}$$

$$= \frac{x_1 + \dots + x_n + \overbrace{\alpha + \dots + \alpha}^{m - n}}{m}$$



Let 
$$m = 2^{l}$$
 such that  $n < m$ .
$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + (m - n)\alpha}{m}$$

$$= \frac{x_1 + \dots + x_n + \overbrace{\alpha + \dots + \alpha}^{m - n}}{m} \ge$$



Let 
$$m = 2^{l}$$
 such that  $n < m$ .
$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + (m - n)\alpha}{m}$$

$$= \frac{x_1 + \dots + x_n + \overbrace{\alpha + \dots + \alpha}^{m - n}}{m} \ge \sqrt[m]{x_1 \dots x_n \alpha^{m - n}}$$



Let 
$$m = 2^{l}$$
 such that  $n < m$ .

$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + (m - n)\alpha}{m}$$

$$= \frac{x_1 + \dots + x_n + \alpha + \dots + \alpha}{m} \ge \sqrt[m]{x_1 \dots x_n \alpha^{m-n}}$$

$$\alpha^m \ge x_1 \dots x_n \alpha^{m-n}$$
,



Let 
$$m = 2^{l}$$
 such that  $n < m$ .

$$\alpha = \frac{x_{1} + \dots + x_{n}}{n} = \frac{\frac{m}{n}(x_{1} + \dots + x_{n})}{m}$$

$$= \frac{x_{1} + \dots + x_{n} + \frac{m - n}{n}(x_{1} + \dots + x_{n})}{m}$$

$$= \frac{x_{1} + \dots + x_{n} + (m - n)\alpha}{m}$$

$$= \frac{x_{1} + \dots + x_{n} + \alpha + \dots + \alpha}{m} \ge \sqrt[m]{x_{1} \dots x_{n}\alpha^{m-n}}$$

$$\alpha^{m} \ge x_{1} \dots x_{n}\alpha^{m-n} , \quad \alpha^{n} \ge x_{1} \dots x_{n} ,$$



Let 
$$m = 2^{l}$$
 such that  $n < m$ .

$$\alpha = \frac{x_{1} + \dots + x_{n}}{n} = \frac{\frac{m}{n}(x_{1} + \dots + x_{n})}{m}$$

$$= \frac{x_{1} + \dots + x_{n} + \frac{m - n}{n}(x_{1} + \dots + x_{n})}{\frac{m}{n}}$$

$$= \frac{x_{1} + \dots + x_{n} + (m - n)\alpha}{m}$$

$$= \frac{x_{1} + \dots + x_{n} + \alpha + \dots + \alpha}{m} \ge \sqrt[m]{x_{1} \dots x_{n}\alpha^{m-n}}$$

$$\alpha^{m} \ge x_{1} \dots x_{n}\alpha^{m-n} , \quad \alpha^{n} \ge x_{1} \dots x_{n} , \quad \alpha \ge \sqrt[n]{x_{1} \dots x_{n}}$$

#### ► Home ► Start ► End

Let 
$$m=2^l$$
 such that  $n < m$ .

$$\alpha = \frac{x_1 + \dots + x_n}{n} = \frac{\frac{m}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + \frac{m - n}{n}(x_1 + \dots + x_n)}{m}$$

$$= \frac{x_1 + \dots + x_n + (m - n)\alpha}{m}$$

$$= \frac{x_1 + \dots + x_n + \overbrace{\alpha + \dots + \alpha}^{m-n}}{m} \ge \sqrt[m]{x_1 \dots x_n}\alpha^{m-n}$$

$$\alpha^m \ge x_1 \dots x_n \alpha^{m-n} , \quad \alpha^n \ge x_1 \dots x_n , \quad \alpha \ge \sqrt[n]{x_1 \dots x_n}$$

$$\therefore \frac{x_1 + \dots + x_n}{n} \ge \sqrt[n]{x_1 \dots x_n} \quad (n \text{ is true.})$$







$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n}$$



$$\frac{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}{n} \geq$$



$$\frac{\frac{1}{x_1} + \dots + \frac{1}{x_n}}{n} \geq \sqrt[n]{\frac{1}{x_1} \cdots \frac{1}{x_n}}$$

#### → Home → Start → End

$$\frac{\frac{1}{x_1} + \dots + \frac{1}{x_n}}{n} \geq \sqrt[n]{\frac{1}{x_1} \dots \frac{1}{x_n}}$$

$$\frac{1}{x_1} + \dots + \frac{1}{x_n}$$

#### ► Home ► Start ► End

$$\frac{\frac{1}{x_1} + \dots + \frac{1}{x_n}}{n} \geq \sqrt[n]{\frac{1}{x_1} \dots \frac{1}{x_n}}$$

$$\frac{\frac{1}{x_1} + \dots + \frac{1}{x_n}}{n} \geq$$

#### → Home → Start → End

$$\frac{\frac{1}{x_1} + \dots + \frac{1}{x_n}}{n} \geq \sqrt[n]{\frac{1}{x_1} \dots \frac{1}{x_n}}$$

$$\frac{\frac{1}{x_1} + \dots + \frac{1}{x_n}}{n} \geq \frac{1}{\sqrt[n]{x_1 \cdots x_n}}$$

#### → Home → Start → End

$$\frac{\frac{1}{x_1} + \dots + \frac{1}{x_n}}{n} \geq \sqrt[n]{\frac{1}{x_1} \dots \frac{1}{x_n}}$$

$$\frac{\frac{1}{x_1} + \dots + \frac{1}{x_n}}{n} \geq \frac{1}{\sqrt[n]{x_1 \cdots x_n}}$$

$$\therefore \sqrt[n]{x_1 \times \cdots \times x_n} \geq$$

#### → Home → Start → End

$$\frac{\frac{1}{x_1} + \dots + \frac{1}{x_n}}{n} \geq \sqrt[n]{\frac{1}{x_1} \dots \frac{1}{x_n}}$$

$$\frac{\frac{1}{x_1} + \dots + \frac{1}{x_n}}{n} \geq \frac{1}{\sqrt[n]{x_1 \dots x_n}}$$

$$\therefore \sqrt[n]{x_1 \times \cdots \times x_n} \ge \frac{n}{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}$$

#### → Home → Start → End

$$\frac{\frac{1}{x_1} + \dots + \frac{1}{x_n}}{n} \geq \sqrt[n]{\frac{1}{x_1} \dots \frac{1}{x_n}}$$

$$\frac{\frac{1}{x_1} + \dots + \frac{1}{x_n}}{n} \geq \frac{1}{\sqrt[n]{x_1 \dots x_n}}$$

$$\therefore \sqrt[n]{x_1 \times \cdots \times x_n} \ge \frac{n}{\frac{1}{x_1} + \cdots + \frac{1}{x_n}}$$

#### Github:

https://min7014.github.io/math20191104001.html

Click or paste URL into the URL search bar, and you can see a picture moving.