치환형 분수방정식 예제 (Example of Substitutional Fractional Equations)



$$\frac{x^2 - 2x - 1}{3} = \frac{1}{x - 2} - \frac{1}{x} \cdots (1)$$

$$\frac{x^2 - 2x - 1}{3} = \frac{1}{x - 2} - \frac{1}{x} \cdots (1)$$

$$\frac{x^2 - 2x - 1}{3} = \frac{2}{x^2 - 2x}$$

$$\frac{x^2 - 2x - 1}{3} = \frac{1}{x - 2} - \frac{1}{x} \cdots (1)$$

$$\frac{x^2 - 2x - 1}{3} = \frac{2}{x^2 - 2x}$$
Let $t = x^2 - 2x$

$$\frac{x^2 - 2x - 1}{3} = \frac{1}{x - 2} - \frac{1}{x} \cdots (1)$$

$$\frac{x^2 - 2x - 1}{3} = \frac{2}{x^2 - 2x}$$

$$\text{Let } t = x^2 - 2x$$

$$\frac{t - 1}{3} = \frac{2}{t}$$

$$\frac{x^2 - 2x - 1}{3} = \frac{1}{x - 2} - \frac{1}{x} \cdots (1)$$

$$\frac{x^2 - 2x - 1}{3} = \frac{2}{x^2 - 2x}$$

$$\text{Let } t = x^2 - 2x$$

$$\frac{t - 1}{3} = \frac{2}{t}$$

$$t^2 - t - 6 = 0$$

$$\frac{x^2 - 2x - 1}{3} = \frac{1}{x - 2} - \frac{1}{x} \cdots (1)$$

$$\frac{x^2 - 2x - 1}{3} = \frac{2}{x^2 - 2x}$$

$$\text{Let } t = x^2 - 2x$$

$$\frac{t - 1}{3} = \frac{2}{t}$$

$$t^2 - t - 6 = 0 \qquad t = 3, -2$$

$$\frac{x^2 - 2x - 1}{3} = \frac{1}{x - 2} - \frac{1}{x} \cdots (1)$$

$$\frac{x^2 - 2x - 1}{3} = \frac{2}{x^2 - 2x}$$

$$\text{Let } t = x^2 - 2x$$

$$\frac{t - 1}{3} = \frac{2}{t}$$

$$t^2 - t - 6 = 0 \qquad t = 3, -2$$

$$x^2 - 2x = 3, \ x^2 - 2x = -2$$

$$\frac{x^2 - 2x - 1}{3} = \frac{1}{x - 2} - \frac{1}{x} \cdots (1)$$

$$\frac{x^2 - 2x - 1}{3} = \frac{2}{x^2 - 2x}$$

$$\text{Let } t = x^2 - 2x$$

$$\frac{t - 1}{3} = \frac{2}{t}$$

$$t^2 - t - 6 = 0 \qquad t = 3, -2$$

$$x^2 - 2x = 3, \quad x^2 - 2x = -2 \qquad x = 3, -1, 1 + i, 1 - i$$

$$\frac{x^2 - 2x - 1}{3} = \frac{1}{x - 2} - \frac{1}{x} \cdots (1)$$

$$\frac{x^2 - 2x - 1}{3} = \frac{2}{x^2 - 2x}$$
Let $t = x^2 - 2x$

$$\frac{t - 1}{3} = \frac{2}{t}$$

$$t^2 - t - 6 = 0 \qquad t = 3, -2$$

$$x^2 - 2x = 3, \quad x^2 - 2x = -2 \qquad x = 3, -1, 1 + i, 1 - i$$

$$(1) 식의 분모를 0 을 만들지 않으므로$$

$$\frac{x^2 - 2x - 1}{3} = \frac{1}{x - 2} - \frac{1}{x} \cdots (1)$$

$$\frac{x^2 - 2x - 1}{3} = \frac{2}{x^2 - 2x}$$
Let $t = x^2 - 2x$

$$\frac{t - 1}{3} = \frac{2}{t}$$

$$t^2 - t - 6 = 0 \qquad t = 3, -2$$

$$x^2 - 2x = 3, \quad x^2 - 2x = -2 \qquad x = 3, -1, 1 + i, 1 - i$$

$$(1) 식의 분모를 0을 만들지 않으므로$$

 $\therefore x = 3, -1, 1 + i, 1 - i$

Github:

https://min7014.github.io/math20210213001.html

Click or paste URL into the URL search bar, and you can see a picture moving.