

성분을 이용한 내적의 공식

(The Formula of Dot Product Using Components)

The Formula of Dot Product Using Components


▶ Start

▶ End

The Formula of Dot Product Using Components

▶ Start


▶ End

$$\vec{a} = (a_1, a_2, a_3)$$


The Formula of Dot Product Using Components

▶ Start


▶ End



The Formula of Dot Product Using Components

▶ Start

▶ End




$$\vec{a} - \vec{b} = (a_1 - b_1, a_2 - b_2, a_3 - b_3)$$

The Formula of Dot Product Using Components

▶ Start


▶ End



The Formula of Dot Product Using Components

▶ Start


▶ End



The Formula of Dot Product Using Components

▶ Start


▶ End



The Formula of Dot Product Using Components

▶ Start


▶ End



The Formula of Dot Product Using Components

▶ Start

▶ End



$|\vec{b}| = \sqrt{b_1^2 + b_2^2 + b_3^2}$

$\vec{b} = (b_1, b_2, b_3)$

$\vec{a} = (a_1, a_2, a_3)$

θ


$\vec{a} - \vec{b} = (a_1 - b_1, a_2 - b_2, a_3 - b_3)$

$|\vec{a} - \vec{b}| = \sqrt{(a_1 - b_1)^2 + (a_2 - b_2)^2 + (a_3 - b_3)^2}$

$|\vec{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$

$|\vec{a} - \vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 - 2|\vec{a}| |\vec{b}| \cos \theta$ (제2 코사인 법칙)

The Formula of Dot Product Using Components



The Formula of Dot Product Using Components

- ▶ Start

$$|\vec{b}| = \sqrt{b_1^2 + b_2^2 + b_3^2}$$

$$\vec{a} - \vec{b} = (a_1 - b_1, a_2 - b_2, a_3 - b_3)$$

$$|\vec{a} - \vec{b}| = \sqrt{(a_1 - b_1)^2 + (a_2 - b_2)^2 + (a_3 - b_3)^2}$$

$$\vec{b} = (b_1, b_2, b_3)$$

$$\vec{a} = (a_1, a_2, a_3)$$

$$|\vec{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$$


$$|\vec{a} - \vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 - 2(\vec{a} \cdot \vec{b})$$

$$\vec{a} \cdot \vec{b} = \frac{1}{2} (|\vec{a}|^2 + |\vec{b}|^2 - |\vec{a} - \vec{b}|^2)$$

The Formula of Dot Product Using Components

▶ Start

▶ End



$|\vec{b}| = \sqrt{b_1^2 + b_2^2 + b_3^2}$

$\vec{b} = (b_1, b_2, b_3)$

$\vec{a} = (a_1, a_2, a_3)$

θ

$\vec{a} - \vec{b} = (a_1 - b_1, a_2 - b_2, a_3 - b_3)$

$|\vec{a} - \vec{b}| = \sqrt{(a_1 - b_1)^2 + (a_2 - b_2)^2 + (a_3 - b_3)^2}$


$|\vec{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$

$$|\vec{a} - \vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 - 2(\vec{a} \cdot \vec{b})$$
$$\vec{a} \cdot \vec{b} = \frac{1}{2} (|\vec{a}|^2 + |\vec{b}|^2 - |\vec{a} - \vec{b}|^2)$$
$$= \frac{1}{2} \{ a_1^2 + a_2^2 + a_3^2 + b_1^2 + b_2^2 + b_3^2 - (a_1 - b_1)^2 - (a_2 - b_2)^2 - (a_3 - b_3)^2 \}$$

The Formula of Dot Product Using Components

▶ Start

▶ End



$$|\vec{a} - \vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 - 2(\vec{a} \cdot \vec{b})$$

$$\vec{a} \cdot \vec{b} = \frac{1}{2} (|\vec{a}|^2 + |\vec{b}|^2 - |\vec{a} - \vec{b}|^2)$$

$$= \frac{1}{2} \{ a_1^2 + a_2^2 + a_3^2 + b_1^2 + b_2^2 + b_3^2 - (a_1 - b_1)^2 - (a_2 - b_2)^2 - (a_3 - b_3)^2 \}$$

$$\therefore \vec{a} \cdot \vec{b} = a_1 b_1 + a_2 b_2 + a_3 b_3$$

Github:

<https://min7014.github.io/math20221102002.html>

Click or paste URL into the URL search bar,
and you can see a picture moving.