

**ANNEX I**

**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE MEDICINAL PRODUCT

Aectura Breezhaler 125 micrograms/62.5 micrograms inhalation powder, hard capsules  
Aectura Breezhaler 125 micrograms/127.5 micrograms inhalation powder, hard capsules  
Aectura Breezhaler 125 micrograms/260 micrograms inhalation powder, hard capsules

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Aectura Breezhaler 125 micrograms/62.5 micrograms inhalation powder, hard capsules

Each capsule contains 150 mcg of indacaterol (as acetate) and 80 mcg of mometasone furoate.

Each delivered dose (the dose that leaves the mouthpiece of the inhaler) contains 125 mcg of indacaterol (as acetate) and 62.5 mcg of mometasone furoate.

Aectura Breezhaler 125 micrograms/127.5 micrograms inhalation powder, hard capsules

Each capsule contains 150 mcg of indacaterol (as acetate) and 160 mcg of mometasone furoate.

Each delivered dose (the dose that leaves the mouthpiece of the inhaler) contains 125 mcg of indacaterol (as acetate) and 127.5 mcg of mometasone furoate.

Aectura Breezhaler 125 micrograms/260 micrograms inhalation powder, hard capsules

Each capsule contains 150 mcg of indacaterol (as acetate) and 320 mcg of mometasone furoate.

Each delivered dose (the dose that leaves the mouthpiece of the inhaler) contains 125 mcg of indacaterol (as acetate) and 260 mcg of mometasone furoate.

Excipient(s) with known effect

Each capsule contains approximately 25 mg of lactose monohydrate.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Inhalation powder, hard capsule (inhalation powder).

Aectura Breezhaler 125 micrograms/62.5 micrograms inhalation powder, hard capsules

Transparent (uncoloured) capsules containing a white powder, with the product code “IM150-80” printed in blue above one blue bar on the body and with the product logo printed in blue and surrounded by two blue bars on the cap.

Aectura Breezhaler 125 micrograms/127.5 micrograms inhalation powder, hard capsules

Transparent (uncoloured) capsules containing a white powder, with the product code “IM150-160” printed in grey on the body and with the product logo printed in grey on the cap.

Aectura Breezhaler 125 micrograms/260 micrograms inhalation powder, hard capsules

Transparent (uncoloured) capsules containing a white powder, with the product code “IM150-320” printed in black above two black bars on the body and with the product logo printed in black and surrounded by two black bars on the cap.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Ateectura Breezhaler is indicated as a maintenance treatment of asthma in adults and adolescents 12 years of age and older not adequately controlled with inhaled corticosteroids and inhaled short-acting beta<sub>2</sub>-agonists.

### **4.2 Posology and method of administration**

#### Posology

##### Adults and adolescents aged 12 years and over

The recommended dose is one capsule to be inhaled once daily.

Patients should be given the strength containing the appropriate mometasone furoate dosage for the severity of their disease and should be regularly reassessed by a healthcare professional.

The maximum recommended dose is 125 mcg/260 mcg once daily.

Treatment should be administered at the same time of the day each day. It can be administered irrespective of the time of the day. If a dose is missed, it should be taken as soon as possible. Patients should be instructed not to take more than one dose in a day.

##### Special populations

###### Elderly population

No dose adjustment is required in elderly patients (65 years of age or older) (see section 5.2).

###### Renal impairment

No dose adjustment is required in patients with renal impairment (see section 5.2).

###### Hepatic impairment

No dose adjustment is required in patients with mild or moderate hepatic impairment. No data are available for the use of the medicinal product in patients with severe hepatic impairment, therefore it should be used in these patients only if the expected benefit outweighs the potential risk (see section 5.2).

###### Paediatric population

The posology in patients 12 years of age and older is the same posology as in adults. The safety and efficacy in paediatric patients below 12 years of age have not been established. No data are available.

### Method of administration

For inhalation use only. The capsules must not be swallowed.

The capsules must be administered only using the inhaler provided (see section 6.6) with each new prescription.

Patients should be instructed on how to administer the medicinal product correctly. Patients who do not experience improvement in breathing should be asked if they are swallowing the medicinal product rather than inhaling it.

The capsules must only be removed from the blister immediately before use.

After inhalation, patients should rinse their mouth with water without swallowing (see sections 4.4 and 6.6).

For instructions on use of the medicinal product before administration, see section 6.6.

### Information for patients using a sensor for Ateectura Breezhaler

The pack may contain an electronic sensor to be attached to the base of the inhaler.

The sensor for Ateectura Breezhaler is for adult use only.

The sensor for Ateectura Breezhaler should not be used for adolescent patients as the App does not have patient consent verification or caregiver account functionality.

The sensor and App are not required for administration of the medicinal product to the patient. The sensor and App do not control or interfere with delivery of the medicinal product using the inhaler.

The prescribing physician may discuss with the patient whether the use of the sensor and App is appropriate.

For detailed instructions on how to use the sensor and the App, see the Instructions for Use provided in the sensor pack and the App.

## **4.3 Contraindications**

Hypersensitivity to the active substances or to any of the excipients listed in section 6.1.

## **4.4 Special warnings and precautions for use**

### Deterioration of disease

This medicinal product should not be used to treat acute asthma symptoms, including acute episodes of bronchospasm, for which a short-acting bronchodilator is required. Increasing use of short-acting bronchodilators to relieve symptoms indicates deterioration of control and patients should be reviewed by a physician.

Patients should not stop treatment without physician supervision since symptoms may recur after discontinuation.

It is recommended that treatment with this medicinal product should not be stopped abruptly. If patients find the treatment ineffective, they should continue treatment but must seek medical attention. Increasing use of reliever bronchodilators indicates a worsening of the underlying condition and warrants a reassessment of the therapy. Sudden and progressive deterioration in the symptoms of asthma is potentially life-threatening and the patient should undergo urgent medical assessment.

## Hypersensitivity

Immediate hypersensitivity reactions have been observed after administration of this medicinal product. If signs suggesting allergic reactions occur, in particular angioedema (including difficulties in breathing or swallowing, swelling of the tongue, lips and face), urticaria or skin rash, treatment should be discontinued immediately and alternative therapy instituted.

## Paradoxical bronchospasm

As with other inhalation therapy, administration of this medicinal product may result in paradoxical bronchospasm, which can be life-threatening. If this occurs, treatment should be discontinued immediately and alternative therapy instituted.

## Cardiovascular effects of beta agonists

Like other medicinal products containing beta<sub>2</sub>-adrenergic agonists, this medicinal product may produce a clinically significant cardiovascular effect in some patients, as measured by increases in pulse rate, blood pressure and/or symptoms. If such effects occur, treatment may need to be discontinued.

This medicinal product should be used with caution in patients with cardiovascular disorders (coronary artery disease, acute myocardial infarction, cardiac arrhythmias, hypertension), convulsive disorders or thyrotoxicosis, and in patients who are unusually responsive to beta<sub>2</sub>-adrenergic agonists.

Patients with unstable ischaemic heart disease, a history of myocardial infarction in last 12 months, New York Heart Association (NYHA) class III/IV left ventricular failure, arrhythmia, uncontrolled hypertension, cerebrovascular disease or history of long QT syndrome and patients being treated with medicinal products known to prolong QTc were excluded from studies in the indacaterol/mometasone furoate clinical development programme. Thus safety outcomes in these populations are considered unknown.

While beta<sub>2</sub>-adrenergic agonists have been reported to produce electrocardiographic (ECG) changes, such as flattening of the T wave, prolongation of QT interval and ST segment depression, the clinical significance of these observations is unknown.

Long-acting beta<sub>2</sub>-adrenergic agonists (LABA) or LABA-containing combination products such as Atecura Breezhaler should therefore be used with caution in patients with known or suspected prolongation of the QT interval or who are being treated with medicinal products affecting the QT interval.

## Hypokalaemia with beta agonists

Beta<sub>2</sub>-adrenergic agonists may produce significant hypokalaemia in some patients, which has the potential to produce adverse cardiovascular effects. The decrease in serum potassium is usually transient, not requiring supplementation. In patients with severe asthma hypokalaemia may be potentiated by hypoxia and concomitant treatment, which may increase the susceptibility to cardiac arrhythmias (see section 4.5).

Clinically relevant hypokalaemia has not been observed in clinical studies of indacaterol/mometasone furoate at the recommended therapeutic dose.

### Hyperglycaemia

Inhalation of high doses of beta<sub>2</sub>-adrenergic agonists and corticosteroids may produce increases in plasma glucose. Upon initiation of treatment, plasma glucose should be monitored more closely in diabetic patients.

This medicinal product has not been investigated in patients with Type I diabetes mellitus or uncontrolled Type II diabetes mellitus.

### Prevention of oropharyngeal infections

In order to reduce the risk of oropharyngeal candida infection, patients should be advised to rinse their mouth or gargle with water without swallowing it or brush their teeth after inhaling the prescribed dose.

### Systemic effects of corticosteroids

Systemic effects of inhaled corticosteroids may occur, particularly at high doses prescribed for prolonged periods. These effects are much less likely to occur than with oral corticosteroids and may vary in individual patients and between different corticosteroid preparations.

Possible systemic effects may include Cushing's syndrome, Cushingoid features, adrenal suppression, growth retardation in children and adolescents, decrease in bone mineral density, cataracts, glaucoma, and, more rarely, a range of psychological or behavioural effects including psychomotor hyperactivity, sleep disorders, anxiety, depression or aggression (particularly in children). It is therefore important that the dose of inhaled corticosteroid is titrated to the lowest dose at which effective control of asthma is maintained.

Visual disturbance may be reported with systemic and topical (including intranasal, inhaled and intraocular) corticosteroid use. Patients presenting with symptoms such as blurred vision or other visual disturbances should be considered for referral to an ophthalmologist for evaluation of possible causes of visual disturbances, which may include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids.

This medicinal product should be administered with caution in patients with pulmonary tuberculosis or in patients with chronic or untreated infections.

### Excipients

This medicinal product contains lactose. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicinal product.

## **4.5 Interaction with other medicinal products and other forms of interaction**

No specific interaction studies were conducted with indacaterol/mometasone furoate. Information on the potential for interactions is based on the potential for each of the monotherapy components.

### Medicinal products known to prolong the QTc interval

Like other medicinal products containing a beta<sub>2</sub>-adrenergic agonist, this medicinal product should be administered with caution to patients being treated with monoamine oxidase inhibitors, tricyclic antidepressants or medicinal products known to prolong the QT interval, as any effect of these on the QT interval may be potentiated. Medicinal products known to prolong the QT interval may increase the risk of ventricular arrhythmia (see sections 4.4 and 5.1).

### Hypokalaemic treatment

Concomitant hypokalaemic treatment with methylxanthine derivatives, steroids or non-potassium-sparing diuretics may potentiate the possible hypokalaemic effect of beta<sub>2</sub>-adrenergic agonists (see section 4.4).

### Beta-adrenergic blockers

Beta-adrenergic blockers may weaken or antagonise the effect of beta<sub>2</sub>-adrenergic agonists. Therefore, this medicinal product should not be given together with beta-adrenergic blockers unless there are compelling reasons for their use. Where required, cardioselective beta-adrenergic blockers should be preferred, although they should be administered with caution.

### Interaction with CYP3A4 and P-glycoprotein inhibitors

Inhibition of CYP3A4 and P-glycoprotein (P-gp) has no impact on the safety of therapeutic doses of Atecura Breezhaler.

Inhibition of the key contributors of indacaterol clearance (CYP3A4 and P-gp) or mometasone furoate clearance (CYP3A4) raises the systemic exposure of indacaterol or mometasone furoate up to two-fold.

Due to the very low plasma concentration achieved after inhaled dosing, clinically significant interactions with mometasone furoate are unlikely. However, there may be a potential for increased systemic exposure to mometasone furoate when strong CYP3A4 inhibitors (e.g. ketoconazole, itraconazole, nelfinavir, ritonavir, cobicistat) are co-administered.

### Other long-acting beta<sub>2</sub>-adrenergic agonists

The co-administration of this medicinal product with other medicinal products containing long-acting beta<sub>2</sub>-adrenergic agonists has not been studied and is not recommended as it may potentiate adverse reactions (see sections 4.8 and 4.9).

## **4.6 Fertility, pregnancy and lactation**

### Pregnancy

There are insufficient data from the use of Atecura Breezhaler or its individual components (indacaterol and mometasone furoate) in pregnant women to determine whether there is a risk.

Indacaterol was not teratogenic in rats and rabbits following subcutaneous administration (see section 5.3). In animal reproduction studies with pregnant mice, rats and rabbits, mometasone furoate caused increased foetal malformations and decreased foetal survival and growth.

Like other medicinal products containing beta<sub>2</sub>-adrenergic agonists, indacaterol may inhibit labour due to a relaxant effect on uterine smooth muscle.

This medicinal product should only be used during pregnancy if the expected benefit to the patient justifies the potential risk to the foetus.

### Breast-feeding

There is no information available on the presence of indacaterol or mometasone furoate in human milk, on the effects on a breast-fed infant, or on the effects on milk production. Other inhaled corticosteroids similar to mometasone furoate are transferred into human milk. Indacaterol (including its metabolites) and mometasone furoate have been detected in the milk of lactating rats.

A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from therapy, taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

### Fertility

Reproduction studies and other data in animals did not indicate a concern regarding fertility in either males or females.

## **4.7 Effects on ability to drive and use machines**

This medicinal product has no or negligible influence on the ability to drive and use machines.

## **4.8 Undesirable effects**

### Summary of the safety profile

The most common adverse reactions over 52 weeks were asthma (exacerbation) (26.9%), nasopharyngitis (12.9%), upper respiratory tract infection (5.9%) and headache (5.8%).

### Tabulated list of adverse reactions

Adverse drug reactions (ADRs) are listed by MedDRA system organ class (Table 1). The frequency of the ADRs is based on the PALLADIUM study. Within each system organ class, the adverse drug reactions are ranked by frequency, with the most frequent reactions first. Within each frequency grouping, adverse drug reactions are presented in order of decreasing seriousness. In addition, the corresponding frequency category for each adverse drug reaction is based on the following convention (CIOMS III): very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1\ 000$  to  $< 1/100$ ); rare ( $\geq 1/10\ 000$  to  $< 1/1\ 000$ ); very rare ( $< 1/10\ 000$ ).



**Table 1 Adverse reactions**

System organ class	Adverse reactions	Frequency category
Infections and infestations	Nasopharyngitis	Very common
	Upper respiratory tract infection	Common
	Candidiasis* <sup>1</sup>	Uncommon
Immune system disorders	Hypersensitivity* <sup>2</sup>	Common
	Angioedema* <sup>3</sup>	Uncommon
Metabolism and nutrition disorders	Hyperglycaemia* <sup>4</sup>	Uncommon
Nervous system disorders	Headache* <sup>5</sup>	Common
Eye disorders	Vision blurred	Uncommon
	Cataract* <sup>6</sup>	Uncommon
Cardiac disorders	Tachycardia* <sup>7</sup>	Uncommon
Respiratory, thoracic and mediastinal disorders	Asthma (exacerbation)	Very common
	Oropharyngeal pain* <sup>8</sup>	Common
	Dysphonia	Common
Skin and subcutaneous tissue disorders	Rash* <sup>9</sup>	Uncommon
	Pruritus* <sup>10</sup>	Uncommon
Musculoskeletal and connective tissue disorders	Musculoskeletal pain* <sup>11</sup>	Common
	Muscle spasms	Uncommon
<p>* Indicates grouping of preferred terms (PTs):</p> <p>1 Oral candidiasis, oropharyngeal candidiasis.</p> <p>2 Drug eruption, drug hypersensitivity, hypersensitivity, rash, rash erythematous, rash pruritic, urticaria.</p> <p>3 Allergic oedema, angioedema, periorbital swelling, swelling of eyelid.</p> <p>4 Blood glucose increased, hyperglycaemia.</p> <p>5 Headache, tension headache.</p> <p>6 Cataract, cataract cortical.</p> <p>7 Heart rate increased, tachycardia, sinus tachycardia, supraventricular tachycardia.</p> <p>8 Oral pain, oropharyngeal discomfort, oropharyngeal pain, throat irritation, odynophagia.</p> <p>9 Drug eruption, rash, rash erythematous, rash pruritic.</p> <p>10 Anal pruritus, eye pruritus, nasal pruritus, pruritus, pruritus genital.</p> <p>11 Back pain, musculoskeletal pain, myalgia, neck pain, musculoskeletal chest pain.</p>		

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

## **4.9 Overdose**

General supportive measures and symptomatic treatment should be initiated in cases of suspected overdose.

An overdose will likely produce signs, symptoms or adverse effects associated with the pharmacological actions of the individual components (e.g. tachycardia, tremor, palpitations, headache, nausea, vomiting, drowsiness, ventricular arrhythmias, metabolic acidosis, hypokalaemia, hyperglycaemia, suppression of hypothalamic pituitary adrenal axis function).

Use of cardioselective beta blockers may be considered for treating beta<sub>2</sub>-adrenergic effects, but only under the supervision of a physician and with extreme caution, since the use of beta<sub>2</sub>-adrenergic blockers may provoke bronchospasm. In serious cases, patients should be hospitalised.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotheapeutic group: Drugs for obstructive airway diseases, adrenergics in combination with corticosteroids or other drugs, excl. anticholinergics, ATC code: R03AK14

#### Mechanism of action

This medicinal product is a combination of indacaterol, a long-acting beta<sub>2</sub>-adrenergic agonist (LABA), and mometasone furoate, an inhaled synthetic corticosteroid (ICS).

#### Indacaterol

The pharmacological effects of beta<sub>2</sub>-adrenoceptor agonists, including indacaterol, are at least in part attributable to increased cyclic-3', 5'-adenosine monophosphate (cyclic AMP) levels, which cause relaxation of bronchial smooth muscle.

When inhaled, indacaterol acts locally in the lung as a bronchodilator. Indacaterol is a partial agonist at the human beta<sub>2</sub>-adrenergic receptor with nanomolar potency. In isolated human bronchus, indacaterol has a rapid onset of action and a long duration of action.

Although beta<sub>2</sub>-adrenergic receptors are the predominant adrenergic receptors in bronchial smooth muscle and beta<sub>1</sub>-receptors are the predominant receptors in the human heart, there are also beta<sub>2</sub>-adrenergic receptors in the human heart comprising 10% to 50% of the total adrenergic receptors.

#### Mometasone furoate

Mometasone furoate is a synthetic corticosteroid with high affinity for glucocorticoid receptors and local anti-inflammatory properties. *In vitro*, mometasone furoate inhibits the release of leukotrienes from leukocytes of allergic patients. In cell culture, mometasone furoate demonstrated high potency in inhibition of synthesis and release of IL-1, IL-5, IL-6 and TNF-alpha. It is also a potent inhibitor of leukotriene production and of the production of the Th2 cytokines IL-4 and IL-5 from human CD4+ T-cells.

#### Pharmacodynamic effects

The pharmacodynamic response profile of this medicinal product is characterised by rapid onset of action within 5 minutes after dosing and sustained effect over the 24-hour dosing interval, as evidenced by improvements in trough forced expiratory volume in the first second (FEV<sub>1</sub>) improvements versus comparators 24 hours after dosing.

No tachyphylaxis to the lung function benefits of this medicinal product was observed over time.

#### QTc interval

The effect of this medicinal product on the QTc interval has not been evaluated in a thorough QT (TQT) study. For mometasone furoate, no QTc-prolonging properties are known.

### Clinical efficacy and safety

Two phase III randomised, double-blind studies (PALLADIUM and QUARTZ) of different durations evaluated the safety and efficacy of Atecura Breezhaler in adult and adolescent patients with persistent asthma.

The PALLADIUM study was a 52-week pivotal study evaluating Atecura Breezhaler 125 mcg/127.5 mcg once daily (N=439) and 125 mcg/260 mcg once daily (N=445) compared to mometasone furoate 400 mcg once daily (N=444) and 800 mcg per day (given as 400 mcg twice daily) (N=442), respectively. A third active control arm included subjects treated with salmeterol/fluticasone propionate 50 mcg/500 mcg twice daily (N=446). All subjects were required to have symptomatic asthma (ACQ-7 score  $\geq 1.5$ ) and were on asthma maintenance therapy using an inhaled synthetic corticosteroid (ICS) with or without LABA for at least 3 months prior to study entry. At screening, 31% of patients had history of exacerbation in the previous year. At study entry, the most common asthma medications reported were medium dose of ICS (20%), high dose of ICS (7%) or low dose of ICS in combination with a LABA (69%).

The primary objective of the study was to demonstrate superiority of either Atecura Breezhaler 125 mcg/127.5 mcg once daily over mometasone furoate 400 mcg once daily or Atecura Breezhaler 125 mcg/260 mcg once daily over mometasone furoate 400 mcg twice daily in terms of trough FEV<sub>1</sub> at week 26.

At week 26, Atecura Breezhaler 125 mcg/127.5 mcg and 125 mcg/260 mcg once daily both demonstrated statistically significant improvements in trough FEV<sub>1</sub> and Asthma Control Questionnaire (ACQ-7) score compared to mometasone furoate 400 mcg once or twice daily, respectively (see Table 2). Findings at week 52 were consistent with week 26.

Atecura Breezhaler 125 mcg/127.5 mcg and 125 mcg/260 mcg once daily both demonstrated a clinically meaningful reduction in the annual rate of moderate or severe exacerbations (secondary endpoint), compared to mometasone furoate 400 mcg once and twice daily (see Table 2).

Results for the most clinically relevant endpoints are described in Table 2.

Table 2 Results of primary and secondary endpoints in PALLADIUM study at weeks 26 and 52

Endpoint	Time point/ Duration	Aectura Breezhaler <sup>1</sup> vs MF <sup>2</sup>		Aectura Breezhaler <sup>1</sup> vs SAL/FP <sup>3</sup>
		Medium dose vs medium dose	High dose vs high dose	High dose vs high dose
<b>Lung function</b>				
<i>Trough FEV<sub>1</sub><sup>4</sup></i>				
Treatment difference P value (95% CI)	Week 26 (primary endpoint)	211 ml <0.001 (167, 255)	132 ml <0.001 (88, 176)	36 ml 0.101 (-7, 80)
	Week 52	209 ml <0.001 (163, 255)	136 ml <0.001 (90, 183)	48 ml 0.040 (2, 94)
<i>Mean morning peak expiratory flow (PEF)*</i>				
Treatment difference (95% CI)	Week 52	30.2 l/min (24.2, 36.3)	28.7 l/min (22.7, 34.8)	13.8 l/min (7.7, 19.8)
<i>Mean evening peak expiratory flow (PEF)*</i>				
Treatment difference (95% CI)	Week 52	29.1 l/min (23.3, 34.8)	23.7 l/min (18.0, 29.5)	9.1 l/min (3.3, 14.9)
<b>Symptoms</b>				
<i>ACQ-7</i>				
Treatment difference P value (95% CI)	Week 26 (key secondary endpoint)	-0.248 <0.001 (-0.334, -0.162)	-0.171 <0.001 (-0.257, -0.086)	-0.054 0.214 (-0.140, 0.031)
	Week 52	-0.266 (-0.354, -0.177)	-0.141 (-0.229, -0.053)	0.010 (-0.078, 0.098)
<i>ACQ responders (percentage of patients achieving minimal clinical important difference (MCID) from baseline with ACQ ≥0.5)</i>				
Percentage	Week 26	76% vs 67%	76% vs 72%	76% vs 76%
Odds ratio (95% CI)	Week 26	1.73 (1.26, 2.37)	1.31 (0.95, 1.81)	1.06 (0.76, 1.46)
Percentage	Week 52	82% vs 69%	78% vs 74%	78% vs 77%
Odds ratio (95% CI)	Week 52	2.24 (1.58, 3.17)	1.34 (0.96, 1.87)	1.05 (0.75, 1.49)
<i>Percentage of rescue medication free days*</i>				
Treatment difference (95% CI)	Week 52	8.6 (4.7, 12.6)	9.6 (5.7, 13.6)	4.3 (0.3, 8.3)
<i>Percentage of days with no symptoms*</i>				
Treatment difference (95% CI)	Week 52	9.1 (4.6, 13.6)	5.8 (1.3, 10.2)	3.4 (-1.1, 7.9)

<b>Annualised rate of asthma exacerbations**</b>				
<i>Moderate or severe exacerbations</i>				
AR	Week 52	0.27 vs 0.56	0.25 vs 0.39	0.25 vs 0.27
RR (95% CI)	Week 52	0.47 (0.35, 0.64)	0.65 (0.48, 0.89)	0.93 (0.67, 1.29)
<i>Severe exacerbations</i>				
AR	Week 52	0.13 vs 0.29	0.13 vs 0.18	0.13 vs 0.14
RR (95% CI)	Week 52	0.46 (0.31, 0.67)	0.71 (0.47, 1.08)	0.89 (0.58, 1.37)
<p>* Mean value for the treatment duration</p> <p>** RR &lt;1.00 favours indacaterol/mometasone furoate.</p> <p>1 Atecura Breezhaler medium dose: 125 mcg/127.5 mcg od; high dose: 125 mcg/260 mcg od.</p> <p>2 MF: mometasone furoate medium dose: 400 mcg od; high dose: 400 mcg bid (content doses). Mometasone furoate 127.5 mcg od and 260 mcg od in Atecura Breezhaler are comparable to mometasone furoate 400 mcg od and 800 mcg per day (given as 400 mcg bid).</p> <p>3 SAL/FP: salmeterol/fluticasone propionate high dose: 50 mcg/500 mcg bid (content dose).</p> <p>4 Trough FEV<sub>1</sub>: the mean of the two FEV<sub>1</sub> values measured at 23 hours 15 min and 23 hours 45 min after the evening dose.</p> <p>Primary endpoint (trough FEV<sub>1</sub> at week 26) and key secondary endpoint (ACQ-7 score at week 26) were part of confirmatory testing strategy and thus controlled for multiplicity. All other endpoints were not part of confirmatory testing strategy.</p> <p>RR = rate ratio, AR = annualised rate od = once daily, bid = twice daily</p>				

### Pre-specified pooled analysis

Atecura Breezhaler 125 mcg/260 mcg once daily was also studied as an active comparator in another phase III study (IRIDIUM) in which all subjects had a history of asthma exacerbation requiring systemic corticosteroids in the past year. A pre-specified pooled analysis across the IRIDIUM and PALLADIUM studies was conducted to compare Atecura Breezhaler 125 mcg/260 mcg once daily to salmeterol/fluticasone 50 mcg/500 mcg twice daily for the endpoints of trough FEV<sub>1</sub> and ACQ-7 at week 26 and annualised rate of exacerbations. The pooled analysis demonstrated that Atecura Breezhaler improved trough FEV<sub>1</sub> by 43 ml (95% CI: 17, 69) and ACQ-7 score by -0.091 (95% CI: -0.153, -0.030) at week 26 and reduced the annualised rate of moderate or severe asthma exacerbations by 22% (RR: 0.78; 95% CI: 0.66, 0.93) and of severe exacerbations by 26% (RR: 0.74; 95% CI: 0.61, 0.91) versus salmeterol/fluticasone.

The QUARTZ study was a 12-week study evaluating Atecura Breezhaler 125 mcg/62.5 mcg once daily (N=398) compared to mometasone furoate 200 mcg once daily (N=404). All subjects were required to be symptomatic and on asthma maintenance therapy using a low-dose ICS (with or without LABA) for at least 1 month prior to study entry. At study entry, the most common asthma medications reported were low-dose ICS (43%) and LABA/low-dose ICS (56%). The primary endpoint of the study was to demonstrate superiority of Atecura Breezhaler 125 mcg/62.5 mcg once daily over mometasone furoate 200 mcg once daily in terms of trough FEV<sub>1</sub> at week 12.

Atecura Breezhaler 125 mcg/62.5 mcg once daily demonstrated a statistically significant improvement in baseline trough FEV<sub>1</sub> at week 12 and Asthma Control Questionnaire (ACQ-7) score compared to mometasone furoate 200 mcg once daily.

Results for the most clinically relevant endpoints are described in Table 3.

**Table 3 Results of primary and secondary endpoints in QUARTZ study at week 12**

Endpoints	Aectura Breezhaler low dose* vs MF low dose**
<b>Lung function</b>	
<i>Trough FEV<sub>1</sub> (primary endpoint)***</i>	
Treatment difference	182 ml
P value	<0.001
(95% CI)	(148, 217)
<i>Mean morning peak expiratory flow (PEF)</i>	
Treatment difference	27.2 l/min
(95% CI)	(22.1, 32.4)
<i>Evening peak expiratory flow (PEF)</i>	
Treatment difference	26.1 l/min
(95% CI)	(21.0, 31.2)
<b>Symptoms</b>	
<i>ACQ-7 (key secondary endpoint)</i>	
Treatment difference	-0.218
P value	<0.001
(95% CI)	(-0.293, -0.143)
<i>Percentage of patients achieving MCID from baseline with ACQ <math>\geq 0.5</math></i>	
Percentage	75% vs 65%
Odds ratio	1.69
(95% CI)	(1.23, 2.33)
<i>Percentage of rescue medication free days</i>	
Treatment difference	8.1
(95% CI)	(4.3, 11.8)
<i>Percentage of days with no symptoms</i>	
Treatment difference	2.7
(95% CI)	(-1.0, 6.4)
* Aectura Breezhaler low dose: 125/62.5 mcg od. ** MF: mometasone furoate low dose: 200 mcg od (content dose). Mometasone furoate 62.5 mcg in Aectura Breezhaler od is comparable to mometasone furoate 200 mcg od (content dose). *** Trough FEV <sub>1</sub> : the mean of the two FEV <sub>1</sub> values measured at 23 hours 15 min and 23 hours 45 min after the evening dose. od = once daily, bid = twice daily	

### Paediatric population

In the PALLADIUM study, which included 106 adolescents (12-17 years old), the improvements in trough FEV<sub>1</sub> at week 26 were 0.173 litres (95% CI: -0.021, 0.368) for Aectura Breezhaler 125 mcg/260 mcg once daily vs mometasone furoate 800 mcg (i.e. high doses) and 0.397 litres (95% CI: 0.195, 0.599) for Aectura Breezhaler 125 mcg/127.5 mcg once daily vs mometasone furoate 400 mcg once daily (i.e. medium doses).

In the QUARTZ study, which included 63 adolescents (12-17 years old), the Least Square means treatment difference for trough FEV<sub>1</sub> at day 85 (week 12) was 0.251 litres (95% CI: 0.130, 0.371).

For the adolescent subgroups, improvements in lung function, symptoms and exacerbation reductions were consistent with the overall population.

The European Medicines Agency has deferred the obligation to submit the results of studies with indacaterol/mometasone furoate in one or more subsets of the paediatric population in asthma (see section 4.2 for information on paediatric use).

## 5.2 Pharmacokinetic properties

### Absorption

Following inhalation of Atecura Breezhaler, the median time to reach peak plasma concentrations of indacaterol and mometasone furoate was approximately 15 minutes and 1 hour, respectively.

Based on the *in vitro* performance data, the dose of each of the monotherapy components delivered to the lung is expected to be similar for the indacaterol/mometasone furoate combination and the monotherapy products. Steady-state plasma exposure to indacaterol and mometasone furoate after inhalation of the combination was similar to the systemic exposure after inhalation of indacaterol maleate or mometasone furoate as monotherapy products.

Following inhalation of the combination, the absolute bioavailability was estimated to be about 45% for indacaterol and less than 10% for mometasone furoate.

### Indacaterol

Indacaterol concentrations increased with repeated once-daily administration. Steady state was achieved within 12 to 14 days. The mean accumulation ratio of indacaterol, i.e. AUC over the 24-h dosing interval on day 14 compared to day 1, was in the range of 2.9 to 3.8 for once-daily inhaled doses between 60 and 480 mcg (delivered dose). Systemic exposure results from a composite of pulmonary and gastrointestinal absorption; about 75% of systemic exposure was from pulmonary absorption and about 25% from gastrointestinal absorption.

### Mometasone furoate

Mometasone furoate concentrations increased with repeated once-daily administration via the Breezhaler inhaler. Steady state was achieved after 12 days. The mean accumulation ratio of mometasone furoate, i.e. AUC over the 24-h dosing interval on day 14 compared to day 1, was in the range of 1.61 to 1.71 for once-daily inhaled doses between 62.5 and 260 mcg as part of the indacaterol/mometasone furoate combination.

Following oral administration of mometasone furoate, the absolute oral systemic bioavailability of mometasone furoate was estimated to be very low (<2%).

### Distribution

### Indacaterol

After intravenous infusion the volume of distribution ( $V_z$ ) of indacaterol was 2 361 to 2 557 litres, indicating an extensive distribution. The *in vitro* human serum and plasma protein binding were 94.1 to 95.3% and 95.1 to 96.2%, respectively.

### Mometasone furoate

After intravenous bolus administration, the  $V_d$  is 332 litres. The *in vitro* protein binding for mometasone furoate is high, 98% to 99% in concentration range of 5 to 500 ng/ml.

## Biotransformation

### Indacaterol

After oral administration of radiolabelled indacaterol in a human ADME (absorption, distribution, metabolism, excretion) study, unchanged indacaterol was the main component in serum, accounting for about one third of total drug-related AUC over 24 hours. A hydroxylated derivative was the most prominent metabolite in serum. Phenolic O-glucuronides of indacaterol and hydroxylated indacaterol were further prominent metabolites. A diastereomer of the hydroxylated derivative, an N-glucuronide of indacaterol, and C- and N-dealkylated products were further metabolites identified.

*In vitro* investigations indicated that UGT1A1 was the only UGT isoform that metabolised indacaterol to the phenolic O-glucuronide. The oxidative metabolites were found in incubations with recombinant CYP1A1, CYP2D6 and CYP3A4. CYP3A4 is concluded to be the predominant isoenzyme responsible for hydroxylation of indacaterol. *In vitro* investigations further indicated that indacaterol is a low-affinity substrate for the efflux pump P-gp.

*In vitro* the UGT1A1 isoform is a major contributor to the metabolic clearance of indacaterol. However, as shown in a clinical study in populations with different UGT1A1 genotypes, systemic exposure to indacaterol is not significantly affected by the UGT1A1 genotype.

### Mometasone furoate

The portion of an inhaled mometasone furoate dose that is swallowed and absorbed in the gastrointestinal tract undergoes extensive metabolism to multiple metabolites. There are no major metabolites detectable in plasma. In human liver microsomes mometasone furoate is metabolised by CYP3A4.

## Elimination

### Indacaterol

In clinical studies which included urine collection, the amount of indacaterol excreted unchanged via urine was generally lower than 2% of the dose. Renal clearance of indacaterol was, on average, between 0.46 and 1.20 litres/hour. Compared with the serum clearance of indacaterol of 18.8 to 23.3 litres/hour, it is evident that renal clearance plays a minor role (about 2 to 6% of systemic clearance) in the elimination of systemically available indacaterol.

In a human ADME study in which indacaterol was given orally, the faecal route of excretion was dominant over the urinary route. Indacaterol was excreted into human faeces primarily as unchanged parent substance (54% of the dose) and, to a lesser extent, hydroxylated indacaterol metabolites (23% of the dose). Mass balance was complete with  $\geq 90\%$  of the dose recovered in the excreta.

Indacaterol serum concentrations declined in a multi-phasic manner with an average terminal half-life ranging from 45.5 to 126 hours. The effective half-life, calculated from the accumulation of indacaterol after repeated dosing, ranged from 40 to 52 hours which is consistent with the observed time to steady state of approximately 12 to 14 days.

### Mometasone furoate

After intravenous bolus administration, mometasone furoate has a terminal elimination  $T_{1/2}$  of approximately 4.5 hours. A radiolabelled, orally inhaled dose is excreted mainly in the faeces (74%) and to a lesser extent in the urine (8%).

## Interactions

Concomitant administration of orally inhaled indacaterol and mometasone furoate under steady-state conditions did not affect the pharmacokinetics of either active substance.



### Linearity/non-linearity

Systemic exposure of mometasone furoate increased in a dose proportional manner following single and multiple doses of Atecura Breezhaler 125 mcg/62.5 mcg and 125 mcg/260 mcg in healthy subjects. A less than proportional increase in steady-state systemic exposure was noted in patients with asthma over the dose range of 125 mcg/62.5 mcg to 125 mcg/260 mcg. Dose proportionality assessments were not performed for indacaterol as only one dose was used across all dose strengths.

### Paediatric population

Atecura Breezhaler may be used in adolescent patients (12 years of age and older) at the same posology as in adults.

### Special populations

A population pharmacokinetic analysis in patients with asthma after inhalation of indacaterol/mometasone furoate indicated no significant effect of age, gender, body weight, smoking status, baseline estimated glomerular filtration rate (eGFR) and FEV<sub>1</sub> at baseline on the systemic exposure to indacaterol and mometasone furoate.

#### *Patients with renal impairment*

Due to the very low contribution of the urinary pathway to total body elimination of indacaterol and mometasone furoate, the effects of renal impairment on their systemic exposure have not been investigated (see section 4.2).

#### *Patients with hepatic impairment*

The effect of indacaterol/mometasone furoate has not been evaluated in subjects with hepatic impairment. However, studies have been conducted with the monotherapy components (see section 4.2).

#### *Indacaterol*

Patients with mild and moderate hepatic impairment showed no relevant changes in C<sub>max</sub> or AUC of indacaterol, nor did protein binding differ between mild and moderate hepatic impaired subjects and their healthy controls. No data are available for subjects with severe hepatic impairment.

#### *Mometasone furoate*

A study evaluating the administration of a single inhaled dose of 400 mcg mometasone furoate by dry powder inhaler to subjects with mild (n=4), moderate (n=4), and severe (n=4) hepatic impairment resulted in only 1 or 2 subjects in each group having detectable peak plasma concentrations of mometasone furoate (ranging from 50 to 105 pcg/ml). The observed peak plasma concentrations appear to increase with severity of hepatic impairment; however, the numbers of detectable levels (assay lower limit of quantification was 50 pcg/ml) were few.

#### *Other special populations*

There were no major differences in total systemic exposure (AUC) for both compounds between Japanese and Caucasian subjects. Insufficient pharmacokinetic data are available for other ethnicities or races.

### 5.3 Preclinical safety data

The non-clinical assessments of each monotherapy and of the combination product are presented below.

#### Indacaterol and mometasone furoate combination

The findings during the 13-week inhalation toxicity studies were predominantly attributable to the mometasone furoate component and were typical pharmacological effects of glucocorticoids. Increased heart rates associated with indacaterol were apparent in dogs after administration of indacaterol/mometasone furoate or indacaterol alone.

#### Indacaterol

Effects on the cardiovascular system attributable to the beta<sub>2</sub>-agonistic properties of indacaterol included tachycardia, arrhythmias and myocardial lesions in dogs. Mild irritation of the nasal cavity and larynx was seen in rodents.

Genotoxicity studies did not reveal any mutagenic or clastogenic potential.

Carcinogenicity was assessed in a two-year rat study and a six-month transgenic mouse study. Increased incidences of benign ovarian leiomyoma and focal hyperplasia of ovarian smooth muscle in rats were consistent with similar findings reported for other beta<sub>2</sub>-adrenergic agonists. No evidence of carcinogenicity was seen in mice.

All these findings occurred at exposures sufficiently in excess of those anticipated in humans.

Following subcutaneous administration in a rabbit study, adverse effects of indacaterol with respect to pregnancy and embryonal/foetal development could only be demonstrated at doses more than 500-fold those achieved following daily inhalation of 150 mcg in humans (based on AUC<sub>0-24 h</sub>).

Although indacaterol did not affect general reproductive performance in a rat fertility study, a decrease in the number of pregnant F1 offspring was observed in the peri- and post-natal developmental rat study at an exposure 14-fold higher than in humans treated with indacaterol. Indacaterol was not embryotoxic or teratogenic in rats or rabbits.

#### Mometasone furoate

All observed effects are typical of the glucocorticoid class of compounds and are related to exaggerated pharmacological effects of glucocorticoids. Mometasone furoate showed no genotoxic activity in a standard battery of *in vitro* and *in vivo* tests.

In carcinogenicity studies in mice and rats, inhaled mometasone furoate demonstrated no statistically significant increase in the incidence of tumours.

Like other glucocorticoids, mometasone furoate is a teratogen in rodents and rabbits. Effects noted were umbilical hernia in rats, cleft palate in mice and gallbladder agenesis, umbilical hernia and flexed front paws in rabbits. There were also reductions in maternal body weight gains, effects on foetal growth (lower foetal body weight and/or delayed ossification) in rats, rabbits and mice, and reduced offspring survival in mice. In studies of reproductive function, subcutaneous mometasone furoate at 15 mcg/kg prolonged gestation and difficult labour occurred, with a reduction in offspring survival and body weight.

Environmental risk assessment studies have shown that mometasone may pose a risk to surface water (see section 6.6).

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Capsule content

Lactose monohydrate

#### Capsule shell

Gelatin

Printing ink

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

3 years.

### **6.4 Special precautions for storage**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

### **6.5 Nature and contents of container**

Inhaler body and cap are made from acrylonitrile butadiene styrene, push buttons are made from methyl methacrylate acrylonitrile butadiene styrene. Needles and springs are made from stainless steel.

PA/Alu/PVC – Alu perforated unit-dose blister. Each blister contains 10 hard capsules.

#### Aectura Breezhaler 125 micrograms/62.5 micrograms inhalation powder, hard capsules

Single pack containing 10 x 1 or 30 x 1 hard capsules, together with 1 inhaler.

Pack containing 30 x 1 hard capsules, together with 1 inhaler and 1 sensor.

Multipacks containing 90 (3 packs of 30 x 1) hard capsules and 3 inhalers.

Multipacks containing 150 (15 packs of 10 x 1) hard capsules and 15 inhalers.

#### Aectura Breezhaler 125 micrograms/127.5 micrograms inhalation powder, hard capsules

Single pack containing 10 x 1 or 30 x 1 hard capsules, together with 1 inhaler.

Pack containing 30 x 1 hard capsules, together with 1 inhaler and 1 sensor.

Multipacks containing 90 (3 packs of 30 x 1) hard capsules and 3 inhalers.

Multipacks containing 150 (15 packs of 10 x 1) hard capsules and 15 inhalers.

#### Aectura Breezhaler 125 micrograms/260 micrograms inhalation powder, hard capsules

Single pack containing 10 x 1 or 30 x 1 hard capsules, together with 1 inhaler.

Pack containing 30 x 1 hard capsules, together with 1 inhaler and 1 sensor.

Multipacks containing 90 (3 packs of 30 x 1) hard capsules and 3 inhalers.

Multipacks containing 150 (15 packs of 10 x 1) hard capsules and 15 inhalers.

Not all pack sizes may be marketed.

## 6.6 Special precautions for disposal and other handling


The inhaler provided with each new prescription should be used. The inhaler in each pack should be disposed of after all capsules in that pack have been used.

This medicinal product may pose a risk to the environment (see section 5.3).

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.


### Instructions for handling and use

Please read the full **Instructions for Use** before using the Atecura Breezhaler.





Step 1b:  
**Open inhaler**




Step 1c:  
**Remove capsule**  
Separate one of the blisters from the blister card.  
Peel open the blister and remove the capsule.  
Do not push the capsule through the foil.  
Do not swallow the capsule.

You should hear a noise as the capsule is pierced.  
Only pierce the capsule once.



Step 2b:  
**Release side buttons**



Step 3b:  
**Inhale medicine deeply**  
Hold the inhaler as shown in the picture. Place the mouthpiece in your mouth and close your lips firmly around it.  
Do not press the side buttons.

Breathe in quickly and as deeply as you can. During inhalation you will hear a whirring noise. You may taste the medicine as you inhale.




Step 3c:  
**Hold breath**  
Hold your breath for up to 5 seconds.

Step 3d:  
**Rinse mouth**  
Rinse your mouth with water after each dose and spit it out.



**Powder remaining**



**Empty**



**Remove empty capsule**  
Put the empty capsule in your household waste. Close the inhaler and replace the cap.



Step 1d:

**Insert capsule**

Never place a capsule directly into the mouthpiece.



Step 1e:


**Close inhaler**

**Important Information**

- Aectura Breezhaler capsules must always be stored in the blister card and only removed immediately before use.
- Do not push the capsule through the foil to remove it from the blister.
- Do not swallow the capsule.
- Do not use the Aectura Breezhaler capsules with any other inhaler.
- Do not use the Aectura Breezhaler inhaler to take any other capsule medicine.
- Never place the capsule into your mouth or the mouthpiece of the inhaler.
- Do not press the side buttons more than once.
- Do not blow into the mouthpiece.
- Do not press the side buttons while inhaling through the mouthpiece.
- Do not handle capsules with wet hands.
- Never wash your inhaler with water.

Your Ateectura Breezhaler Inhaler pack contains:

- One Ateectura Breezhaler inhaler
- One or more blister cards, each containing 10 Ateectura Breezhaler capsules to be used in the inhaler



## Frequently Asked Questions

### Why didn't the inhaler make a noise when I inhaled?

The capsule may be stuck in the capsule chamber. If this happens, carefully loosen the capsule by tapping the base of the inhaler. Inhale the medicine again by repeating steps 3a to 3d.

### What should I do if there is powder left inside the capsule?

You have not received enough of your medicine. Close the inhaler and repeat steps 3a to 3d.

### I coughed after inhaling – does this matter?

This may happen. As long as the capsule is empty you have received enough of your medicine.

### I felt small pieces of the capsule on my tongue – does this matter?

This can happen. It is not harmful. The chances of the capsule breaking into small pieces will be increased if the capsule is pierced more than once.

## Cleaning the inhaler

Wipe the mouthpiece inside and outside with a clean, dry, lint-free cloth to remove any powder residue. Keep the inhaler dry. Never wash your inhaler with water.

## Disposing of the inhaler after use

Each inhaler should be disposed of after all capsules have been used. Ask your pharmacist how to dispose of medicines and inhalers that are no longer required.

For detailed instructions on use of the sensor and the App, see the Instructions for Use provided in the sensor pack and the App.

## 7. MARKETING AUTHORISATION HOLDER

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**8. MARKETING AUTHORISATION NUMBER(S)**

Aectura Breezhaler 125 micrograms/62.5 micrograms inhalation powder, hard capsules

EU/1/20/1439/001-004

EU/1/20/1439/013

Aectura Breezhaler 125 micrograms/127.5 micrograms inhalation powder, hard capsules

EU/1/20/1439/005-008

EU/1/20/1439/014

Aectura Breezhaler 125 micrograms/260 micrograms inhalation powder, hard capsules

EU/1/20/1439/009-012

EU/1/20/1439/015

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

30 May 2020

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.



## **ANNEX II**

- A. MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE**

### Name and address of the manufacturers responsible for batch release

Novartis Farmacéutica, S.A.  
Gran Via de les Corts Catalanes, 764  
08013 Barcelona  
Spain

Novartis Pharma GmbH  
Roonstraße 25  
D-90429 Nuremberg  
Germany

Novartis Pharma GmbH  
Sophie-Germain-Strasse 10  
90443 Nuremberg  
Germany

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to medical prescription.

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

### **• Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

The marketing authorisation holder (MAH) shall submit the first PSUR for this product within 6 months following authorisation.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

### **• Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

## **PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

### **OUTER CARTON OF UNIT PACK**

#### **1. NAME OF THE MEDICINAL PRODUCT**

Aectura Breezhaler 125 micrograms/62.5 micrograms inhalation powder, hard capsules  
indacaterol/mometasone furoate

#### **2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each delivered dose contains 125 micrograms indacaterol (as acetate) and 62.5 micrograms mometasone furoate.

#### **3. LIST OF EXCIPIENTS**

Also contains lactose. See package leaflet for further information.

#### **4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

10 x 1 capsules + 1 inhaler

30 x 1 capsules + 1 inhaler

#### **5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

For use only with the inhaler provided in the pack.

Do not swallow capsules.

Inhalation use

‘QR code to be included’

Scan for more or visit: [www.breezhaler-asthma.eu/atectura](http://www.breezhaler-asthma.eu/atectura)

#### **6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

#### **7. OTHER SPECIAL WARNING(S), IF NECESSARY**

#### **8. EXPIRY DATE**

EXP

The inhaler in each pack should be disposed of after all capsules in that pack have been used.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/20/1439/001

10 x 1 capsules + 1 inhaler

EU/1/20/1439/002

30 x 1 capsules + 1 inhaler

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Aectura Breezhaler 125 micrograms/62.5 micrograms

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****OUTER CARTON OF UNIT PACK CONTAINING SENSOR****1. NAME OF THE MEDICINAL PRODUCT**

Aectura Breezhaler 125 micrograms/62.5 micrograms inhalation powder, hard capsules  
indacaterol/mometasone furoate

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each delivered dose contains 125 micrograms indacaterol (as acetate) and 62.5 micrograms  
mometasone furoate.

**3. LIST OF EXCIPIENTS**

Also contains lactose. See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

30 x 1 capsules + 1 inhaler + 1 sensor for adults only

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
For use only with the inhaler provided in the pack.  
Do not swallow capsules.  
Inhalation use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

The inhaler in each pack should be disposed of after all capsules in that pack have been used.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/20/1439/013

30 x 1 capsules + 1 inhaler + 1 sensor

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Aectura Breezhaler 125 micrograms/62.5 micrograms

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN



**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****INTERMEDIATE CARTON OF UNIT PACK CONTAINING SENSOR****1. NAME OF THE MEDICINAL PRODUCT**

Aectura Breezhaler 125 micrograms/62.5 micrograms inhalation powder, hard capsules  
indacaterol/mometasone furoate

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each delivered dose contains 125 micrograms indacaterol (as acetate) and 62.5 micrograms  
mometasone furoate.

**3. LIST OF EXCIPIENTS**

Also contains lactose. See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

30 x 1 capsules + 1 inhaler

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
For use only with the inhaler provided in the pack.  
Do not swallow capsules.  
Inhalation use

‘QR code to be included’  
Scan for more or visit: [www.breezhaler-asthma.eu/atectura](http://www.breezhaler-asthma.eu/atectura)

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP  
The inhaler in each pack should be disposed of after all capsules in that pack have been used.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/20/1439/013

30 x 1 capsules + 1 inhaler + 1 sensor

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Aectura Breezhaler 125 micrograms/62.5 micrograms

**17. UNIQUE IDENTIFIER – 2D BARCODE****18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****OUTER CARTON OF MULTIPACK (INCLUDING BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Aectura Breezhaler 125 micrograms/62.5 micrograms inhalation powder, hard capsules  
indacaterol/mometasone furoate

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each delivered dose contains 125 micrograms indacaterol (as acetate) and 62.5 micrograms mometasone furoate.

**3. LIST OF EXCIPIENTS**

Also contains lactose. See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

Multipack: 90 (3 packs of 30 x 1) capsules + 3 inhalers.

Multipack: 150 (15 packs of 10 x 1) capsules + 15 inhalers.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

For use only with the inhaler provided in the pack.

Do not swallow capsules.

Inhalation use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

The inhaler in each pack should be disposed of after all capsules in that pack have been used.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/20/1439/003

90 (3 packs of 30 x 1) capsules + 3 inhalers

EU/1/20/1439/004

150 (15 packs of 10 x 1) capsules + 15 inhalers

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Aectura Breezhaler 125 micrograms/62.5 micrograms

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

## **PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

### **INTERMEDIATE CARTON OF MULTIPACK (WITHOUT BLUE BOX)**

#### **1. NAME OF THE MEDICINAL PRODUCT**

Aectura Breezhaler 125 micrograms/62.5 micrograms inhalation powder, hard capsules  
indacaterol/mometasone furoate

#### **2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each delivered dose contains 125 micrograms indacaterol (as acetate) and 62.5 micrograms mometasone furoate.

#### **3. LIST OF EXCIPIENTS**

Also contains lactose. See package leaflet for further information.

#### **4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

10 x 1 capsules + 1 inhaler. Component of a multipack. Not to be sold separately.

30 x 1 capsules + 1 inhaler. Component of a multipack. Not to be sold separately.

#### **5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

For use only with the inhaler provided in the pack.

Do not swallow capsules.

Inhalation use

‘QR code to be included’

Scan for more or visit: [www.breezhaler-asthma.eu/atectura](http://www.breezhaler-asthma.eu/atectura)

#### **6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

#### **7. OTHER SPECIAL WARNING(S), IF NECESSARY**

#### **8. EXPIRY DATE**

EXP

The inhaler in each pack should be disposed of after all capsules in that pack have been used.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/20/1439/003

90 (3 packs of 30 x 1) capsules + 3 inhalers

EU/1/20/1439/004

150 (15 packs of 10 x 1) capsules + 15 inhalers

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Aectura Breezhaler 125 micrograms/62.5 micrograms

**17. UNIQUE IDENTIFIER – 2D BARCODE****18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**INNER LID OF**

- **OUTER CARTON OF UNIT PACK**
- **INTERMEDIATE CARTON OF UNIT PACK CONTAINING SENSOR, AND**
- **INTERMEDIATE CARTON OF MULTIPACK**

**1. OTHER**

- 1 Insert
  - 2 Pierce and release
  - 3 Inhale deeply
- Check Check capsule is empty

Read the leaflet before use.

<b>PARTICULARS TO APPEAR ON THE OUTER PACKAGING</b>
---

<b>INNER LID OF OUTER CARTON OF UNIT PACK CONTAINING SENSOR</b>
---

<b>1. OTHER</b>
-----------------

Please read the package leaflet in the Atecura Breezhaler pack and the instructions for use provided in the sensor pack.



<b>MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS</b>
--

<b>BLISTERS</b>
-----------------

<b>1. NAME OF THE MEDICINAL PRODUCT</b>
---

Aectura Breezhaler 125 mcg/62.5 mcg inhalation powder  
indacaterol/mometasone furoate

<b>2. NAME OF THE MARKETING AUTHORISATION HOLDER</b>
--

Novartis Europharm Limited

<b>3. EXPIRY DATE</b>
-----------------------

EXP

<b>4. BATCH NUMBER</b>
------------------------

Lot

<b>5. OTHER</b>
-----------------

Inhalation use only

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****OUTER CARTON OF UNIT PACK****1. NAME OF THE MEDICINAL PRODUCT**

Aectura Breezhaler 125 micrograms/127.5 micrograms inhalation powder, hard capsules  
indacaterol/mometasone furoate

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each delivered dose contains 125 micrograms indacaterol (as acetate) and 127.5 micrograms mometasone furoate.

**3. LIST OF EXCIPIENTS**

Also contains lactose. See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

10 x 1 capsules + 1 inhaler

30 x 1 capsules + 1 inhaler

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

For use only with the inhaler provided in the pack.

Do not swallow capsules.

Inhalation use

‘QR code to be included’

Scan for more or visit: [www.breezhaler-asthma.eu/atectura](http://www.breezhaler-asthma.eu/atectura)

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

The inhaler in each pack should be disposed of after all capsules in that pack have been used.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/20/1439/005

10 x 1 capsules + 1 inhaler

EU/1/20/1439/006

30 x 1 capsules + 1 inhaler

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Aectura Breezhaler 125 micrograms/127.5 micrograms

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****OUTER CARTON OF UNIT PACK CONTAINING SENSOR****1. NAME OF THE MEDICINAL PRODUCT**

Aectura Breezhaler 125 micrograms/127.5 micrograms inhalation powder, hard capsules  
indacaterol/mometasone furoate

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each delivered dose contains 125 micrograms indacaterol (as acetate) and 127.5 micrograms  
mometasone furoate.

**3. LIST OF EXCIPIENTS**

Also contains lactose. See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

30 x 1 capsules + 1 inhaler + 1 sensor for adults only

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
For use only with the inhaler provided in the pack.  
Do not swallow capsules.  
Inhalation use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

The inhaler in each pack should be disposed of after all capsules in that pack have been used.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/20/1439/014

30 x 1 capsules + 1 inhaler + 1 sensor

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Aectura Breezhaler 125 micrograms/127.5 micrograms

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****INTERMEDIATE CARTON OF UNIT PACK CONTAINING SENSOR****1. NAME OF THE MEDICINAL PRODUCT**

Aectura Breezhaler 125 micrograms/127.5 micrograms inhalation powder, hard capsules  
indacaterol/mometasone furoate

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each delivered dose contains 125 micrograms indacaterol (as acetate) and 127.5 micrograms mometasone furoate.

**3. LIST OF EXCIPIENTS**

Also contains lactose. See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

30 x 1 capsules + 1 inhaler

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
For use only with the inhaler provided in the pack.  
Do not swallow capsules.  
Inhalation use

‘QR code to be included’  
Scan for more or visit: [www.breezhaler-asthma.eu/atectura](http://www.breezhaler-asthma.eu/atectura)

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP  
The inhaler in each pack should be disposed of after all capsules in that pack have been used.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/20/1439/014

30 x 1 capsules + 1 inhaler + 1 sensor

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Aectura Breezhaler 125 micrograms/127.5 micrograms

**17. UNIQUE IDENTIFIER – 2D BARCODE****18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****OUTER CARTON OF MULTIPACK (INCLUDING BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Aectura Breezhaler 125 micrograms/127.5 micrograms inhalation powder, hard capsules  
indacaterol/mometasone furoate

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each delivered dose contains 125 micrograms indacaterol (as acetate) and 127.5 micrograms mometasone furoate.

**3. LIST OF EXCIPIENTS**

Also contains lactose. See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

Multipack: 90 (3 packs of 30 x 1) capsules + 3 inhalers.

Multipack: 150 (15 packs of 10 x 1) capsules + 15 inhalers.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

For use only with the inhaler provided in the pack.

Do not swallow capsules.

Inhalation use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

The inhaler in each pack should be disposed of after all capsules in that pack have been used.



**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/20/1439/007

90 (3 packs of 30 x 1) capsules + 3 inhalers

EU/1/20/1439/008

150 (15 packs of 10 x 1) capsules + 15 inhalers

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Aectura Breezhaler 125 micrograms/127.5 micrograms

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****INTERMEDIATE CARTON OF MULTIPACK (WITHOUT BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Aectura Breezhaler 125 micrograms/127.5 micrograms inhalation powder, hard capsules  
indacaterol/mometasone furoate

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each delivered dose contains 125 micrograms indacaterol (as acetate) and 127.5 micrograms mometasone furoate.

**3. LIST OF EXCIPIENTS**

Also contains lactose. See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

10 x 1 capsules + 1 inhaler. Component of a multipack. Not to be sold separately.  
30 x 1 capsules + 1 inhaler. Component of a multipack. Not to be sold separately.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
For use only with the inhaler provided in the pack.  
Do not swallow capsules.  
Inhalation use

‘QR code to be included’  
Scan for more or visit: [www.breezhaler-asthma.eu/atectura](http://www.breezhaler-asthma.eu/atectura)

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP  
The inhaler in each pack should be disposed of after all capsules in that pack have been used.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/20/1439/007

90 (3 packs of 30 x 1) capsules + 3 inhalers

EU/1/20/1439/008

150 (15 packs of 10 x 1) capsules + 15 inhalers

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Aectura Breezhaler 125 micrograms/127.5 micrograms

**17. UNIQUE IDENTIFIER – 2D BARCODE****18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**INNER LID OF**

- **OUTER CARTON OF UNIT PACK**
- **INTERMEDIATE CARTON OF UNIT PACK CONTAINING SENSOR, AND**
- **INTERMEDIATE CARTON OF MULTIPACK**

**1. OTHER**

- 1 Insert
  - 2 Pierce and release
  - 3 Inhale deeply
- Check Check capsule is empty

Read the leaflet before use.

<b>PARTICULARS TO APPEAR ON THE OUTER PACKAGING</b>
---

<b>INNER LID OF OUTER CARTON OF UNIT PACK CONTAINING SENSOR</b>
---

<b>1. OTHER</b>
-----------------

Please read the package leaflet in the Atecura Breezhaler pack and the instructions for use provided in the sensor pack.

<b>MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS</b>
--

<b>BLISTERS</b>
-----------------

<b>1. NAME OF THE MEDICINAL PRODUCT</b>
---

Aectura Breezhaler 125 mcg/127.5 mcg inhalation powder  
indacaterol/mometasone furoate

<b>2. NAME OF THE MARKETING AUTHORISATION HOLDER</b>
--

Novartis Europharm Limited

<b>3. EXPIRY DATE</b>
-----------------------

EXP

<b>4. BATCH NUMBER</b>
------------------------

Lot

<b>5. OTHER</b>
-----------------

Inhalation use only

## **PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

### **OUTER CARTON OF UNIT PACK**

#### **1. NAME OF THE MEDICINAL PRODUCT**

Aectura Breezhaler 125 micrograms/260 micrograms inhalation powder, hard capsules  
indacaterol/mometasone furoate

#### **2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each delivered dose contains 125 micrograms indacaterol (as acetate) and 260 micrograms mometasone furoate.

#### **3. LIST OF EXCIPIENTS**

Also contains lactose. See package leaflet for further information.

#### **4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

10 x 1 capsules + 1 inhaler

30 x 1 capsules + 1 inhaler

#### **5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

For use only with the inhaler provided in the pack.

Do not swallow capsules.

Inhalation use

‘QR code to be included’

Scan for more or visit: [www.breezhaler-asthma.eu/atectura](http://www.breezhaler-asthma.eu/atectura)

#### **6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

#### **7. OTHER SPECIAL WARNING(S), IF NECESSARY**

#### **8. EXPIRY DATE**

EXP

The inhaler in each pack should be disposed of after all capsules in that pack have been used.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/20/1439/009

10 x 1 capsules + 1 inhaler

EU/1/20/1439/010

30 x 1 capsules + 1 inhaler

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Aectura Breezhaler 125 micrograms/260 micrograms

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN



**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****OUTER CARTON OF UNIT PACK CONTAINING SENSOR****1. NAME OF THE MEDICINAL PRODUCT**

Aectura Breezhaler 125 micrograms/260 micrograms inhalation powder, hard capsules  
indacaterol/mometasone furoate

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each delivered dose contains 125 micrograms indacaterol (as acetate) and 260 micrograms  
mometasone furoate.

**3. LIST OF EXCIPIENTS**

Also contains lactose. See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

30 x 1 capsules + 1 inhaler + 1 sensor for adults only

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
For use only with the inhaler provided in the pack.  
Do not swallow capsules.  
Inhalation use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

The inhaler in each pack should be disposed of after all capsules in that pack have been used.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/20/1439/015

30 x 1 capsules + 1 inhaler + 1 sensor

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Aectura Breezhaler 125 micrograms/260 micrograms

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****INTERMEDIATE CARTON OF UNIT PACK CONTAINING SENSOR****1. NAME OF THE MEDICINAL PRODUCT**

Aectura Breezhaler 125 micrograms/260 micrograms inhalation powder, hard capsules  
indacaterol/mometasone furoate

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each delivered dose contains 125 micrograms indacaterol (as acetate) and 260 micrograms  
mometasone furoate.

**3. LIST OF EXCIPIENTS**

Also contains lactose. See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

30 x 1 capsules + 1 inhaler

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
For use only with the inhaler provided in the pack.  
Do not swallow capsules.  
Inhalation use

‘QR code to be included’  
Scan for more or visit: [www.breezhaler-asthma.eu/atectura](http://www.breezhaler-asthma.eu/atectura)

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP  
The inhaler in each pack should be disposed of after all capsules in that pack have been used.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/20/1439/015

30 x 1 capsules + 1 inhaler + 1 sensor

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Aectura Breezhaler 125 micrograms/260 micrograms

**17. UNIQUE IDENTIFIER – 2D BARCODE****18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****OUTER CARTON OF MULTIPACK (INCLUDING BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Aectura Breezhaler 125 micrograms/260 micrograms inhalation powder, hard capsules  
indacaterol/mometasone furoate

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each delivered dose contains 125 micrograms indacaterol (as acetate) and 260 micrograms  
mometasone furoate.

**3. LIST OF EXCIPIENTS**

Also contains lactose. See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

Multipack: 90 (3 packs of 30 x 1) capsules + 3 inhalers.

Multipack: 150 (15 packs of 10 x 1) capsules + 15 inhalers.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

For use only with the inhaler provided in the pack.

Do not swallow capsules.

Inhalation use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

The inhaler in each pack should be disposed of after all capsules in that pack have been used.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/20/1439/011

90 (3 packs of 30 x 1) capsules + 3 inhalers

EU/1/20/1439/012

150 (15 packs of 10 x 1) capsules + 15 inhalers

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Aectura Breezhaler 125 micrograms/260 micrograms

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****INTERMEDIATE CARTON OF MULTIPACK (WITHOUT BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Aectura Breezhaler 125 micrograms/260 micrograms inhalation powder, hard capsules  
indacaterol/mometasone furoate

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each delivered dose contains 125 micrograms indacaterol (as acetate) and 260 micrograms  
mometasone furoate.

**3. LIST OF EXCIPIENTS**

Also contains lactose. See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

10 x 1 capsules + 1 inhaler. Component of a multipack. Not to be sold separately.  
30 x 1 capsules + 1 inhaler. Component of a multipack. Not to be sold separately.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
For use only with the inhaler provided in the pack.  
Do not swallow capsules.  
Inhalation use

‘QR code to be included’  
Scan for more or visit: [www.breezhaler-asthma.eu/atectura](http://www.breezhaler-asthma.eu/atectura)

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP  
The inhaler in each pack should be disposed of after all capsules in that pack have been used.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/20/1439/011

90 (3 packs of 30 x 1) capsules + 3 inhalers

EU/1/20/1439/012

150 (15 packs of 10 x 1) capsules + 15 inhalers

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Aectura Breezhaler 125 micrograms/260 micrograms

**17. UNIQUE IDENTIFIER – 2D BARCODE****18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**



**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**INNER LID OF**

- **OUTER CARTON OF UNIT PACK**
- **INTERMEDIATE CARTON OF UNIT PACK CONTAINING SENSOR,AND**
- **INTERMEDIATE CARTON OF MULTIPACK**

**1. OTHER**

- 1 Insert
  - 2 Pierce and release
  - 3 Inhale deeply
- Check Check capsule is empty

Read the leaflet before use.

<b>PARTICULARS TO APPEAR ON THE OUTER PACKAGING</b>
---

<b>INNER LID OF OUTER CARTON OF UNIT PACK CONTAINING SENSOR</b>
---

<b>1. OTHER</b>
-----------------

Please read the package leaflet in the Atecura Breezhaler pack and the instructions for use provided in the sensor pack.

<b>MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS</b>
--

<b>BLISTERS</b>
-----------------

<b>1. NAME OF THE MEDICINAL PRODUCT</b>
---

Aectura Breezhaler 125 mcg/260 mcg inhalation powder  
indacaterol/mometasone furoate

<b>2. NAME OF THE MARKETING AUTHORISATION HOLDER</b>
--

Novartis Europharm Limited

<b>3. EXPIRY DATE</b>
-----------------------

EXP

<b>4. BATCH NUMBER</b>
------------------------

Lot

<b>5. OTHER</b>
-----------------

Inhalation use only

## **B. PACKAGE LEAFLET**

## **Package leaflet: Information for the user**

**Aectura Breezhaler 125 micrograms/62.5 micrograms inhalation powder, hard capsules**  
**Aectura Breezhaler 125 micrograms/127.5 micrograms inhalation powder, hard capsules**  
**Aectura Breezhaler 125 micrograms/260 micrograms inhalation powder, hard capsules**  
indacaterol/mometasone furoate

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

### **What is in this leaflet:**

1. What Aectura Breezhaler is and what it is used for
  2. What you need to know before you use Aectura Breezhaler
  3. How to use Aectura Breezhaler
  4. Possible side effects
  5. How to store Aectura Breezhaler
  6. Contents of the pack and other information
- Instructions for use of Aectura Breezhaler inhaler

### **1. What Aectura Breezhaler is and what it is used for**

#### **What Aectura Breezhaler is and how it works**

Aectura Breezhaler contains two active substances called indacaterol and mometasone furoate.

Indacaterol belongs to a group of medicines called bronchodilators. It relaxes the muscles of the small airways in the lungs. This helps to open the airways and makes it easier for air to get in and out of the lungs. When it is taken regularly, it helps the small airways to remain open.

Mometasone furoate belongs to a group of medicines called corticosteroids (or steroids). Corticosteroids reduce the swelling and irritation (inflammation) in the small airways in the lungs and so gradually ease breathing problems. Corticosteroids also help to prevent attacks of asthma.

#### **What Aectura Breezhaler is used for**

Aectura Breezhaler is used regularly as treatment for asthma in adults and adolescents (12 years of age and older).

Asthma is a serious, long-term lung disease where the muscles surrounding the smaller airways become tight (bronchoconstriction) and inflamed. Symptoms come and go and include shortness of breath, wheezing, chest tightness and cough.

You should use Aectura Breezhaler every day and not only when you have breathing problems or other symptoms of asthma. This will ensure that it controls your asthma properly. Do not use this medicine to relieve a sudden attack of breathlessness or wheezing.

If you have any questions about how Aectura Breezhaler works or why this medicine has been prescribed for you, ask your doctor.

## **2. What you need to know before you use Atecura Breezhaler**

Follow all the doctor's instructions carefully.

### **Do not use Atecura Breezhaler**

- if you are allergic to indacaterol, mometasone furoate or any of the other ingredients of this medicine (listed in section 6). If you think you may be allergic, ask your doctor for advice.

### **Warnings and precautions**

Talk to your doctor, pharmacist or nurse **before** using Atecura Breezhaler if any of the following applies to you:

- if you have heart problems, including an irregular or fast heartbeat.
- if you have thyroid gland problems.
- if you have ever been told you have diabetes or high blood sugar.
- if you suffer from seizures or fits.
- if you have a low level of potassium in your blood.
- if you have severe liver problems.
- if you have tuberculosis (TB) of the lung, or any long-standing or untreated infections.

### **During treatment with Atecura Breezhaler**

**Stop using this medicine and get medical help immediately** if you have any of the following:

- tightness of the chest, coughing, wheezing or breathlessness immediately after using Atecura Breezhaler (signs the medicine is unexpectedly tightening the airways, known as paradoxical bronchospasm).
- difficulty breathing or swallowing, swelling of the tongue, lips or face, skin rash, itching and hives (signs of allergic reaction).

### **Children and adolescents**

Do not give this medicine to children below 12 years of age because it has not been studied in this age group.

### **Other medicines and Atecura Breezhaler**

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines. In particular, tell your doctor or pharmacist if you are using:

- medicines that decrease the level of potassium in your blood. These include diuretics (which increase urine production and can be used to treat high blood pressure, e.g. hydrochlorothiazide), other bronchodilators such as methylxanthines used for breathing problems (e.g. theophylline) or corticosteroids (e.g. prednisolone).
- tricyclic antidepressants or monoamine oxidase inhibitors (medicines used in the treatment of depression).
- any medicines that may be similar to Atecura Breezhaler (contain similar active substances); using them together may increase the risk of possible side effects.
- medicines called beta blockers used to treat high blood pressure or other heart problems (e.g. propranolol) or to treat glaucoma (e.g. timolol).
- ketoconazole or itraconazole (medicines used to treat fungal infections)
- ritonavir, nelfinavir or cobicistat (medicines used to treat HIV infection).

### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you whether you can use Atecura Breezhaler.

### **Driving and using machines**

It is unlikely that this medicine will affect your ability to drive and use machines.

### **Aectura Breezhaler contains lactose**

This medicine contains about 25 mg of lactose per capsule. If you have been told by your doctor that you have an intolerance to some sugars, speak with your doctor before taking this medicine.

### **3. How to use Aectura Breezhaler**

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

#### **How much Aectura Breezhaler to inhale**

There are three different strengths of Aectura Breezhaler capsules. Your doctor will decide which is best for you.

The usual dose is to inhale the content of one capsule each day. You only need to use the medicine once a day. Do not use more than your doctor tells you to use.

You should use Aectura Breezhaler every day, even when your asthma is not troubling you.

#### **When to inhale Aectura Breezhaler**

Inhale Aectura Breezhaler at the same time each day. This will help control your symptoms throughout the day and night. It will also help you to remember to use it.

#### **How to inhale Aectura Breezhaler**

- Aectura Breezhaler is for inhalation use.
- In this pack, you will find an inhaler and capsules that contain the medicine. The inhaler enables you to inhale the medicine in the capsule. Only use the capsules with the inhaler provided in this pack. The capsules should remain in the blister until you need to use them.
- Peel the backing away from the blister to open it, **do not push the capsule through the foil.**
- When you start a new pack, use the new inhaler supplied in this new pack.
- Dispose of the inhaler in each pack after all capsules in that pack have been used.
- Do not swallow the capsules.
- **Please read the instructions for use on the other side of this leaflet for more information on how to use the inhaler.**

#### **If your pack contains a sensor for Aectura Breezhaler**

- The sensor for Aectura Breezhaler is for adult use only. It should not be used in adolescents (12 years of age and older) as the App does not have patient consent verification or caregiver account functionality.
- The sensor and App are not required in order for you to take your medicine. The sensor does not need to be connected to the App when taking your medicine.
- Your doctor will decide whether use of the sensor and App is appropriate for you.
- The electronic sensor for Aectura Breezhaler is to be attached to the base of the Aectura Breezhaler inhaler.
- The sensor confirms that you have used the Aectura Breezhaler inhaler by recording and monitoring its actuations and the whirring noise of the spinning capsule during inhalation but it will not monitor whether you have received the dose of your medicine.
- The sensor is to be used with the Propeller App on your smart phone or another suitable device. The sensor links with the Propeller App via Bluetooth.
- Please read the instructions for use provided in the sensor pack and the App for more information on how to use the sensor for Aectura Breezhaler and the App.
- After all Aectura Breezhaler capsules in a pack have been used, move the sensor to the new inhaler in the next Aectura Breezhaler pack.

#### **If your symptoms do not improve**

If your asthma is not getting better or if it gets worse after you have started using Aectura Breezhaler, talk to your doctor.

**If you use more Ateectura Breezhaler than you should**

If you accidentally inhale too much of this medicine, contact your doctor or hospital for advice immediately. You may need medical attention.

**If you forget to use Ateectura Breezhaler**

If you forget to inhale a dose at the usual time, inhale one as soon as possible on that day. Then inhale the next dose at the usual time on the next day. Do not inhale two doses on the same day.

**If you stop using Ateectura Breezhaler**

Do not stop using Ateectura Breezhaler unless your doctor tells you to. Your asthma symptoms may come back if you stop using it.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

**4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Some side effects could be serious**

Stop using Ateectura Breezhaler and get medical help immediately if you have any of the following:

**Common:** may affect up to 1 in every 10 people

- difficulty breathing or swallowing, swelling of the tongue, lips, or face, skin rash, itching and hives (signs of allergic reaction).

**Uncommon:** may affect up to 1 in every 100 people

- swelling mainly of the tongue, lips, face or throat (possible signs of angioedema).

**Other side effects**

Other side effects include the following listed below. If these side effects become severe, please tell your doctor, pharmacist or nurse.

**Very common:** may affect more than 1 in 10 people

- sore throat
- runny nose
- sudden difficulty breathing and feeling of tightness in chest with wheezing or coughing

**Common:** may affect up to 1 in every 10 people

- voice alteration (hoarseness)
- blocked nose
- sneezing, cough
- headache
- pain in muscles, bones or joints (signs of musculoskeletal pain)

**Uncommon:** may affect up to 1 in every 100 people

- fast heart beat
- oral thrush (sign of candidiasis)
- high level of sugar in the blood
- muscle spasm
- skin itching
- rash
- clouding of the lens of your eyes (signs of cataract)
- blurred vision



## Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store Ateectura Breezhaler

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton and blister after “EXP”. The expiry date refers to the last day of that month.
- Do not store above 30°C.
- Store the capsules in the original blister in order to protect from light and moisture, and do not remove until immediately before use.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.
- If your pack contains an electronic sensor for Ateectura Breezhaler, see the Instructions for Use provided in the sensor pack for detailed instructions on how to store and when to discard it.

## 6. Contents of the pack and other information

### What Ateectura Breezhaler contains

- The active substances are indacaterol (as acetate) and mometasone furoate.

#### Ateectura Breezhaler 125 micrograms/62.5 micrograms

Each capsule contains 173 micrograms of indacaterol acetate (equivalent to 150 micrograms of indacaterol) and 80 micrograms of mometasone furoate. The delivered dose (the dose that leaves the mouthpiece of the inhaler) is equivalent to 125 micrograms of indacaterol and 62.5 micrograms of mometasone furoate.

#### Ateectura Breezhaler 125 micrograms/127.5 micrograms

Each capsule contains 173 micrograms of indacaterol acetate (equivalent to 150 micrograms of indacaterol) and 160 micrograms of mometasone furoate. The delivered dose (the dose that leaves the mouthpiece of the inhaler) is equivalent to 125 micrograms of indacaterol and 127.5 micrograms of mometasone furoate.

#### Ateectura Breezhaler 125 micrograms/260 micrograms

Each capsule contains 173 micrograms of indacaterol acetate (equivalent to 150 micrograms of indacaterol) and 320 micrograms of mometasone furoate. The delivered dose (the dose that leaves the mouthpiece of the inhaler) is equivalent to 125 micrograms of indacaterol and 260 micrograms of mometasone furoate.

- The other ingredient is lactose monohydrate (see “Ateectura Breezhaler contains lactose” in section 2).

**What Ateectura Breezhaler looks like and content of the pack**

In this pack, you will find an inhaler together with capsules in blisters. Some packs also contain a sensor device. The capsules are transparent and contain a white powder.

- Ateectura Breezhaler 125 micrograms/62.5 micrograms capsules have a blue product code “IM150-80” printed above one blue bar on the body with a logo printed in blue and surrounded by two blue bars on the cap.
- Ateectura Breezhaler 125 micrograms/127.5 micrograms capsules have a grey product code “IM150-160” printed on the body with a logo printed in grey on the cap.
- Ateectura Breezhaler 125 micrograms/260 micrograms capsules have a black product code “IM150-320” printed above two black bars on the body with a logo printed in black and surrounded by two black bars on the cap.

The following pack sizes are available:

Single pack containing 10 x 1 or 30 x 1 hard capsules, together with 1 inhaler.

Pack containing 30 x 1 hard capsules, together with 1 inhaler and 1 sensor.

Multipacks comprising 3 cartons, each containing 30 x 1 hard capsules together with 1 inhaler.

Multipacks comprising 15 cartons, each containing 10 x 1 hard capsules together with 1 inhaler.

Not all pack sizes may be available in your country.

**Marketing Authorisation Holder**

Novartis Europharm Limited  
Vista Building  
Elm Park, Merrion Road  
Dublin 4  
Ireland

**Manufacturer**

Novartis Farmacéutica, S.A.  
Gran Via de les Corts Catalanes, 764  
08013 Barcelona  
Spain

Novartis Pharma GmbH  
Roonstraße 25  
D-90429 Nuremberg  
Germany

Novartis Pharma GmbH  
Sophie-Germain-Strasse 10  
90443 Nuremberg  
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

Novartis Pharma N.V.  
Tél/Tel: +32 2 246 16 11

**България**

Novartis Bulgaria EOOD  
Тел: +359 2 489 98 28

**Česká republika**

Novartis s.r.o.  
Tel: +420 225 775 111

**Danmark**

Novartis Healthcare A/S  
Tlf: +45 39 16 84 00

**Deutschland**

**APONTIS PHARMA** Deutschland GmbH & Co.  
KG  
Tel: +49 2173 8955 4949

**Eesti**

SIA Novartis Baltics Eesti filiaal  
Tel: +372 66 30 810

**Ελλάδα**

Novartis (Hellas) A.E.B.E.  
Τηλ: +30 210 281 17 12

**España**

Laboratorios Menarini, S.A.  
Tel: +34 93 462 88 00

**France**

Novartis Pharma S.A.S.  
Tél: +33 1 55 47 66 00

**Hrvatska**

Novartis Hrvatska d.o.o.  
Tel. +385 1 6274 220

**Ireland**

Novartis Ireland Limited  
Tel: +353 1 260 12 55

**Ísland**

Vistor hf.  
Sími: +354 535 7000

**Italia**

Novartis Farma S.p.A.  
Tel: +39 02 96 54 1

**Lietuva**

SIA Novartis Baltics Lietuvos filialas  
Tel: +370 5 269 16 50

**Luxembourg/Luxemburg**

Novartis Pharma N.V.  
Tél/Tel: +32 2 246 16 11

**Magyarország**

Novartis Hungária Kft.  
Tel.: +36 1 457 65 00

**Malta**

Novartis Pharma Services Inc.  
Tel: +356 2122 2872

**Nederland**

Novartis Pharma B.V.  
Tel: +31 88 04 52 111

**Norge**

Novartis Norge AS  
Tlf: +47 23 05 20 00

**Österreich**

Novartis Pharma GmbH  
Tel: +43 1 86 6570

**Polska**

Novartis Poland Sp. z o.o.  
Tel.: +48 22 375 4888

**Portugal**

Novartis Farma - Produtos Farmacêuticos, S.A.  
Tel: +351 21 000 8600

**România**

Novartis Pharma Services Romania SRL  
Tel: +40 21 31299 01

**Slovenija**

Novartis Pharma Services Inc.  
Tel: +386 1 300 75 50

**Slovenská republika**

Novartis Slovakia s.r.o.  
Tel: +421 2 5542 5439

**Suomi/Finland**

Novartis Finland Oy  
Puh/Tel: +358 (0)10 6133 200

**Κύπρος**

Novartis Pharma Services Inc.  
Τηλ: +357 22 690 690

**Sverige**

Novartis Sverige AB  
Tel: +46 8 732 32 00

**Latvija**

SIA Novartis Baltics  
Tel: +371 67 887 070

**United Kingdom (Northern Ireland)**

Novartis Ireland Limited  
Tel: +44 1276 698370

**This leaflet was last revised in**













**Other sources of information**


Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>.

## Instructions for Use of Atecura Breezhaler

Please read full instructions for use of Atecura Breezhaler inhaler before use. These instructions are also available by scanning the QR code or visiting [www.breezhaler-asthma.eu/atecura](http://www.breezhaler-asthma.eu/atecura)

“QR code to be included”.

 <p><b>Insert</b></p> <p><b>1</b></p>	 <p><b>Pierce and release</b></p> <p><b>2</b></p>	 <p><b>Inhale deeply</b></p> <p><b>3</b></p>	 <p><b>Check capsule is empty</b></p> <p><b>Check</b></p>
 <p><b>Step 1a:</b> <b>Pull off cap</b></p>  <p><b>Step 1b:</b> <b>Open inhaler</b></p>	 <p><b>Step 2a:</b> <b>Pierce capsule once</b> Hold the inhaler upright. Pierce capsule by firmly pressing both side buttons at the same time.</p> <p>You should hear a noise as the capsule is pierced. <u>Only pierce the capsule once.</u></p>  <p><b>Step 2b:</b> <b>Release side buttons</b></p>	<p><b>Step 3a:</b> <b>Breathe out fully</b> <u>Do not blow into the inhaler.</u></p>  <p><b>Step 3b:</b> <b>Inhale medicine deeply</b> Hold the inhaler as shown in the picture. Place the mouthpiece in your mouth and close your lips firmly around it. <u>Do not press the side buttons.</u></p>	 <p><b>Check capsule is empty</b> Open the inhaler to see if any powder is left in the capsule.</p> <p>If there is powder left in the capsule:</p> <ul style="list-style-type: none"> <li>• Close the inhaler.</li> <li>• Repeat steps 3a to 3d.</li> </ul> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p><b>Powder remaining</b></p> </div> <div style="text-align: center;">  <p><b>Empty</b></p> </div> </div>



Step 1c:

**Remove capsule**

Separate one of the blisters from the blister card.

Peel open the blister and remove the capsule.

Do not push the capsule through the foil.

Do not swallow the capsule.

Breathe in quickly and as deeply as you can.  
During inhalation you will hear a whirring noise.  
You may taste the medicine as you inhale.



Step 3c:

**Hold breath**

Hold your breath for up to 5 seconds.

Step 3d:

**Rinse mouth**

Rinse your mouth with water after each dose and spit it out.



**Remove empty capsule**

Put the empty capsule in your household waste.  
Close the inhaler and replace the cap.



Step 1d:

**Insert capsule**

Never place a capsule directly into the mouthpiece.



Step 1e:


**Close inhaler**

**Important Information**

- Aectura Breezhaler capsules must always be stored in the blister card and only removed immediately before use.
- Do not push the capsule through the foil to remove it from the blister.
- Do not swallow the capsule.
- Do not use the Aectura Breezhaler capsules with any other inhaler.
- Do not use the Aectura Breezhaler inhaler to take any other capsule medicine.
- Never place the capsule into your mouth or the mouthpiece of the inhaler.
- Do not press the side buttons more than once.
- Do not blow into the mouthpiece.
- Do not press the side buttons while inhaling through the mouthpiece.
- Do not handle capsules with wet hands.
- Never wash your inhaler with water.

Your Ateectura Breezhaler Inhaler pack contains:

- One Ateectura Breezhaler inhaler
- One or more blister cards, each containing 10 Ateectura Breezhaler capsules to be used in the inhaler



## Frequently Asked Questions

### Why didn't the inhaler make a noise when I inhaled?

The capsule may be stuck in the capsule chamber. If this happens, carefully loosen the capsule by tapping the base of the inhaler. Inhale the medicine again by repeating steps 3a to 3d.

### What should I do if there is powder left inside the capsule?

You have not received enough of your medicine. Close the inhaler and repeat steps 3a to 3d.

### I coughed after inhaling – does this matter?

This may happen. As long as the capsule is empty you have received enough of your medicine.

### I felt small pieces of the capsule on my tongue – does this matter?

This can happen. It is not harmful. The chances of the capsule breaking into small pieces will be increased if the capsule is pierced more than once.

## Cleaning the inhaler

Wipe the mouthpiece inside and outside with a clean, dry, lint-free cloth to remove any powder residue. Keep the inhaler dry. Never wash your inhaler with water.

## Disposing of the inhaler after use

Each inhaler should be disposed of after all capsules have been used. Ask your pharmacist how to dispose of medicines and inhalers that are no longer required.