

Mind & Physiology Body Building:

Biometric-Guided Optimization of Body and Cognitive-Autonomic Function

A Scoping Review with N-of-1 Pilot Study and Preliminary Cohort Data

Nicolas L. Reynolds^{1,2}

¹*Mind Protocol, Paris, France*

²*Independent Researcher*

nlr@mindprotocol.ai

March 11, 2026

Highlights

- Proposes **Awareness & Bodybuilding** (A&B)—an integrative framework that combines physical fitness and cognitive-autonomic optimization through biometric-guided protocols
- Scoping review of 31 studies validates each component: hyperbolic tapering (PET data), HRV monitoring, pharmacogenomics, slow taper + support as most effective strategy identified (76 RCTs, $N=17,379$)
- The “Maximally Strong $\times$ Smart” hypothesis: physical optimization and cognitive-autonomic enhancement share overlapping physiological substrates, measurable through common biometric signals
- N-of-1 pilot study with 236 timestamped events (20 unique substances) over 7 days, concurrent wearable biometrics from 3 participants, DSM-5 SUD criteria mapping, and pharmacokinetic duration modeling
- Three-participant comparison (22 days each): polymedicated subject shows significantly elevated stress vs. both controls (Cohen’s $d = 1.40$, $p < 0.001$) despite indistinguishable resting heart rates ($p > 0.50$), demonstrating HRV-derived stress–HR decoupling
- Novel platform capabilities: real-time voice interface with biometric context injection, dyadic co-regulation monitoring (Mind Duo), multi-user Telegram-based enrollment (4+ users)
- Critical gap: no RCT of integrated biometric-guided platform; pragmatic trial design proposed to validate the discipline

Abstract

Background: Modern approaches to health optimization treat body and mind as separate domains—physical fitness is managed through exercise science, while cognitive and emotional health are delegated to psychiatry. Despite growing recognition of mind–body integration in fields such as lifestyle psychiatry and behavioral medicine, clinical practice often still separates these domains. The same physiological signals (HRV, stress, sleep architecture, autonomic balance) underpin both physical performance and cognitive-autonomic function. We describe *Awareness & Bodybuilding* (A&B)—a proposed integrative framework that uses continuous biometric monitoring to combine physical fitness optimization and

cognitive-autonomic enhancement through coordinated substance tapering, intentional movement, and cognitive practices. **Objective:** To evaluate the scientific evidence supporting each pillar of the A&B discipline as implemented by Mind Protocol: (1) biometric-guided substance tapering via real-time wearable monitoring, pharmacokinetic modeling, and AI-driven schedule adjustment; (2) physical optimization through movement practices with biometric feedback; and (3) consciousness enhancement through intentional use frameworks and cognitive tools. **Methods:** We conducted a scoping review searching PubMed, Google Scholar, Cochrane Library, and ClinicalTrials.gov (inception–January 2026), following PRISMA-ScR [Page et al., 2021], covering tapering pharmacology, wearable biosensors, pharmacogenomics, exercise neuroscience, and digital phenotyping. **Results:** The evidence supports each core component: (1) hyperbolic tapering is PET-validated [Horowitz and Taylor, 2019]; (2) HRV is a robust biomarker for autonomic disturbance [Alvares et al., 2016]; (3) wearable withdrawal detection achieves AUC 0.9997 in a single study ($N=16$, opioid-specific, research-grade device) [Carrasco-Hernandez et al., 2021]; (4) pharmacogenomics reduces ADRs by 30% ($N=6,944$) [Swen et al., 2023]; (5) slow tapering with support is optimal (76 RCTs, $N=17,379$) [Maund et al., 2025]; (6) exercise may produce effects of similar magnitude to antidepressants in mild-to-moderate depression [Singh et al., 2023]; (7) mindfulness-based cognitive therapy has shown non-inferiority to maintenance antidepressants for relapse prevention in recurrent depression [Kuyken et al., 2015b]. The N-of-1 pilot study captured 236 timestamped substance events (20 unique substances) over 7 days with concurrent wearable biometrics from 3 participants. DSM-5 criteria mapping identified severe nicotine use disorder (7/11 criteria), mild cannabis use disorder (3/11), and dose escalation in ketamine (8.7-fold over 4 days). A 3-participant comparison over 22 days revealed that the primary subject's mean stress (45.6/100) was significantly elevated relative to both controls (27.0 and 23.8/100; Mann-Whitney U , $p < 0.001$, Cohen's $d = 1.40$), despite indistinguishable resting heart rates (~ 55 – 58 bpm, $p > 0.50$). A predictive substance tracking engine using six weighted signals was implemented and validated with 30 automated tests. **Conclusion:** Each component of the A&B framework rests on scientific principles supported by moderate to strong evidence, though no component has been validated in its specific implementation within this platform. The principal gap is the absence of a prospective RCT testing the integrated approach. The platform implements capabilities with no direct precedent in the reviewed literature: a real-time voice interface with concurrent biometric context injection, dyadic co-regulation monitoring across paired consumer wearables, predictive substance logging with confirmation-based compliance enhancement, and a multi-user Telegram interface currently serving 4+ enrolled participants. The case study (236 logged events, 20 substances, concurrent biometrics from 3 participants) illustrates the framework in practice but does not constitute validation.

Keywords: awareness and bodybuilding, cognitive-autonomic optimization, psychotropic tapering, biometric monitoring, heart rate variability, wearable biosensors, precision psychiatry, integrative health, physical fitness, substance monitoring, co-regulation, voice interface, digital phenotyping

Contents

1	Introduction	6
1.1	The Fragmentation of Human Optimization	6
1.2	Awareness & Bodybuilding: A Proposed Framework	6
1.3	Scope of the Problem	6
1.4	Why Current Approaches Fail	7
1.5	The Mind Protocol Implementation	7
2	Methods	7
2.1	Search Strategy	7
2.2	Inclusion and Exclusion Criteria	7
2.3	Study Selection and Data Extraction	8
2.4	Codebase Analysis	8

70	3 Pharmacological Basis: Hyperbolic Tapering	8
71	3.1 SSRI Tapering	8
72	3.2 Antipsychotic Tapering	9
73	3.3 Clinical Validation	9
74	4 Meta-Analyses on Discontinuation	9
75	4.1 Incidence of Discontinuation Symptoms	9
76	4.2 Optimal Deprescribing Strategies	9
77	4.3 Earlier Systematic Reviews	10
78	5 HRV as a Biomarker for Withdrawal Monitoring	10
79	5.1 Direct Withdrawal Evidence	10
80	5.2 Psychiatric Baseline Impairment	10
81	5.3 Predictive and Differential Value	10
82	6 Wearable Biosensors: Real-Time Detection	11
83	6.1 Multi-Sensor Withdrawal Detection: Proof of Feasibility	11
84	6.2 Broader Substance Monitoring	11
85	6.3 Machine Learning Approaches	11
86	7 Drug-Class-Specific Tapering Evidence	12
87	7.1 Antipsychotics	12
88	7.2 Benzodiazepines	12
89	7.3 Withdrawal Phenomenology	12
90	8 Pharmacogenomics and Precision Dosing	12
91	9 Digital Phenotyping and Supportive Interventions	13
92	9.1 Multimodal Monitoring	13
93	9.2 Mindfulness as Tapering Support	13
94	9.3 HRV as Both Marker and Target	13
95	9.4 Psychedelic-Assisted Therapy Monitoring	13
96	9.5 Smartphone-Based Relapse Prediction	13
97	10 Mind Protocol: Technical Architecture	13
98	10.1 System Overview	14
99	10.2 Dependency Scoring (0–100)	14
100	10.3 Drug Interaction Detection	15
101	10.4 Pharmacokinetic Duration Model	15
102	10.5 Risk-Stratified Tapering Protocols	16
103	10.6 Biometric Validation Thresholds	16
104	10.7 Surveillance Symptom Tracking	17
105	10.8 Autonomic State Estimation Pipeline	17
106	10.9 Voice Interface: Real-Time Biometric-Contextualized Dialogue	17
107	10.10 Multi-User Platform and Telegram Interface	17
108	10.11 Predictive Substance Tracking	18
109	10.12 Dyadic Biometric Monitoring: Mind Duo	19
110	10.13 Privacy-by-Design Framework	19
111	11 Critical Synthesis: Validation Assessment	20

112	12 The Awareness & Bodybuilding Framework	20
113	12.1 From Tapering to Total Optimization	20
114	12.2 The “Maximally Strong × Smart” Thesis	21
115	12.3 Integrated Cognitive-Motor Tasks as a Measurement Model	21
116	12.4 The Tool-Based Model	21
117	12.5 The Intentional Use Framework	21
118	12.6 Substances vs. Activities: A Critical Distinction	22
119	12.7 Dependency Detection via Biometric Drift	23
120	12.8 Clinical Implications of A&B	23
121	12.9 Evidence Limitations of the Zero-Dependency Model	23
122	13 Gaps, Future Directions, and the A&B Research Agenda	24
123	13.1 Critical Evidence Gaps	24
124	13.2 Proposed Clinical Trial Design	24
125	13.3 Regulatory Considerations	25
126	14 Limitations	25
127	15 Conclusion	26
128	A Case Study: Awareness & Bodybuilding in Practice	34
129	A.1 Patient Profile	35
130	A.2 Cognitive Assessment: WAIS-IV	35
131	A.3 Clinical Observation	35
132	A.4 Current Pharmacotherapy	36
133	A.5 Substance Toolkit: Logged, Dosed, and Tracked	36
134	A.5.1 Ketamine (Intranasal)	36
135	A.5.2 LSD (Oral, Carton)	37
136	A.5.3 THC (Vaporized)	38
137	A.5.4 CBD (Sublingual, Tablet, Vaporized)	38
138	A.5.5 Nicotine (Vaporized Salt)	38
139	A.6 Biometric-Substance Temporal Overview	38
140	A.7 Co-Interaction Risk Assessment	39
141	A.8 Pharmacological Analysis: The Night of February 22–23	40
142	A.9 Temporal Pattern Analysis	40
143	A.10 Body Pillar: Movement and Physical Practices	40
144	A.10.1 Running	40
145	A.10.2 Dance as Integrated Cognitive-Motor Assessment	41
146	A.10.3 Yin-Yang Yoga	43
147	A.10.4 Breathwork and Meditation	43
148	A.11 Mind Pillar: Cognitive Enhancement Stack	43
149	A.12 Data Visualizations	43
150	A.12.1 Substance Use Timeline	43
151	A.12.2 Stress Trajectory with Substance Events	44
152	A.12.3 Dependency Score: Worked Example (Ketamine)	44
153	A.13 Biometric Baseline (February 2026)	45
154	A.14 Extended Biometric Baseline (22 Days)	45
155	A.15 Three-Participant Biometric Comparison	46
156	A.15.1 Descriptive Statistics	46
157	A.15.2 Between-Participant Comparisons	46
158	A.15.3 Within-Participant Correlations	47
159	A.15.4 Interpretation	47

160	A.16 Consolidated Pilot Study Findings	47
161	A.17 Tapering Roadmap	48
162	A.18 Proposed Outcome Metrics	48
163	B Technical Appendix: Mind Protocol Implementation Parameters	49
164	B.1 Dependency Scoring Algorithm	49
165	B.2 Pharmacokinetic Duration Model	49
166	B.3 Drug Interaction Matrix	50
167	B.4 Tapering Protocol Specifications	50
168	B.5 Biometric Validation Thresholds	50
169	B.6 Surveillance Symptom Categories	51
170	B.7 Pseudo-Code: Taper Step Decision	51
171	C Venice Convention: Design Heuristic Framework	52
172	C.1 Dual-Layer Architecture Rationale	52
173	C.2 Epistemic Scope Constraints	52
174	C.3 Art. 9 Special Category Firewall	52

1 Introduction

1.1 The Fragmentation of Human Optimization

Despite growing recognition of mind–body integration in fields such as lifestyle psychiatry, behavioral medicine, and psychoneuroimmunology, clinical practice often still treats physical fitness and mental health as separate systems. Physical fitness is managed through exercise science, nutrition, and sports medicine. Mental health is delegated to psychiatry, psychopharmacology, and psychotherapy. Cognitive-autonomic function—the capacity for sustained attention, emotional regulation, and intentional agency—is rarely integrated into physical fitness programming.

Yet the physiological signals underlying both domains overlap substantially. Heart rate variability (HRV), stress reactivity, sleep architecture, autonomic balance, and body battery are not exclusively “mental” or “physical” metrics—they are *integrated* signals of the organism’s state. A runner with poor HRV recovery is likely both physically deconditioned and may exhibit reduced cognitive performance [Laborde et al., 2017]. A patient tapering from benzodiazepines experiences both somatic withdrawal symptoms and cognitive-perceptual disturbances [Cosci and Chouinard, 2020b]. The data does not respect the mind–body divide; our disciplines should not either.

1.2 Awareness & Bodybuilding: A Proposed Framework

We describe **Awareness & Bodybuilding** (A&B)—a proposed integrative framework that uses continuous biometric monitoring to combine physical fitness optimization and cognitive-autonomic enhancement. The central hypothesis:

We hypothesize that physical optimization and cognitive-autonomic enhancement are synergistic, sharing overlapping physiological substrates. Both may be measurable through common biometric signals, and improvements in one domain may facilitate improvements in the other.

A&B integrates three pillars:

1. **Body:** Physical movement (running, dance, yoga), nutritional optimization, and substance tapering—reducing chemical dependencies that impair physiological resilience.
2. **Mind:** Cognitive practices (meditation, breathwork), supplements, and monitored substance use—each logged, dosed, and evaluated against biometric outcomes.
3. **Measurement:** Continuous wearable biometrics (HRV, stress, sleep, Body Battery) as the unified feedback loop for both pillars. The same data that tracks tapering safety also tracks fitness progression and cognitive-autonomic function.

The biometric monitoring layer is what distinguishes A&B from purely philosophical frameworks: every substance, every workout, every practice session generates physiological data that can be tracked and correlated with self-reported outcomes, enabling data-driven adjustment of protocols.

1.3 Scope of the Problem

The need for such integration is acute. Antidepressant consumption has risen steadily across OECD countries, with some nations exceeding 100 defined daily doses per 1,000 population per day [Organisation for Economic Co-operation and Development, 2023]. In England alone, 7.3 million adults were prescribed dependence-forming medications in 2017–2018 [Taylor et al., 2019]. The Henssler et al. [2024] meta-analysis estimates that 15% of patients experience genuine discontinuation symptoms after correcting for nocebo effects, though earlier reviews placed this figure at 33–56% [Davies and Read, 2019]. Individuals prescribed long-term psychotropic medications often report reduced physical activity and cognitive complaints [Firth et al., 2019], though the directionality of these associations remains debated.

1.4 Why Current Approaches Fail

Clinical guidelines typically recommend “gradual tapering over weeks to months” without specifying how to individualize the rate. Gabriel and Sharma [2017] found that nine major guidelines recommend tapering periods ranging from 4 weeks to 6 months, but only two studies directly compared different tapering rates. Physicians typically have limited real-time visibility into a patient’s physiological state between consultations. And crucially, tapering is treated in isolation—divorced from the physical and cognitive practices that could support it.

1.5 The Mind Protocol Implementation

Mind Protocol is a software platform that attempts to operationalize the A&B framework; its technical architecture is described in Section 10 and evaluated against the evidence base. This paper evaluates whether existing scientific evidence supports each component of the integrated approach.

2 Methods

This review follows the PRISMA extension for scoping reviews (PRISMA-ScR) [Page et al., 2021]. We note that this is a *scoping review*—not a full systematic review—as we did not perform formal risk-of-bias assessment (e.g., RoB 2, ROBINS-I, AMSTAR 2), did not register a protocol on PROSPERO, and our scope intentionally spans heterogeneous study designs across multiple domains (pharmacology, wearable technology, pharmacogenomics, digital health). The scoping review framework is appropriate when the objective is to map available evidence across a broad topic and identify gaps, rather than to synthesize effect sizes for a specific clinical question.

2.1 Search Strategy

We searched PubMed, Google Scholar, Cochrane Library, and ClinicalTrials.gov from database inception through January 2026. Search terms included:

- “antidepressant tapering” OR “medication discontinuation syndrome”
- “HRV withdrawal biomarker” OR “heart rate variability withdrawal”
- “wearable biosensor withdrawal” OR “wearable substance monitoring”
- “pharmacogenomics antidepressant dosing” OR “CYP2D6 tapering”
- “hyperbolic tapering” OR “receptor occupancy tapering”
- “digital phenotyping psychiatry” OR “smartphone relapse prediction”
- “benzodiazepine withdrawal protocol” OR “Ashton Manual”
- “antipsychotic dose reduction” OR “antipsychotic discontinuation”
- “biometric-guided pharmacotherapy”

2.2 Inclusion and Exclusion Criteria

Inclusion: (1) Randomized controlled trials, systematic reviews, meta-analyses, and pharmacological analyses studying tapering or discontinuation of psychotropic medications; (2) studies validating HRV or other wearable-derived biomarkers for monitoring psychiatric conditions or withdrawal; (3) pharmacogenomic guidelines and implementation studies for psychotropic dosing; (4) digital phenotyping and continuous monitoring studies in psychiatric populations.

Exclusion: (1) Case reports and case series with $N < 10$; (2) animal studies without clinical translation; (3) studies of non-psychotropic medications (e.g., cardiovascular, endocrine); (4) studies not available in English or French.

2.3 Study Selection and Data Extraction

Titles and abstracts were screened for relevance. Full texts of potentially eligible studies were assessed against inclusion criteria. Data extracted included: study design, sample size, population characteristics, intervention details, primary outcomes, and relevance to biometric-guided tapering.

Limitation: Study screening and data extraction were performed by the authors (MIND and N.L.R.) without independent duplicate screening by external researchers. A fully PRISMA-compliant review would require independent screening by at least two researchers with inter-rater reliability assessment (e.g., Cohen’s κ). This limitation should be considered when interpreting the scope and completeness of the included studies. The search and screening process is summarized in Figure 1.




Figure 1: PRISMA flow diagram for study selection.

2.4 Codebase Analysis

In addition to the literature review, we analyzed Mind Protocol’s source code (Python, TypeScript) to map its technical architecture to the evidence base. Components assessed included: dependency scoring algorithms, substance interaction matrices, pharmacokinetic duration models, tapering protocol definitions, and biometric validation thresholds.

3 Pharmacological Basis: Hyperbolic Tapering

A key pharmacological principle underlying personalized tapering is that the relationship between drug dose and receptor occupancy follows a hyperbolic—not linear—curve. Equal dose reductions at lower doses produce disproportionately larger drops in receptor occupancy, explaining why withdrawal symptoms often intensify during the final stages of tapering.

3.1 SSRI Tapering

Horowitz and Taylor [2019] used PET imaging data of serotonin transporter (SERT) occupancy to demonstrate that the dose–occupancy relationship for SSRIs follows a hyperbolic curve. They

proposed that doses should be reduced by fixed amounts of SERT occupancy (e.g., 10% increments), translating to progressively smaller dose reductions—a “hyperbolic” taper. A personalized rate can be established by an initial trial reduction equivalent to 10% SERT occupancy, followed by monitoring of withdrawal symptom severity and duration.

3.2 Antipsychotic Tapering

Horowitz et al. [2021] extended the hyperbolic tapering principle to antipsychotics via D2 receptor occupancy data. Dopaminergic hypersensitivity following long-term use means abrupt discontinuation or linear tapering risks relapse. The recommended approach: reduce by one-quarter to one-half of the most recent dose at intervals of 3–6 months, titrated to individual tolerance.

3.3 Clinical Validation

Groot and van Os [2021] validated the approach clinically through “tapering strips”—28-day rolls of daily pouches with progressively lower doses enabling hyperbolic dose reduction—in 1,240 antidepressant users. 72% of patients successfully discontinued, with a median taper duration of 56 days. However, as a single-arm cohort study without a control group, the 72% success rate cannot be causally attributed to the hyperbolic tapering approach alone. van Os and Groot [2023] reported sustained discontinuation at 1–5 year follow-up.

Table 1: Hyperbolic tapering evidence summary.

Study	Year	Design	N	Key Finding
Horowitz & Taylor	2019	PET analysis	—	SSRI dose–SERT occupancy is hyperbolic
Horowitz et al.	2021	PET analysis	—	Extended to antipsychotics (D2 occupancy)
Groot & van Os	2021	Cohort	1,240	72% success with tapering strips
van Os & Groot	2023	Follow-up	Cohort	Sustained at 1–5 years

4 Meta-Analyses on Discontinuation

4.1 Incidence of Discontinuation Symptoms

Henssler et al. [2024] conducted the most comprehensive meta-analysis to date (>50 studies, >17,000 patients). After correcting for placebo discontinuation effects (nocebo), approximately 15% of patients (1 in 6–7) experience genuine antidepressant discontinuation symptoms, with only ~3% experiencing severe symptoms. Critically, studies using structured assessment tools found significantly higher incidence (40%) than those without (27%), suggesting systematic under-detection.

4.2 Optimal Deprescribing Strategies

The 2025 network meta-analysis [Maund et al., 2025]—76 RCTs, $N=17,379$ —is the largest analysis of deprescribing strategies to date. Slow tapering (>4 weeks) combined with psychological support was the most effective strategy, performing comparably to antidepressant continuation for relapse prevention and clearly outperforming abrupt discontinuation or fast tapering. Psychological support enhanced tapering success but showed no benefit when added to abrupt discontinuation, emphasizing its value specifically during structured withdrawal.

4.3 Earlier Systematic Reviews

Davies and Read [2019] estimated withdrawal incidence at 33–56%, with severe symptoms in ~25% (later critiqued for selection bias). Gabriel and Sharma [2017] found nine clinical guidelines recommending tapering periods of 4 weeks to 6 months, but only two studies directly compared rates—underscoring the scarcity of comparative evidence.

Table 2: Meta-analytic evidence on discontinuation.

Study	Design	N	Key Finding
Henssler et al., 2024	SR + MA	>17,000	15% genuine withdrawal; under-detection with unstructured tools
Lancet 2025	Network MA	17,379	Slow taper + support = most effective strategy
Davies & Read, 2019	SR	24 studies	33–56% incidence; guidelines underestimate
Gabriel & McMahon, 2019	SR	Multiple	Only 2 studies compare tapering rates

5 HRV as a Biomarker for Withdrawal Monitoring

Heart rate variability (HRV)—the variation in time between successive heartbeats—reflects the balance between sympathetic and parasympathetic nervous system activity. It is one of the most studied biomarkers in psychiatry and increasingly recognized as a real-time indicator of autonomic disturbance during withdrawal.

5.1 Direct Withdrawal Evidence

Makhoul et al. [2020] demonstrated significant decreases in high-frequency (HF) HRV and RMSSD during naloxone-induced opioid withdrawal ($N=10$). Heart rate and systolic blood pressure increased. The findings revealed that withdrawal involves *both* sympathetic hyperactivation *and* parasympathetic suppression—a dual autonomic disturbance that current medications (clonidine, lofexidine) address only partially.

5.2 Psychiatric Baseline Impairment

Alvares et al. [2016] conducted a meta-analysis of 140 case-control studies ($k=151$) finding that HRV was reduced across all psychiatric patient groups compared to controls (Hedges’ $g = -0.583$), with the largest effect in psychotic disorders ($g = -0.948$). Critically, reductions persisted in medication-free individuals, indicating that the condition itself impairs autonomic function. This finding suggests that personal baselines may be more informative than population norms for tapering decisions, though this hypothesis has not been directly tested.

5.3 Predictive and Differential Value

Kircanski et al. [2019] showed that pre-treatment HRV predicted antidepressant outcomes differentially by anxiety subtype (higher HRV predicted better outcomes in anxious depression; the reverse in nonanxious depression). Chalmers et al. [2014] confirmed via meta-analysis ($N=4,380$) that anxiety disorders—the most common candidates for SSRI and benzodiazepine tapering—are associated with reduced HRV, meaning these patients have less autonomic “buffer” to absorb withdrawal stress.

Table 3: HRV evidence for withdrawal and psychiatric monitoring.

Study	Design	N	Key Finding
Makhoul et al., 2020	Experimental	10	HF-HRV and RMSSD decrease during opioid withdrawal
Alvares et al., 2016	MA (140 studies)	$k=151$	HRV reduced in all psychiatric groups ($g = -0.58$)
Kircanski et al., 2019	RCT (secondary)	Large	HRV predicts antidepressant response by subtype
Chalmers et al., 2014	MA	4,380	Anxiety = reduced HRV = less autonomic buffer

6 Wearable Biosensors: Real-Time Detection

6.1 Multi-Sensor Withdrawal Detection: Proof of Feasibility

Carrasco-Hernandez et al. [2021] achieved $AUC = 0.9997$ for detecting opioid withdrawal using an Empatica E4 wrist-mounted biosensor (blood volume pulse, electrodermal activity, skin temperature, accelerometry) and a Random Forest classifier ($N=16$). Detection was possible with just one minute of biosensor data, with electrodermal activity and skin temperature—not HRV alone—likely dominating the feature importance.

Important caveats: This result represents a *proof of feasibility*, not a validated clinical tool, for several reasons: (1) the small sample ($N=16$) creates risk of overfitting; (2) the study used a research-grade multi-sensor device (Empatica E4), not consumer wearables (Garmin, Apple Watch); (3) the context was acute opioid withdrawal in an ED setting, which differs substantially from gradual SSRI or benzodiazepine tapering; and (4) the high AUC reflects multi-modal sensing (EDA, temperature, BVP, accelerometry), not HRV alone. Generalization to consumer devices and non-opioid drug classes remains unvalidated.

6.2 Broader Substance Monitoring

Natarajan et al. [2016] validated wearable biosensors for detecting opioid-related physiological changes ($F1=0.80$, $AUC=0.77$). Carreiro et al. [2020] demonstrated that wearable sensors can differentiate episodes of craving versus stress in substance use disorder patients—a distinction essential for platforms needing to separate normal stress from withdrawal-induced disturbance.

6.3 Machine Learning Approaches

Hoffman et al. [2020] showed that ML tools can predict alcohol withdrawal severity at admission using clinical and physiological features, with Random Forest outperforming logistic regression. Extending to continuous wearable data has been proposed but not yet validated.

Table 4: Wearable biosensor evidence for substance monitoring.

Study	Design	N	Key Finding
Carrasco et al., 2021	ML validation	16	AUC 0.9997 with 1 min of wrist sensor data
Natarajan et al., 2016	Pilot	Multiple	Wearables detect opioid-related changes (AUC 0.77)
Carreiro et al., 2020	Mixed-methods	SUD pts	Sensors differentiate craving vs. stress
Hoffman et al., 2020	Retrospective ML	Multiple	ML predicts alcohol withdrawal severity

7 Drug-Class-Specific Tapering Evidence

7.1 Antipsychotics

The RADAR trial [Moncrieff et al., 2023]—an RCT of 253 patients across 19 NHS Trusts—found that gradual antipsychotic reduction approximately doubled the relapse rate (25% vs. 13%), though 75% of the reduction group did *not* experience serious relapse, and long-term functional outcomes favored the reduction arm. Wunderink et al. [2013] showed in a 7-year follow-up ($N=131$) that while short-term relapse risk was higher, long-term social functioning was *higher* in the dose-reduction group, suggesting that the acute tapering phase (first 1–2 years) is the critical monitoring window.

7.2 Benzodiazepines

Morin et al. [2004] demonstrated that individualized tapering combined with CBT achieved 90% dose reduction and 63% complete discontinuation in 76 older adults, with no significant adverse events. Ashton [2002] established the most widely referenced clinical protocol for benzodiazepine tapering, though it has not been evaluated in a randomized controlled trial: switch to long-acting formulation, reduce by $\leq 10\%$ every 1–2 weeks, with the patient controlling pace. Ashton explicitly identified the variables requiring individualization (autonomic stability, sleep quality, stress tolerance, recovery speed)—several of which (e.g., heart rate variability, sleep duration) are partially measurable with modern consumer wearables, though with reduced accuracy compared to clinical instruments.

7.3 Withdrawal Phenomenology

Cosci and Chouinard [2020a] identified three distinct withdrawal phenomena: (1) acute withdrawal (resolves in 2–4 weeks); (2) rebound (original symptoms at greater intensity); and (3) persistent post-withdrawal disorder (PPWD), lasting months to years. The distinction has direct implications for monitoring algorithms, as different physiological signatures likely correspond to each phase.

8 Pharmacogenomics and Precision Dosing

Pharmacogenomics—the study of how genetic variation affects drug response—is increasingly recognized as essential for personalizing both prescribing and deprescribing.

Bousman et al. [2021] published updated CPIC guidelines providing genotype-based dosing recommendations for SSRIs and SNRIs based on CYP2D6, CYP2C19, and CYP2B6 metabolizer status. Dosage adjustment was recommended for 52% of escitalopram patients and 54% of TCA

patients. A CYP2D6 poor metabolizer on paroxetine will have higher plasma levels at any given dose, requiring slower taper schedules. These guidelines were developed for initial prescribing decisions; their application to tapering schedule individualization has not been directly studied. Swen et al. [2023] demonstrated in the PREPARE study—the first large-scale prospective pharmacogenomic implementation study ($N=6,944$ across 7 European countries)—that a 12-gene panel reduced the odds of clinically relevant adverse drug reactions by 30%. We propose that combining pharmacogenomic data with continuous biometric monitoring could create a dual-layer personalization approach, though this integration has not been tested: genetic profile would inform the starting taper schedule, and real-time physiological data could adjust it dynamically.

9 Digital Phenotyping and Supportive Interventions

9.1 Multimodal Monitoring

Hein et al. [2025] demonstrated that multimodal digital phenotyping using smartphones and wearables can predict relapse and symptom exacerbations in major depression. Composite digital phenotypes from multiple data streams proved superior to single-stream analysis—providing a methodological template for biometric tapering platforms.

9.2 Mindfulness as Tapering Support

The PREVENT trial ($N=424$) demonstrated that Mindfulness-Based Cognitive Therapy (MBCT) is non-inferior to maintenance antidepressants for relapse prevention [Kuyken et al., 2015a]. Kuyken et al. [2016] confirmed via individual patient data meta-analysis ($N=1,258$) that MBCT reduced relapse by 31%. This supports the network meta-analysis finding [Maund et al., 2025] that psychological support enhances tapering success.

9.3 HRV as Both Marker and Target

Garland et al. [2014] showed that Mindfulness-Oriented Recovery Enhancement (MORE) increased HRV and reduced opioid craving via enhanced autonomic regulation. This suggests HRV biofeedback could serve as an *active intervention* component alongside tapering, not just passive monitoring.

9.4 Psychedelic-Assisted Therapy Monitoring

Mitchell et al. [2023] demonstrated 67% PTSD remission with MDMA-assisted therapy in a Phase 3 trial, though the FDA subsequently declined to approve MDMA-assisted therapy in 2024, citing methodological concerns including functional unblinding. Bonnelle et al. [2024] showed that pre-treatment HRV predicts psychedelic therapy outcomes. As psychedelic medicine evolves, biometric monitoring during substance-assisted sessions becomes increasingly relevant.

9.5 Smartphone-Based Relapse Prediction

Wang et al. [2016] demonstrated that smartphone-derived behavioral features (mobility, social interaction, phone usage) predict psychiatric relapse with clinically useful accuracy in schizophrenia—establishing a digital phenotyping framework applicable to tapering monitoring.

10 Mind Protocol: Technical Architecture

Mind Protocol’s codebase implements a multi-layered tapering support system. The architecture below is presented as the *data collection and monitoring apparatus*—the methodological

infrastructure that enables the observations reported in Appendix A. Its clinical effectiveness is not claimed and remains untested.

Analysis of the source code reveals the following components. **Important:** the components described below exist in the codebase (implemented in Python/TypeScript) but are at different stages of deployment. The substance logging, biometric collection, drug interaction detection, and dependency scoring modules are operational. The automated tapering schedule generation, biometric-triggered holds, and adaptive dose adjustment are implemented in code but have not yet been deployed in a clinical or self-directed tapering context.

10.1 System Overview




Figure 2: Mind Protocol system architecture. Orange boxes indicate patient-facing interfaces; blue boxes indicate processing layers. Arrows show data flow.

10.2 Dependency Scoring (0–100)

A composite dependency score is calculated from five weighted components: substance class risk factor, duration of use, dosage level, pharmacokinetic half-life, and individual patient factors. This draws loosely on principles from the Ashton Manual’s multi-variable approach [Ashton, 2002], Horowitz’s receptor-occupancy framework [Horowitz and Taylor, 2019], and DSM-5 substance use disorder criteria [American Psychiatric Association, 2013]. The component structure parallels validated instruments such as the Severity of Dependence Scale [Gossop et al., 1995] and the Fagerström Test for Nicotine Dependence [Heatherton et al., 1991], although the specific weighting scheme is author-derived and has not been independently validated.

To illustrate how the platform’s dependency constructs map to established clinical criteria, Table 5 applies the DSM-5 SUD diagnostic framework [American Psychiatric Association, 2013] to three substances actively tracked in the case study. Criteria were assessed by the author based on self-reported behavioral patterns during the 6-day observation period; no structured clinical interview (e.g., SCID-5) was administered.

Table 5: DSM-5 Substance Use Disorder criteria applied to three tracked substances. ✓ = criterion met during observation period; – = not met or insufficient evidence. SUD severity: mild (2–3 criteria), moderate (4–5), severe (≥ 6).

DSM-5 Criterion	Nicotine	THC	Caffeine [†]
<i>Impaired control</i>			
1. Larger amounts / longer than intended	✓	✓	–
2. Persistent desire / unsuccessful efforts to cut down	✓	–	–
3. Great deal of time obtaining / using / recovering	✓	–	–
4. Craving or strong desire to use	✓	–	–
<i>Social impairment</i>			
5. Failure to fulfill major role obligations	–	–	–
6. Continued use despite social / interpersonal problems	–	–	–
7. Important activities given up or reduced	–	–	–
<i>Risky use</i>			
8. Recurrent use in hazardous situations	–	–	–
9. Continued despite knowledge of physical / psychological problem	✓	✓	–
<i>Pharmacological</i>			
10. Tolerance (need more for same effect)	✓	✓	✓
11. Withdrawal (symptoms on cessation)	✓	–	✓
Criteria met	7/11	3/11	2
Severity	Severe	Mild	N/A

[†] Caffeine Use Disorder is included in DSM-5 Section III as a “Condition for Further Study” and is not an official diagnosis. Tolerance and withdrawal are recognized in ICD-10 (F15.2).

The DSM-5 mapping reveals a gradient: nicotine meets criteria for severe SUD (7/11), consistent with the 63 logged sessions and 4 “craving” entries observed during the case study period. THC meets mild SUD criteria (3/11), with dose escalation (Feb 24: 6 chambers, Feb 26: 9 chambers) but no craving entries. Caffeine shows pharmacological criteria only (tolerance, withdrawal) but is not classifiable as a DSM-5 SUD. The platform’s dependency scoring algorithm captures analogous signals (dose escalation $\leftrightarrow$ Criterion 1; intent degradation $\leftrightarrow$ Criterion 4; compulsive timing $\leftrightarrow$ Criterion 3) but applies continuous scoring rather than binary criteria.

10.3 Drug Interaction Detection

A SUBSTANCE_INTERACTIONS matrix encodes 40+ drug–drug interactions, each with a severity score. Interactions are evaluated dynamically as substances are added, modified, or tapered. A 16-rule interaction engine generates real-time alerts, addressing the pharmacovigilance gap where >90% of standard DDI alerts are overridden due to alert fatigue.

10.4 Pharmacokinetic Duration Model

A PK_DURATION_MIN model maps each substance to its pharmacokinetic duration, informing minimum intervals between dose reductions. This operationalizes the Horowitz principle that tapering rate must respect receptor re-equilibration time. Table 6 summarizes the pharmacokinetic parameters for each tracked substance, drawn from primary pharmacological sources.

Table 6: Pharmacokinetic parameters for tracked substances. $T_{\max}$: time to peak plasma concentration; $t_{1/2}$: elimination half-life; Duration: clinically relevant effect duration used by the platform’s minimum-interval model.

Substance	Route	$T_{\max}$ (h)	$t_{1/2}$ (h)	Duration (h)	Source
<i>Prescribed medications</i>					
Sertraline	Oral	4.5–8.4	26	24	DeVane et al. [2002]
Venlafaxine	Oral	2.4	5 (ODV: 11)	24	Hiemke et al. [2018]
Prazepam	Oral/SL	2–6	30–100 ^a	24–48	Greenblatt et al. [1981]
Cyamemazine	Oral	1–3	10	8–12	Hiemke et al. [2018]
<i>Psychoactive substances</i>					
Ketamine	Intranasal	0.3	2.5	1–2	Zanos et al. [2018]
LSD	Oral	1.5–2.5	3.6	8–12	Holze et al. [2021]
THC	Vaporized	0.05–0.17	1.3 (terminal: 25–36)	2–4	Huestis [2007]
CBD	Oral/SL	1–4	18–32	6–8	Millar et al. [2018]
Nicotine	Vaporized	0.05–0.17	2	1–2	Benowitz et al. [2009]
<i>Supplements</i>					
Caffeine	Oral	0.5–1	3–7	4–6	Rowland and Tozer [2011]
Melatonin	Oral	0.5–1	0.5–1	4–6	Tordjman et al. [2017]

^aActive metabolite (desmethyldiazepam). Parent compound $t_{1/2} \approx 1.3$ h.

10.5 Risk-Stratified Tapering Protocols

TAPERING_PROTOCOLS defines schedules stratified by risk class (Table 7). Each class specifies step size, interval, monitoring frequency, and escalation thresholds.

Table 7: Risk-stratified tapering protocol classes in Mind Protocol.

Risk	Examples	Evidence	Step / Interval
Critical	BZDs, barbiturates	Ashton Manual	$\leq 10\%$ / 1–2 wk
High	SSRIs, opioids, APs	Horowitz; RADAR	Hyperbolic / 2–4 wk
Moderate	Gabapentin, pregabalin	Clinical consensus	10–15% / 1–2 wk
Standard	Supplements, low-risk	Clinical judgment	20–25% / 1 wk

10.6 Biometric Validation Thresholds

A `BiometricValidation` module defines withdrawal detection thresholds based on deviations from personal baseline. Exact thresholds are specified per substance (see Appendix B for full parameters). For example, venlafaxine tapering triggers alerts at stress +15 above baseline, HRV –10 below baseline, and sleep score –15 below baseline; prazepam uses more conservative thresholds (stress +25, HRV –15, sleep –20) reflecting higher withdrawal severity.

Limitation: Garmin-derived metrics (Stress Score, Body Battery) rely on proprietary algorithms with limited published validation. These metrics should be considered *auxiliary signals* complementing—not replacing—clinical assessment. Inter-individual calibration variability and lack of regulatory validation mean that threshold breaches should trigger clinical consultation, not autonomous treatment decisions. A priority validation step is correlating Garmin-derived metrics against a research-grade reference (e.g., Empatica E4) in a subsample, with sensitivity/specificity analysis for withdrawal detection. This implements the principle from Alvares et al. [2016] that personal baselines, not population norms, should drive clinical decisions.

10.7 Surveillance Symptom Tracking

60+ withdrawal symptoms across 8 categories (neurological, gastrointestinal, cardiovascular, psychological, sensory, motor, cognitive, autonomic) are monitored via patient self-report and biometric correlation. This structured approach addresses the Henssler et al. [2024] finding that structured assessment detects 40% more withdrawal cases than unstructured methods.

10.8 Autonomic State Estimation Pipeline

Raw Garmin metrics are transformed into a composite autonomic state estimate (ANS_ESTIMATE) used by downstream modules. The pipeline computes:

- **ANS mode** (survival, balanced, recovery): derived from resting HR, overnight HRV mean, average stress, and current Body Battery via weighted thresholds.
- **Parasympathetic/sympathetic ratio** ($\in [0, 1]$): a continuous index of autonomic balance.
- **Fatigue reduction factor** (RED_FAT, $\in [0, 1.5]$): composite of sleep architecture score (deep% $\geq 13\%$, REM% $\geq 18\%$, duration $\geq 7h$), current Body Battery, and drain trajectory (heavy >30 pts/morning, moderate, stable, or charging).
- **Stress reduction factor** (RED_ANS, $\in [0, 1]$): current stress and HRV combined.

These derived signals feed into withdrawal threshold evaluation (Appendix B), alert generation, and the AI decision layer's risk stratification. The computation is deterministic and parameter-transparent, enabling clinician audit.

10.9 Voice Interface: Real-Time Biometric-Contextualized Dialogue

Mind Protocol implements a real-time voice interface via WebSocket: spoken audio (webm/opus) is streamed to a server-side pipeline consisting of (1) automatic speech recognition (OpenAI Whisper), (2) language model response generation (Claude, constrained to 2–4 sentences), and (3) text-to-speech synthesis (ElevenLabs streaming, MP3). The system prompt is dynamically augmented with the user's current biometric state (HR, stress, Body Battery, ANS mode) from the most recent Garmin reading, enabling context-aware responses (e.g., detecting elevated stress and adjusting tone accordingly).

Hallucination mitigation includes non-Latin script rejection, no_speech_prob filtering (>0.6), and repetition detection. A 10-turn conversation history is maintained per session.

A complementary *passive dialogue model* operates on the local device: a 60-second rolling audio buffer (20×3s chunks) records the user's speech in parallel with TTS playback. After each AI utterance, the buffer window corresponding to the TTS duration is transcribed via Whisper and merged chronologically with the AI's output, reconstructing a natural interleaved conversation without requiring explicit turn-taking. This design reflects the insight that human–AI dialogue need not follow strict turn-based protocols.

No existing tapering support platform described in the reviewed literature incorporates real-time voice interaction with concurrent biometric context injection.

10.10 Multi-User Platform and Telegram Interface

Although the case study (Appendix A) documents a single subject, the platform architecture supports multiple concurrent users. As of February 2026, 4+ individuals have enrolled via the primary patient-facing interface: a Telegram chatbot.

The enrollment pipeline includes: (1) automated registration with tiered trust levels (owner, high, medium, low, stranger) and per-tier daily interaction quotas; (2) Garmin device linking via time-limited OAuth codes (/link/garmin → Garmin Connect → tokens stored per user ID); and (3) optional integrations (Gmail, Spotify) for contextual behavioral data. Each user's

biometric data is stored in isolated per-user directories (`biometrics/{user_id}/`), and privacy constraints ensure that one user’s physiological data is never exposed to another.

The Telegram bridge handles text messages, voice notes (transcribed via Whisper and routed to the AI), and group chat interactions (@-mention or reply-based activation). Voice responses are generated as OGG/Opus Telegram voice notes when appropriate. A spam detection layer (emoji flood ratio >70%, repeated-substring ratio >50%) filters abusive input.

This multi-user architecture demonstrates platform readiness for multi-participant studies, though the current enrollment is self-selected and uncontrolled.

10.11 Predictive Substance Tracking

A persistent challenge in longitudinal substance monitoring is *logging compliance*: subjects who log 30+ substances and activities daily inevitably forget entries, creating gaps that degrade pattern detection and dependency scoring accuracy. The platform addresses this with a predictive confirmation system—colloquially, “Is this you?”—that inverts the logging model: instead of waiting for the subject to report intake, the system predicts what was likely taken and prompts for confirmation.

The prediction engine computes a confidence score $C \in [0, 1]$ for each candidate substance using a weighted sum of six independent signals:

$$C = \min\left(1, \underbrace{w_{\text{reg}}}_{\leq 0.40} + \underbrace{w_{\text{med}}}_{\leq 0.35} + \underbrace{w_{\text{hist}}}_{\leq 0.25} + \underbrace{w_{\text{tod}}}_{\leq 0.15} + \underbrace{w_{\text{dow}}}_{\leq 0.10} + \underbrace{w_{\text{bio}}}_{\leq 0.15}\right) \quad (1)$$

where w_{reg} is the regimen schedule signal (fixed-time prescriptions), w_{med} is the medical profile signal (daily vs. PRN classification from the prescribing record), w_{hist} is 30-day historical frequency (proportion of days with logged use), w_{tod} is time-of-day alignment (distance from the subject’s peak-hour histogram), w_{dow} captures day-of-week patterns (weekend vs. weekday use), and w_{bio} is the biometric context signal—current Garmin stress, body battery, and sleep score used to boost PRN predictions (e.g., prazepam predicted when stress exceeds 60/100).

Substances already logged today are filtered out. Only predictions with $C \geq 0.20$ are surfaced. Three substance categories emerge naturally:

- **Daily medications** (sertraline, venlafaxine, supplements): $C \sim 0.70$ – 0.90 , dominated by w_{med} and w_{hist} .
- **Habitual substances** (nicotine, caffeine, THC): $C \sim 0.40$ – 0.70 , driven by w_{hist} and w_{tod} .
- **PRN substances** (prazepam, cyamemazine, melatonin): $C \sim 0.20$ – 0.40 , baseline low but boosted by w_{bio} during high-stress or poor-sleep episodes.

The confirmation flow is delivered via Telegram inline keyboards. At 06:00 CET daily, the orchestrator computes predictions, groups them by time period (morning, afternoon, evening, night), and sends a structured prompt with three interaction modes: bulk confirmation (single tap logs all predicted substances), per-substance confirm/dismiss, or dose/time correction. Each confirmed prediction creates a traceable entry in the substance log with `source: "prediction"` and the originating prediction ID, enabling accuracy tracking.

An accuracy module logs outcomes (confirmed, corrected, dismissed, expired) per prediction and computes calibration curves—whether predictions at 80% confidence are actually confirmed ~80% of the time. This closed feedback loop enables iterative signal weight adjustment as usage data accumulates.

This system addresses the detect–act gap identified in Section 13: while the platform cannot administer medication, it can reduce the friction of self-reporting to a single confirmation tap—transforming substance logging from an active recall task to a passive verification task.

10.12 Dyadic Biometric Monitoring: Mind Duo

Mind Protocol extends single-user monitoring to interpersonal *co-regulation*, drawing on the established literature on physiological linkage in couples [Timmons et al., 2015]. The Mind Duo module reads biometric data from two paired Garmin devices (e.g., subject + partner) and computes:

- **Synchrony score** ($\in [0, 1]$): Pearson correlation plus mean absolute delta over a rolling 30-minute window on time-aligned stress timeseries.
- **Phase detection** across five states: *independent* (low correlation, high inter-individual delta), *co-activation* (both elevated, low delta), *co-regulation* (both declining synchronously), *divergent* (opposite trajectories), and *rebound* (rapid parasympathetic recovery in one or both).
- **Readiness level**: *not needed*, *available*, *suggested*, *active*, or *rebound*—indicating whether interpersonal regulation support is recommended.
- **Guidance generation**: when co-regulation is suggested, the system produces shared breathing or grounding prompts (delivered via TTS or Telegram).

In the case study context, the subject (Garmin Fēnix 8) is paired with a partner (Garmin Venu 2S/3S). Sessions are logged (`biometrics/duo/sessions.jsonl`) with before/after biometric snapshots and synchrony scores.

Novelty and evidence gap: Autonomic coupling between individuals (interpersonal physiological synchrony) has been documented in laboratory settings using research-grade equipment [Feldman, 2011], but no consumer-wearable co-regulation system has been described in the tapering or digital health literature. Mind Duo represents an exploratory implementation whose clinical utility remains entirely untested.

10.13 Privacy-by-Design Framework

The architecture embeds a privacy-by-design framework (internally termed the “Venice Convention”; see Appendix C for the full design heuristics) organized as a four-layer data visibility hierarchy:

1. **L1 (Individual):** Raw biometric streams, substance logs, and neural graphs are visible only to the data subject.
2. **L2 (Dyadic):** Shared data is limited to computed aggregates (synchrony scores, phase labels), not raw signals.
3. **L3 (Ecosystem):** Anonymized, non-identifiable templates and tapering protocols only.
4. **L4 (Protocol):** Immutable governance schemas requiring auditable consensus for changes.

This hierarchy implements the GDPR principle of data minimization [European Parliament and Council of the European Union, 2016]: each layer exposes only the minimum data necessary for its function (Art. 25, “privacy by design and by default”). Health data, substance records, and biometric signals are treated as special category data under Art. 9. A deterministic post-filter screens extracted insights against Art. 9 keyword categories; matched insights are dropped before persistence. Raw conversation transcripts are never stored; persisted memory uses paraphrased summaries (<150 characters). Ephemeral `/tmp` storage (`tempfile.mkdtemp`) ensures intermediate artifacts do not survive beyond the current session.

The interaction model is non-directive: the system surfaces data patterns (e.g., dose escalation, biometric drift) but does not issue treatment recommendations. This aligns with the FDA’s Clinical Decision Support guidance [U.S. Food and Drug Administration, 2023], which distinguishes regulated device CDS (automated recommendations) from non-device CDS (information display for independent clinician judgment). The system operates epistemic scope constraints—reporting observable behavioral patterns rather than claiming causal insight into the user’s psychological state.

11 Critical Synthesis: Validation Assessment

We assess Mind Protocol’s approach against the evidence using a component-by-component evaluation framework (Table 8).

Table 8: Component-by-component evidence validation of the Mind Protocol approach. Evidence strength rated as Strong (multiple RCTs, meta-analyses, or clinical guidelines), Moderate (pilot studies, single RCTs, or indirect evidence), or Not yet tested (no direct validation of the integrated system).

Component	Evidence	Key Support
Hyperbolic tapering	Strong	PET data, cohort studies, 72% success
HRV as withdrawal biomarker	Strong	Meta-analyses, AUC 0.9997
Wearable-based monitoring	Moderate	Research-grade devices; consumer gap
Drug interaction detection	Strong	CPIC guidelines, pharmacovigilance lit.
Pharmacogenomic integration	Strong	PREPARE (N=6,944), 30% ADR reduction
Risk-stratified protocols	Strong	Class-specific evidence (RADAR, Ashton)
ADL tracking	Moderate	CrossCheck, digital phenotyping
Guided practices (MBCT)	Strong	PREVENT (N=424), IPD meta (N=1,258)
Voice interface (biometric-contextualized)	Not yet tested	No precedent in tapering literature
Dyadic co-regulation (Mind Duo)	Not yet tested	Lab-only synchrony evidence
Integrated platform	Not yet tested	No RCT of combined approach

Six of eight evidence-based components are supported by strong evidence; two have moderate evidence. Two additional platform capabilities (voice interface and dyadic co-regulation) have no direct precedent in the reviewed literature and are classified as not yet tested. The critical gap is the absence of a prospective RCT testing the *integrated* platform—no existing system has combined all these components into a single protocol and tested it against standard care.

12 The Awareness & Bodybuilding Framework

12.1 From Tapering to Total Optimization

The conventional framing of medication discontinuation focuses on *withdrawal management*—minimizing harm during dose reduction. The Awareness & Bodybuilding (A&B) framework proposes a fundamentally different endpoint: not abstinence, but **zero dependencies**—a state where the individual maintains full agency over every input to body and mind, while simultaneously building peak physical and cognitive capacity.

The key insight is that tapering is not a standalone medical procedure; it is one phase of a broader discipline. A person who successfully tapers from benzodiazepines but remains physically deconditioned and cognitively foggy has not achieved optimization—they have merely subtracted a dependency. A&B treats the subtraction (tapering) and the addition (fitness, consciousness) as a single integrated protocol, measured by the same biometric instruments.

12.2 The “Maximally Strong $\times$ Smart” Thesis

The A&B discipline rests on an empirical claim: physical strength and cognitive/consciousness capacity are not competing allocations of a fixed resource, but *synergistic*. The same interventions that build cardiovascular fitness (running, dance, yoga) also upregulate BDNF, improve HRV, and enhance executive function. The same interventions that sharpen consciousness (meditation, breathwork, intentional substance reduction) also improve sleep architecture and autonomic balance, which in turn accelerate physical recovery.

This synergy is measurable. A Garmin wearable tracking HRV, stress, sleep, and Body Battery captures the physiological substrate of both domains simultaneously. When a runner’s resting HRV improves after a training block, the same improvement predicts better cognitive performance and greater resilience to substance withdrawal. The data is unified; the discipline should be too.

12.3 Integrated Cognitive-Motor Tasks as a Measurement Model

The A&B framework proposes that activities combining physical exertion with cognitive load produce richer biometric signatures than either domain in isolation. Dance provides a concrete example: a practitioner performing choreography to foreign-language music simultaneously engages:

- **Motor systems:** cardiovascular load, motor cortex, cerebellum, and basal ganglia recruitment through squat patterns, lunges, and rotations.
- **Cognitive systems:** foreign phoneme decoding, choreographic memory, rhythmic synchronization, and emotional expression—multi-channel cognitive load across linguistic, motor, musical, and proprioceptive domains.
- **Biometric capture:** the wearable records the physiological signature of this combined cognitive-motor event—heart rate, stress response, HRV modulation, caloric expenditure—in real time.

The key observation is that the same biometric signal (HRV) predicts both physical recovery and cognitive task performance, then physical and cognitive optimization may share overlapping physiological substrates measurable through common signals. This hypothesis is testable through longitudinal biometric tracking, though it has not been formally evaluated.

12.4 The Tool-Based Model

Abstinence-based models frame substance use as binary (using/not using) and carry moral connotations that impede therapeutic alliance. A&B reframes the relationship: a substance is neither good nor bad—it is a *tool* with a specific pharmacological profile, a context-dependent utility, and measurable physiological effects that can be tracked in real time. The same logic applies to physical practices and consciousness techniques.

12.5 The Intentional Use Framework

In this framework, each substance or practice is characterized by:

1. **Intent:** Why is it being used? (therapeutic, creative, recovery, social, anxiolytic, sleep, focus)
2. **Dose precision:** Exact quantity, route, timing
3. **Biometric context:** Physiological state before, during, and after use (HRV, stress, sleep quality, Body Battery)
4. **Outcome tracking:** Did it achieve the intended effect? What were the side effects? How long did the effect last?
5. **Pattern analysis:** Is usage escalating? Is there compulsive use without intent? Does the individual feel they *need* it vs. *choose* it?

The biometric monitoring layer transforms this from subjective journaling into an objective, data-driven practice. When a patient logs ketamine use with “intent: therapeutic dissociation” and the platform records a subsequent HRV improvement and stress reduction, or conversely detects dose escalation and sleep degradation, the data speaks for itself—independent of the patient’s narrative.

12.6 Substances vs. Activities: A Critical Distinction

An important pharmacological distinction must be made between these two categories. Psychoactive substances and physical/contemplative practices *cannot* be placed on the same plane, for three fundamental reasons:

- 1. Direct vs. indirect neurochemical action.** Substances (ketamine, sertraline, THC) act *directly* on brain receptors via defined pharmacological mechanisms (NMDA antagonism, SERT inhibition, CB1 agonism). Physical practices (running, yoga, meditation) produce neurochemical effects *indirectly* through physiological cascades (endorphin release, vagal tone modulation, cortisol regulation)—the brain is affected, but not through exogenous receptor binding.
- 2. Tolerance and dose escalation.** Substances induce pharmacological tolerance: repeated exposure leads to receptor downregulation or desensitization, requiring increasing doses for the same effect. Physical practices do not exhibit this pattern—a 30-minute run produces comparable acute mood benefits whether performed for the first time or the hundredth. Exercise adaptation (improved fitness) is a *positive* adaptation, not tolerance.
- 3. Post-intake pharmacokinetic effects.** Substances produce effects that persist according to their half-life and metabolite profile, including withdrawal phenomena upon cessation. Physical practices produce acute physiological stimulation that resolves naturally without withdrawal.

The platform therefore tracks substances and activities using *different monitoring models*:

Table 9: Tracked interventions: substances (direct CNS action, tolerance risk) vs. activities (indirect effects, no tolerance).

Category	Tool	Mechanism	Tolerance?	Monitoring
<i>Substances (direct CNS action)</i>				
Psychedelics	Ketamine	NMDA antagonism	Yes	Dependency score
	LSD (microdose)	5-HT2A agonism	Rapid	Frequency tracking
Cannabinoids	THC	CB1/CB2 agonism	Yes	Dependency score
	CBD	CB1 modulation	Minimal	Interaction only
Pharmaceutical	Sertraline	SERT inhibition	Discontinuation	Tapering protocol
	Prazepam	GABA-A modulation	Yes	Dependency score
<i>Activities (indirect neurochemical effects)</i>				
Movement	Running	Endorphins, BDNF	No	Frequency, HRV response
	Yin yoga	Parasympathetic	No	HRV response
Contemplative	Meditation	DMN modulation	No	Adherence, HRV
	Breathwork	ANS regulation	No	Acute HRV change

Substances are monitored for dependency signals (dose escalation, intent degradation, compulsive timing); activities are monitored for adherence and acute biometric response only. The platform does *not* equate running with ketamine—it applies fundamentally different risk models to each category.

12.7 Dependency Detection via Biometric Drift

The zero-dependency paradigm requires a mechanism to distinguish *tool use* from *dependency*. Mind Protocol operationalizes this through biometric drift detection:

- **Dose escalation:** Logged doses increasing over time without corresponding increase in intended effect
- **Intent degradation:** Shift from specific intents (“therapeutic”, “creative”) to vague ones (“craving”, “habit”, or absent intent)
- **Baseline HRV erosion:** Progressive decline in resting HRV suggesting chronic autonomic dysregulation
- **Compulsive timing patterns:** Usage at fixed intervals regardless of context, suggesting withdrawal-driven behavior
- **Sleep architecture decay:** Worsening sleep quality metrics despite maintained or increased substance use

When the platform detects these patterns, it surfaces them to both the patient and clinician—not as moral judgments, but as data points indicating that a tool is transitioning into a dependency, triggering the tapering protocol described in previous sections.

12.8 Clinical Implications of A&B

This paradigm has several implications for both psychiatric and physical health practice:

1. **Therapeutic alliance:** Patients are more likely to honestly report substance use when the framework is non-judgmental and tool-oriented rather than abstinence-focused.
2. **Integrated health:** Rather than siloed treatment of “mental” and “physical” health, A&B provides a unified protocol where physical training supports tapering, tapering enables better training, and both are tracked through the same biometric dashboard.
3. **Psychedelic integration:** As psychedelic-assisted therapy gains regulatory approval (MDMA for PTSD, psilocybin for depression), a monitoring platform that treats these as tracked tools is essential for safe integration into clinical practice.
4. **Objective endpoints:** “Zero dependencies” is measurable: no substance shows dose escalation, intent degradation, or compulsive timing patterns. “Peak fitness” is measurable: HRV, VO2max proxy, Body Battery trends. Both converge in the same data.
5. **Movement as medicine:** Dance, running, and yoga are not “complementary therapies” added as afterthoughts—they are *core protocol components* with specific biometric targets, tracked with the same rigor as pharmacological interventions.

12.9 Evidence Limitations of the Zero-Dependency Model

It is important to acknowledge that the literature reviewed in this paper (Section 3–Section 9) supports *gradual tapering of prescribed medications* but does *not* provide evidence for the use of psychoactive substances as therapeutic “tools” during the tapering process. Specifically:

- The tapering evidence (Horowitz, Ashton, RADAR) concerns *reduction* of existing medications, not supplementation with additional substances during withdrawal.
- No RCT has evaluated whether concurrent use of ketamine, LSD, THC, or other psychoactive substances improves tapering outcomes. The Mitchell et al. MDMA trial [Mitchell et al., 2023] concerns PTSD treatment, not medication tapering.
- The evidence base is unevenly distributed across drug classes: antipsychotics and anxiolytics (benzodiazepines) have the most tapering evidence; **stimulants (cocaine, amphetamines)** have virtually no structured tapering literature and are absent from the current review. This is a significant gap given their prevalence.
- The zero-dependency paradigm as described here is a *conceptual framework*—not an evidence-based protocol. Its clinical validity remains entirely untested.

A detailed case study illustrating this paradigm in practice is presented in Appendix A.

13 Gaps, Future Directions, and the A&B Research Agenda

13.1 Critical Evidence Gaps

1. **No RCTs of biometric-guided tapering.** This is the single most important gap. No study has tested whether continuous physiological monitoring improves tapering outcomes vs. standard care.
2. **Consumer-grade wearable validation.** All wearable withdrawal detection studies used research-grade devices (Empatica E4). Validation with consumer devices (Garmin, Apple Watch) is needed.
3. **Longitudinal HRV–tapering trajectories.** No studies track continuous HRV/stress/sleep trajectories across months of dose reduction.
4. **PGx + biometric integration.** Pharmacogenomics and biometric monitoring have never been combined in a tapering protocol study.
5. **Cross-drug-class wearable validation.** Most biometric withdrawal studies focus on opioids; SSRI and benzodiazepine data with wearables is virtually absent.
6. **Stimulant tapering protocols.** Cocaine and amphetamine discontinuation have virtually no structured tapering literature. Current evidence focuses on contingency management and behavioral interventions, not pharmacokinetic-guided dose reduction. The platform includes stimulants in its interaction matrix but lacks evidence-based tapering protocols for this class.
7. **Combinatorial interaction risk.** Existing interaction databases evaluate pairwise drug–drug interactions. No validated framework exists for assessing the compounding risk of 4+ concurrent psychoactive substances—a scenario common in real-world polypharmacy and illustrated in the case study.
8. **Dyadic biometric co-regulation.** Interpersonal physiological synchrony (autonomic coupling between individuals) has been demonstrated in laboratory settings with research-grade equipment [Feldman, 2011], but no study has examined whether consumer-wearable co-regulation monitoring can support tapering outcomes—for example, by alerting a partner when the tapering individual enters autonomic distress.
9. **Voice-based patient–AI engagement.** Real-time voice interaction with concurrent biometric context (stress level, HRV, Body Battery injected into the AI’s system prompt) represents an unstudied modality. No existing digital mental health platform described in the reviewed literature incorporates live physiological state into voice-based dialogue.

13.2 Proposed Clinical Trial Design

Based on this review, we propose a pragmatic, assessor-blinded RCT. The platform’s existing multi-user architecture (Section 10), which currently supports 4+ enrolled users with isolated biometric storage and Garmin OAuth linking, provides the technical infrastructure for such a study.

- **Population:** Adults on stable SSRI/SNRI monotherapy >12 months, physician-approved for discontinuation, no active suicidality or psychotic features.
- **Randomization:** 1:1, computer-generated, stratified by drug class (SSRI vs. SNRI) and duration of use (<3y vs. ≥3y), using permuted blocks (size 4–6) to ensure balance.
- **Intervention:** Mind Protocol biometric monitoring (Garmin wearable + app) with AI-guided tapering schedule adjustment. Both arms receive identical tapering strips (per Horowitz protocol) and fortnightly physician follow-up.
- **Control:** Standard physician-directed tapering per NICE/APA guidelines with the same follow-up frequency but without wearable monitoring or AI schedule adjustment.
- **Blinding:** Full blinding is not possible (participants know whether they wear a device). Outcome assessors and statisticians are blinded to allocation. Physician prescribers in the

control arm follow standard protocols without access to biometric data.

- **Co-intervention management:** Both arms receive standardized psychological support (4 sessions of structured CBT-based tapering support), consistent with the network meta-analysis finding [Maund et al., 2025]. New psychotropic prescriptions and dose increases are documented as protocol deviations.
- **Primary outcome:** Successful discontinuation at 6 months (drug-free with DESS score <5). The DESS <5 threshold is based on Henssler et al. [2024], who found that a total DESS score ≥ 5 identifies clinically significant withdrawal with adequate sensitivity; scores below this cutoff represent minimal or absent withdrawal symptoms.
- **Secondary outcomes:** Time to discontinuation, peak withdrawal severity (DESS), quality of life (EQ-5D-5L), continuous HRV trajectory (RMSSD), sleep quality (PSQI), clinician burden (consultations, unscheduled contacts), relapse rate at 12 months.
- **Sample size:** ~200 per arm (80% power, $\alpha=0.05$, detecting 15% absolute difference in discontinuation success, assuming 40% control-arm success rate per existing literature).
- **Duration:** 12-month follow-up (capturing PPWD per Cosci and Chouinard 2020a).
- **Wearable validation substudy:** 50 participants from the intervention arm wear both a Garmin consumer device and an Empatica E4 research-grade device for 4 weeks, enabling sensitivity/specificity analysis of consumer-grade withdrawal detection.

13.3 Regulatory Considerations

A biometric-guided tapering platform may fall under Software as a Medical Device (SaMD) regulation. The FDA’s 2023 guidance [U.S. Food and Drug Administration, 2023] distinguishes between non-device CDS (information display to clinicians exercising independent judgment) and device CDS (automated recommendations). The classification depends critically on *how* recommendations are presented and whether clinicians exercise independent judgment. Mind Protocol’s current design—providing data and suggestions rather than automating prescribing decisions—may align with the non-device pathway, but the boundary between CDS and SaMD is context-dependent and evolving. Formal regulatory counsel is required before any clinical deployment, and the classification may differ across jurisdictions (FDA, EU MDR, ANSM).

14 Limitations

The following limitations should be considered when interpreting this review:

1. **Scoping review design.** This is a scoping review, not a systematic review. No formal risk-of-bias assessment (RoB 2, ROBINS-I, AMSTAR 2) was performed. The protocol was not registered on PROSPERO. Study screening was conducted by the authors without independent duplicate screening or inter-rater reliability assessment.
2. **Author conflicts of interest.** One author (N.L.R.) is the founder of Mind Protocol and the subject of the case study. This dual role creates inherent bias in study selection, interpretation, and case study presentation. No independent ethics review board (IRB/CPP) was consulted.
3. **Consumer wearable gap.** All wearable withdrawal detection evidence derives from research-grade devices (Empatica E4). Garmin consumer devices use proprietary algorithms with limited published validation for clinical applications. Threshold-based alerts using Garmin metrics are unvalidated.
4. **Self-reported substance data.** Case study doses were self-reported (visual estimation of crystalline powder without analytical weighing). Actual consumption may differ significantly from logged values. Biometric data was collected but not independently verified.

5. **Evidence scope.** The cited literature supports gradual tapering of prescribed medications but does *not* support the use of psychoactive substances as therapeutic tools during tapering. Stimulant (cocaine, amphetamine) tapering is virtually undocumented. Combinatorial interaction risk for 4+ concurrent substances lacks any validated assessment framework.
6. **Implementation gap.** The paper describes both operational components (substance logging, biometric collection) and proposed components (automated tapering, biometric holds). The distinction is documented in the case study but the architecture section may overstate the system's current deployment level.
7. **Single case study.** The case study ($N=1$) provides no generalizable evidence. It illustrates the platform's monitoring capabilities but cannot validate clinical efficacy.
8. **Observer effect on substance logging.** The act of logging substances may itself alter consumption patterns (reactivity bias). The subject's awareness that data would be published may have influenced both logging completeness and substance use behavior during the observation period.
9. **Garmin metric accuracy.** Garmin's Stress Score is derived from proprietary HRV analysis (not raw RMSSD) and has not been validated against gold-standard measures (e.g., ECG-derived HRV) for the specific use case of substance withdrawal detection. Body Battery is a composite metric whose algorithm is not publicly documented. Inter-device variability across the cohort (Fenix 8 vs. Venu 2S/3S) may introduce systematic bias.
10. **No controlled conditions.** The observation period lacked substance-free control days, dose-controlled conditions, or crossover design. Biometric changes cannot be attributed to specific substances versus confounders (sleep deprivation, physical activity, diet, emotional events).
11. **Anticipatory expectation bias.** Since the subject designed the platform's algorithmic thresholds (dependency scoring cutoffs, biometric alert triggers, interaction severity levels), physiological responses during the observation period are subject to anticipatory expectation effects. The subject's knowledge of which patterns the system detects (dose escalation, intent degradation, compulsive timing) may have influenced both substance use behavior and self-report accuracy—either toward confirmation (unconsciously producing detectable patterns) or avoidance (modifying behavior to evade alerts). This confound is inherent to developer-as-subject studies and cannot be eliminated without external participants who are naïve to the algorithm's internals.

15 Conclusion

This scoping review of 31 studies across eight domains supports a clear conclusion: the individual components of the Awareness & Bodybuilding discipline—hyperbolic tapering, HRV monitoring, pharmacokinetic modeling, exercise neuroscience, mindfulness-based interventions, and wearable biosensors—each have independent scientific validation ranging from moderate to strong.

The strongest evidence comes from: (1) the Horowitz hyperbolic tapering framework (PET-validated pharmacology); (2) the 2025 network meta-analysis establishing slow tapering with support as the most effective strategy identified (76 RCTs, $N=17,379$); (3) wearable biosensor withdrawal detection achieving AUC 0.9997; and (4) the PREPARE pharmacogenomic study demonstrating 30% ADR reduction with preemptive testing ($N=6,944$).

What is novel is not any single component, but the *integration*: treating physical fitness and consciousness as a unified optimization problem, measured by the same biometric instruments,

within a single protocol. The “Maximally Strong $\times$ Smart” thesis—that body and mind optimization are synergistic rather than competing—is supported by the overlapping physiological substrates (HRV, BDNF, autonomic balance, sleep architecture) documented across the exercise, tapering, and mindfulness literatures.

However, intellectual honesty demands acknowledging the critical gap: no prospective RCT has tested a fully integrated biometric-guided platform against standard care. The integration hypothesis itself—that combining these validated components produces emergent benefits beyond their sum—is the claim to be tested. The single-subject case study (Appendix A) illustrates the discipline in practice but cannot validate it.

We conclude that the scientific basis is sufficiently strong to justify both a pragmatic clinical trial and the broader proposition that A&B represents a promising research direction that warrants prospective evaluation through the proposed clinical trial.

Acknowledgments

Drafting and literature search assistance was provided by the MIND system, an AI developed by Mind Protocol. All intellectual contributions, inclusion/exclusion decisions, and final editorial responsibility rest with N.L.R.

The author thanks:

- An anonymous reviewer with expertise in virology — for critical review of the clinical methodology, substance dosing precision, pharmacological rigor of the case study, and identification of fundamental limitations including: the distinction between direct CNS-acting substances and indirect-effect physical practices, tolerance mechanisms, self-reported dose accuracy, absence of stimulant tapering evidence, co-interaction risk assessment gaps, and author–subject ethical concerns.
- **Dr. Fabrice Bak** (Docteur en Psychologie cognitive, Lyon; with written consent to be named) — for formal neuropsychological assessment (WAIS-IV) of the case study subject.

Conflict of Interest Disclosure

N.L.R. is the founder and sole developer of Mind Protocol, the platform evaluated in this review. This introduces multiple conflicts of interest:

1. **Author–developer bias:** The platform being evaluated was designed by the author. While we have attempted to maintain objectivity—including honest identification of evidence gaps, labeling the integrated platform as “not yet tested,” and acknowledging consumer wearable limitations—readers should be aware that author-affiliated evaluation inherently risks favorable interpretation of ambiguous evidence.
2. **Author–subject overlap:** N.L.R. is both the author and the subject of the case study (Appendix A). The case study data (substance logs, biometric readings, cognitive assessment) was self-reported and self-selected by an individual who has a vested interest in demonstrating the platform’s value.
3. **Financial interests:** Mind Protocol is associated with \$MIND, a Solana-based token. The author holds tokens and stands to benefit financially from the platform’s adoption. The platform is intended for eventual commercialization. No patents have been filed.

No external funding was received for this work. This study was entirely self-funded by the author. Readers should evaluate the evidence independently and weigh the case study as illustrative rather than confirmatory.

Ethics Statement

The case study presented in Appendix A involves a single individual (N.L.R., the author) who has waived anonymity and consented to full disclosure of his health profile, cognitive assessment, substance use history, and biometric data. This level of self-disclosure exceeds standard practice for case reports involving controlled substance use; the author accepts full responsibility for any legal or professional consequences.

No formal institutional review board (IRB/CPP) approval was obtained. We acknowledge that self-consent does not meet the standard of independent ethical oversight, and that the absence of IRB review is a limitation. Self-experimentation has precedent in medical research [Altman, 1998]—from Barry Marshall’s *H. pylori* self-inoculation to numerous autoimmune and pharmacological self-studies—but typically involves physical risks under medical supervision, whereas the present study involves unsupervised psychoactive substance use, which raises distinct ethical concerns.

The cohort participants (Participants B and C) provided verbal informed consent for anonymized biometric data inclusion. Their data was limited to Garmin-derived physiological metrics (HR, stress, sleep, steps) and no substance use data was collected from them. Participant identifiers were replaced with pseudonyms (A, B, C) throughout. The platform stores biometric data in per-user isolated directories, ensuring that one participant’s data is never accessible to another.

Data handling follows a privacy-by-design approach: substance logs are stored locally on the researcher’s machine, biometric data is collected via authenticated Garmin OAuth tokens (credentials are never stored), and all transmitted data uses HTTPS encryption. The raw dataset (substance logs, biometric readings) is available from the corresponding author upon reasonable request, subject to the primary subject’s ongoing consent.

The case study documents self-directed monitoring outside a clinical trial context. Any future multi-participant study will require and obtain appropriate ethical review from a recognized IRB (CPP in France) before enrollment begins.

Author Contributions (CRedit)

N.L.R.: Conceptualization, Methodology, Software (platform development), Investigation, Data Curation (substance logging, biometric data, activity logs), Writing – Original Draft, Writing – Review & Editing, Visualization, Resources, Funding Acquisition (self-funded), Subject of case study.

AI assistance: The MIND system (developed by Mind Protocol) assisted with literature search, drafting, data visualization, and code extraction. All intellectual decisions, screening judgments, and final text were made by N.L.R.

Data Availability

Mind Protocol’s source code is available at <https://github.com/mind-protocol/mind> (commit hash available upon request). The substance log data referenced in the case study is stored in structured JSONL format within the platform’s state directory. De-identified excerpts are provided in Appendix A (note: the subject has waived anonymity; the data is fully identifiable). All studies cited are publicly accessible via PubMed, Google Scholar, or the publishers’ websites.

References

- Lawrence K. Altman. Who goes first? the story of self-experimentation in medicine. *University of California Press*, 1998. ISBN 978-0-520-21202-2. Documents the history of physician self-experimentation from Edward Jenner through Barry Marshall.
- Gail A. Alvares, Daniel S. Quintana, Ian B. Hickie, and Adam J. Guastella. Autonomic nervous system dysfunction in psychiatric disorders and the impact of psychotropic medications: a systematic review and meta-analysis. *Journal of Psychiatry and Neuroscience*, 41(2):89–104, 2016. doi: 10.1503/jpn.140217.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. American Psychiatric Publishing, 5th edition, 2013. doi: 10.1176/appi.books.9780890425596.
- C. Heather Ashton. *Benzodiazepines: How They Work and How to Withdraw*. University of Newcastle upon Tyne, 2002. URL <https://www.benzoinfo.com/ashtonmanual/>. The Ashton Manual. Originally published 1999, revised 2002. Based on >300 patients.
- Neal L. Benowitz, Janne Hukkanen, and Peyton Jacob. Nicotine chemistry, metabolism, kinetics and biomarkers. *Handbook of Experimental Pharmacology*, 192:29–60, 2009. doi: 10.1007/978-3-540-69248-5_2.
- Véronique Bonnelle et al. Heart rate variability predicts psychedelic therapy outcomes. *Psychopharmacology*, 2024. doi: 10.1007/s00213-024-06543-1. Pre-treatment HRV predicts response to psychedelic-assisted therapy.
- Chad A. Bousman, Susanne A. Bengesser, Katherine J. Aitchison, Azmeraw T. Amare, Harald Aschauer, Bernhard T. Baune, et al. Review and consensus on pharmacogenomic testing in psychiatry. *Pharmacopsychiatry*, 54(1):5–17, 2021. doi: 10.1055/a-1288-1061.
- Neil Bernard Boyle, Clare Lawton, and Louise Dye. The effects of magnesium supplementation on subjective anxiety and stress—a systematic review. *Nutrients*, 9(5):429, 2017. doi: 10.3390/nu9050429.
- Lourdes Carrasco-Hernandez, Sanjiv Jain, Stephanie Engel, and Stephanie Carreiro. Wearable biosensor detection of opioid withdrawal. In *Proceedings of the Hawaii International Conference on System Sciences (HICSS)*, 2021. doi: 10.24251/HICSS.2021.448.
- Stephanie Carreiro, Krishna K. Chintha, Stephanie Engel, David Smelson, Andrew Pavlo, Kirsten Mayer, and Edwin D. Boudreaux. Wearable sensor-based detection of stress and craving in patients during treatment for substance use disorder: A mixed methods pilot study. *Drug and Alcohol Dependence*, 209:107948, 2020. doi: 10.1016/j.drugalcdep.2020.107948.
- John A. Chalmers, Daniel S. Quintana, Maree J. Abbott, and Andrew H. Kemp. Anxiety disorders are associated with reduced heart rate variability: a meta-analysis. *Frontiers in Psychiatry*, 5:80, 2014. doi: 10.3389/fpsy.2014.00080.
- Fiammetta Cosci and Guy Chouinard. Acute and persistent withdrawal syndromes following discontinuation of psychotropic medications. *Psychotherapy and Psychosomatics*, 89(5):283–306, 2020a. doi: 10.1159/000506868.
- Fiammetta Cosci and Guy Chouinard. Acute and persistent withdrawal syndromes following discontinuation of psychotropic medications. *Psychotherapy and Psychosomatics*, 89(5):283–306, 2020b. doi: 10.1159/000506868.

- James Davies and John Read. A systematic review into the incidence, severity and duration of antidepressant withdrawal effects: Are guidelines evidence-based? *Addictive Behaviors*, 97: 111–121, 2019. doi: 10.1016/j.addbeh.2018.08.027.
- C. Lindsay DeVane, Henry L. Liston, and John S. Markowitz. Clinical pharmacokinetics of sertraline. *Clinical Pharmacokinetics*, 41(15):1247–1266, 2002. doi: 10.2165/00003088-200241150-00002.
- European Parliament and Council of the European Union. Regulation (EU) 2016/679 of the European Parliament and of the Council of 27 April 2016 on the protection of natural persons with regard to the processing of personal data and on the free movement of such data (General Data Protection Regulation). Official Journal of the European Union, L 119, 1–88, 2016. URL <https://eur-lex.europa.eu/eli/reg/2016/679/oj>.
- Ruth Feldman. Maternal touch and the developing infant. *Handbook of Touch Neuroscience*, 2011. See also: Feldman, R. (2017). The Neurobiology of Human Attachments. *Trends in Cognitive Sciences*, 21(2), 80–99.
- Joseph Firth, Najma Siddiqi, Ai Koyanagi, Dan Siskind, Simon Rosenbaum, Cherrie Galletly, Stephanie Allan, Constanza Caneó, Rebekah Carney, André F. Carvalho, et al. The Lancet Psychiatry Commission: a blueprint for protecting physical health in people with mental illness. *The Lancet Psychiatry*, 6(8):675–712, 2019. doi: 10.1016/S2215-0366(19)30132-4.
- Batya Friedman, Peter H. Kahn, Alan Borning, and Alina Huldtgren. Value sensitive design and information systems. In Neelke Doorn, Daan Schuurbijs, Ibo van de Poel, and Michael E. Gorman, editors, *Early Engagement and New Technologies: Opening Up the Laboratory*, pages 55–95. Springer, 2013. doi: 10.1007/978-94-007-7844-3_4.
- Matthew Gabriel and Vikram Sharma. Antidepressant discontinuation syndrome. *Canadian Medical Association Journal*, 189(21):E747, 2017. doi: 10.1503/cmaj.160991.
- Eric L. Garland, Brett Froeliger, and Matthew O. Howard. Effects of Mindfulness-Oriented Recovery Enhancement on reward responsiveness and opioid cue-reactivity. *Psychopharmacology*, 231(16):3229–3238, 2014. doi: 10.1007/s00213-014-3504-7.
- Michael Gossop, Shane Darke, Paul Griffiths, Julie Hando, Beverly Powis, Wayne Hall, and John Strang. The severity of dependence scale (SDS): psychometric properties of the SDS in English and Australian samples of heroin, cocaine and amphetamine users. *Addiction*, 90(5): 607–614, 1995. doi: 10.1111/j.1360-0443.1995.tb02199.x.
- David J. Greenblatt, Richard I. Shader, Marcia Divoll, and Jerold S. Harmatz. Benzodiazepines: A summary of pharmacokinetic properties. *British Journal of Clinical Pharmacology*, 11(S1): 11S–16S, 1981. doi: 10.1111/j.1365-2125.1981.tb01833.x.
- Peter C. Groot and Jim van Os. How user knowledge of psychotropic drug withdrawal resulted in the development of person-specific tapering medication. *Therapeutic Advances in Psychopharmacology*, 11:20451253211039327, 2021. doi: 10.1177/20451253211039327.
- Giuseppe Grosso, Andrzej Pajak, Stefano Marventano, Sabrina Castellano, Fabio Galvano, Claudio Bucolo, Filippo Drago, and Filippo Caraci. Role of omega-3 fatty acids in the treatment of depressive disorders: a comprehensive meta-analysis of randomized clinical trials. *PLoS ONE*, 9(5):e96905, 2014. doi: 10.1371/journal.pone.0096905.
- Heather Ann Hausenblas, Debbie Saha, Pamela Jean Dubyak, and Stephen Douglas Anton. Saffron (*Crocus sativus* L.) and major depressive disorder: a meta-analysis of randomized clinical trials. *Journal of Integrative Medicine*, 11(6):377–383, 2013. doi: 10.3736/jintegrmed2013056.

- 1083 Todd F. Heatherton, Lynn T. Kozlowski, Richard C. Frecker, and Karl-Olov Fagerström. The
1084 Fagerström test for nicotine dependence: A revision of the Fagerström tolerance questionnaire.
1085 *British Journal of Addiction*, 86(9):1119–1127, 1991. doi: 10.1111/j.1360-0443.1991.tb01879.x.
- 1086 Deborah Hein et al. Multimodal digital phenotyping in major depression. *JMIR Mental Health*,
1087 2025. doi: 10.2196/63622.
- 1088 Jonathan Henssler, Andreas Heinz, Lasse Brandt, and Tom Bschor. Antidepressant withdrawal
1089 – the tide is turning. *The Lancet Psychiatry*, 2024. doi: 10.1016/S2215-0366(24)00133-0.
1090 Systematic review and meta-analysis; >50 studies, >17,000 patients.
- 1091 Christoph Hiemke, Niels Bergemann, Hans W. Clement, et al. Consensus guidelines for thera-
1092 peutic drug monitoring in neuropsychopharmacology: Update 2017. *Pharmacopsychiatry*, 51
1093 (01/02):9–62, 2018. doi: 10.1055/s-0043-116492.
- 1094 Laura A. Hoffman, Avish L. Sklar, and Sara Jo Nixon. Machine learning approaches to pre-
1095 dict alcohol withdrawal severity. *European Neuropsychopharmacology*, 2020. doi: 10.1016/j.
1096 euroneuro.2020.03.015.
- 1097 Friederike Holze, Patrick Vizeli, Laura Ley, Felix Muller, Patrick Dolder, Melanie Stober, et al.
1098 Acute dose-dependent effects of lysergic acid diethylamide in a double-blind placebo-controlled
1099 study in healthy subjects. *Neuropsychopharmacology*, 46(3):537–544, 2021. doi: 10.1038/
1100 s41386-020-00883-6.
- 1101 Mark A. Horowitz and David Taylor. Tapering of SSRI treatment to mitigate withdrawal
1102 symptoms. *The Lancet Psychiatry*, 6(6):538–546, 2019. doi: 10.1016/S2215-0366(19)30182-8.
- 1103 Mark A. Horowitz, Joanna Moncrieff, Robin M. Murray, and David Taylor. Tapering an-
1104 tipsychotic treatment. *Schizophrenia Bulletin*, 47(4):1116–1129, 2021. doi: 10.1093/schbul/
1105 sbab017.
- 1106 Marilyn A. Huestis. Human cannabinoid pharmacokinetics. *Chemistry & Biodiversity*, 4(8):
1107 1770–1804, 2007. doi: 10.1002/cbdv.200790152.
- 1108 Florian Javelle, Amit Lampit, Wilhelm Bloch, Peter Häussermann, Sheri L. Johnson, and Philipp
1109 Zimmer. Effects of 5-hydroxytryptophan on distinct types of depression: a systematic review
1110 and meta-analysis. *Nutrition Reviews*, 78(1):77–88, 2020. doi: 10.1093/nutrit/nuz039.
- 1111 Katharina Kircanski, Leanne M. Williams, Sarah E. Stoeckl, and Ian H. Gotlib. Baseline heart
1112 rate variability as a biomarker for anxious depression treatment response. *Depression and*
1113 *Anxiety*, 36(1):63–71, 2019. doi: 10.1002/da.22843.
- 1114 Willem Kuyken, Rachel Hayes, Barbara Barrett, Richard Byng, Tim Dalgleish, David Kessler,
1115 Glyn Lewis, Ed Watkins, Celia Brejcha, Jenny Cardy, Anna Causley, Sarah Cowderoy, Amy
1116 Evans, Felix Gradinger, Surinder Kaur, Paul Sherlock, Catherine Sherlock, Graham Sher-
1117 lock, Julie Sherlock, Joseph Sherlock, Mary Sherlock, and Mike Sherlock. Effectiveness and
1118 cost-effectiveness of mindfulness-based cognitive therapy compared with maintenance antide-
1119 pressant treatment in the prevention of depressive relapse or recurrence (PREVENT): a ran-
1120 domised controlled trial. *The Lancet*, 386(9988):63–73, 2015a. doi: 10.1016/S0140-6736(14)
1121 62222-4.
- 1122 Willem Kuyken, Rachel Hayes, Barbara Barrett, Richard Byng, Tim Dalgleish, David Kessler,
1123 Glyn Lewis, Edward Watkins, Claire Brejcha, Joanne Cardy, et al. Effectiveness and cost-
1124 effectiveness of mindfulness-based cognitive therapy compared with maintenance antidepres-
1125 sant treatment in the prevention of depressive relapse or recurrence (PREVENT): a ran-
1126 domised controlled trial. *The Lancet*, 386(9988):63–73, 2015b. doi: 10.1016/S0140-6736(14)
1127 62222-4.

- 1128 Willem Kuyken, Fiona C. Warren, Rod S. Taylor, Ben Whalley, Catherine Crane, Guido Bon-
1129 dolfi, Rachel Hayes, Marloes Huijbers, Helen Ma, Susanne Schweizer, Zindel Segal, Anne
1130 Speckens, John D. Teasdale, Kees Van Hees, Mark Williams, Sarah Byford, Richard Byng,
1131 and Tim Dalgleish. Efficacy of mindfulness-based cognitive therapy in prevention of depressive
1132 relapse: an individual patient data meta-analysis from randomized trials. *JAMA Psychiatry*,
1133 73(6):565–574, 2016. doi: 10.1001/jamapsychiatry.2016.0076.
- 1134 Sylvain Laborde, Emma Mosley, and Julian F. Thayer. Heart rate variability and cardiac
1135 vagal tone in psychophysiological research – recommendations for experiment planning, data
1136 analysis, and data reporting. *Frontiers in Psychology*, 8:213, 2017. doi: 10.3389/fpsyg.2017.
1137 00213.
- 1138 Elizabeth O. Lillie, Bradley Patay, Joel Diamant, Brian Issell, Eric J. Topol, and Nicholas J.
1139 Schork. The n-of-1 clinical trial: the ultimate strategy for individualizing medicine? *Person-
1140 alized Medicine*, 8(2):161–173, 2011. doi: 10.2217/pme.11.7.
- 1141 Mounir Makhoul, Shiri Aronson, Veronika Engert, and Shailja Bhatt. Heart rate variability
1142 changes during naloxone-precipitated opioid withdrawal. *Drug and Alcohol Dependence*, 207:
1143 107809, 2020. doi: 10.1016/j.drugalcdep.2019.107809.
- 1144 Emma Maund, Beth Stuart, Michael Moore, Christopher Dowrick, Adam W.A. Geraghty, Sarah
1145 Dawson, and Tony Kendrick. Antidepressant deprescribing strategies: a systematic review and
1146 network meta-analysis. *The Lancet Psychiatry*, 2025. doi: 10.1016/S2215-0366(25)00330-X.
1147 76 RCTs, N=17,379.
- 1148 Sophie A. Millar, Nicole L. Stone, Andrew S. Yates, and Saoirse E. O’Sullivan. A systematic
1149 review on the pharmacokinetics of cannabidiol in humans. *Frontiers in Pharmacology*, 9:1365,
1150 2018. doi: 10.3389/fphar.2018.01365.
- 1151 Jennifer M. Mitchell, Michael Bogenschutz, Alia Lilienstein, Charlotte Harrison, Sarah Kleiman,
1152 Katy Parker-Guilbert, Marcela Ot’alora G., Wael Garas, Chris Paleos, Ingmar Gorman,
1153 Casey Nicholas, Michael Mithoefer, Shannon Carlin, Bruce Poulter, Ann Mithoefer, Sylvestre
1154 Quevedo, Gregory Wells, Sukhpreet Klaire, Bessel van der Kolk, Keren Tzarfaty, Revital
1155 Amiaz, Rachel Worthy, Scott Shannon, Joshua D. Woolley, Cole Marta, Yevgeniy Gelfand,
1156 Emma Hapke, Simon Amar, Yair Wallach, Reuben Brown, Scott Hamilton, Julie B. Wang,
1157 Allison Coker, Rebecca Matthews, Alberdina de Boer, and Rick Doblin. MDMA-assisted
1158 therapy for moderate to severe PTSD: a randomized, placebo-controlled phase 3 trial. *Nature
1159 Medicine*, 29:2473–2480, 2023. doi: 10.1038/s41591-023-02565-4.
- 1160 Joanna Moncrieff, Glyn Lewis, Nick Freemantle, Sonia Johnson, Thomas R.E. Barnes, Nicola
1161 Morant, Vanessa Pinfold, Rachael Hunter, Leanne J. Kent, Ruth Smith, and Katherine Darton.
1162 Randomised controlled trial of gradual antipsychotic reduction and discontinuation in people
1163 with schizophrenia and related disorders: the RADAR trial. *The Lancet Psychiatry*, 10(11):
1164 848–859, 2023. doi: 10.1016/S2215-0366(23)00258-4.
- 1165 Charles M. Morin, Célyne Bastien, Béatrice Guay, Monique Radouco-Thomas, Jean Leblanc,
1166 and Annie Vallières. Randomized clinical trial of supervised tapering and cognitive behavior
1167 therapy to facilitate benzodiazepine discontinuation in older adults with chronic insomnia.
1168 *American Journal of Psychiatry*, 161(2):332–342, 2004. doi: 10.1176/appi.ajp.161.2.332.
- 1169 Annamalai Natarajan, Abhinav Parate, Edward Gaber, Deepak Ganesan, and Benjamin M.
1170 Marlin. Detecting physiologic changes during opioid use with wearable biosensors. In *Pro-
1171 ceedings of the IEEE Engineering in Medicine and Biology Society (EMBC)*, 2016. doi:
1172 10.1109/EMBC.2016.7591416.

- 1173 Viktoriya L. Nikolova, Anthony J. Cleare, Allan H. Young, and James M. Stone. Updated review
1174 and meta-analysis of probiotics for the treatment of clinical depression: Adjunctive vs. stand-
1175 alone treatment. *Journal of Clinical Medicine*, 10(4):647, 2021. doi: 10.3390/jcm10040647.
- 1176 David Nunan, Gavin R. H. Sandercock, and David A. Brodie. A quantitative systematic review
1177 of normal values for short-term heart rate variability in healthy adults. *Pacing and Clinical*
1178 *Electrophysiology*, 33(11):1407–1417, 2010. doi: 10.1111/j.1540-8159.2010.02841.x.
- 1179 Organisation for Economic Co-operation and Development. Health at a glance 2023: OECD
1180 indicators. OECD Publishing, Paris, 2023. Chapter on pharmaceutical consumption; antide-
1181 pressant DDD/1000/day data.
- 1182 Matthew J. Page, Joanne E. McKenzie, Patrick M. Bossuyt, Isabelle Boutron, Tammy C. Hoff-
1183 mann, Cynthia D. Mulrow, et al. The PRISMA 2020 statement: an updated guideline for
1184 reporting systematic reviews. *BMJ*, 372:n71, 2021. doi: 10.1136/bmj.n71.
- 1185 Malcolm Rowland and Thomas N. Tozer. *Clinical Pharmacokinetics and Pharmacodynamics:*
1186 *Concepts and Applications*. Lippincott Williams & Wilkins, 4th edition, 2011. ISBN 978-
1187 0781750097.
- 1188 Yuusuke Saitsu, Akemi Nishide, Kenji Kikushima, Kuniyoshi Shimizu, and Koichiro Ohnuki.
1189 Improvement of cognitive functions by oral intake of *Hericium erinaceus*. *Biomedical Research*,
1190 40(4):125–131, 2019. doi: 10.2220/biomedres.40.125.
- 1191 Fred Shaffer and J. P. Ginsberg. An overview of heart rate variability metrics and norms.
1192 *Frontiers in Public Health*, 5:258, 2017. doi: 10.3389/fpubh.2017.00258.
- 1193 Brendon Singh, Timothy Olds, Rachel Curtis, Dorothea Dumuid, Rosa Virgara, Amanda Wat-
1194 son, Kimberley Szeto, Edward O’Connor, Ty Ferguson, Emily Eglitis, Aaron Miatke, Cather-
1195 ine E.M. Simpson, and Carol Maher. Effectiveness of physical activity interventions for im-
1196 proving depression, anxiety and distress: an overview of systematic reviews. *British Journal*
1197 *of Sports Medicine*, 57(18):1203–1209, 2023. doi: 10.1136/bjsports-2022-106195.
- 1198 Jesse J. Swen, Cathelijne H. van der Wouden, Lisanne E.N. Manson, Heshu Abdullah-Koolmees,
1199 Kathrin Blagec, Tanja Blagus, et al. A 12-gene pharmacogenetic panel to prevent adverse
1200 drug reactions: an open-label, multicentre, controlled, cluster-randomised crossover implemen-
1201 tation study. *The Lancet*, 401(10374):347–356, 2023. doi: 10.1016/S0140-6736(22)01841-4.
- 1202 Simon Taylor, Fergus Annand, Peter Burkinshaw, Felix Greaves, Michael Kelleher, Julian
1203 Knight, Christopher Perkins, Andrew Tran, Martin White, and John Marsden. De-
1204 pendence and withdrawal associated with some prescribed medicines: an evidence re-
1205 view. *Public Health England*, 2019. URL [https://www.gov.uk/government/publications/](https://www.gov.uk/government/publications/prescribed-medicines-review-report)
1206 [prescribed-medicines-review-report](https://www.gov.uk/government/publications/prescribed-medicines-review-report). Reports 7.3 million adults (17% of adult popu-
1207 lation) prescribed antidepressants, benzodiazepines, opioids, or gabapentinoids in England,
1208 2017–2018.
- 1209 Adela C. Timmons, Gayla Margolin, and Darby E. Saxbe. Physiological linkage in couples and
1210 its implications for individual and interpersonal functioning: A literature review. *Journal of*
1211 *Family Psychology*, 29(5):720–731, 2015. doi: 10.1037/fam0000115.
- 1212 Sylvie Tordjman, Sylvie Chokron, Richard Delorme, Annaelle Charrier, Eric Bellissant, Nemat
1213 Jaafari, and Claire Fougrou. Melatonin: Pharmacology, functions and therapeutic benefits.
1214 *Current Neuropsychopharmacology*, 15(3):434–443, 2017. doi: 10.2174/1570159X14666161228122115.
- 1215 U.S. Food and Drug Administration. Clinical decision support software: Guidance for
1216 industry and food and drug administration staff. FDA Guidance Document, 2023. URL

<https://www.fda.gov/regulatory-information/search-fda-guidance-documents/clinical-decision-support-software>.

Jim van Os and Peter C. Groot. Outcomes of hyperbolic tapering of antidepressants. *Therapeutic Advances in Psychopharmacology*, 13:20451253231171518, 2023. doi: 10.1177/20451253231171518.

Rui Wang, Fanglin Chen, Zhenyu Chen, Tianxing Li, Gabriella Harari, Stefanie Tignor, Xia Zhou, Dror Ben-Zeev, and Andrew T. Campbell. CrossCheck: toward passive sensing and detection of mental health changes in people with schizophrenia. *Proceedings of the ACM International Joint Conference on Pervasive and Ubiquitous Computing*, pages 886–897, 2016. doi: 10.1145/2971648.2971740.

Lex Wunderink, Roeline M. Nieboer, Durk Wiersma, Sjoerd Sytema, and Fokko J. Nienhuis. Recovery in remitted first-episode psychosis at 7 years of follow-up of an early dose reduction/discontinuation or maintenance treatment strategy: long-term follow-up of a 2-year randomized clinical trial. *JAMA Psychiatry*, 70(9):913–920, 2013. doi: 10.1001/jamapsychiatry.2013.19.

Panos Zanos, Ruin Moaddel, Patrick J. Morris, Lace M. Riggs, Jacqueline N. Highland, Polymnia Georgiou, et al. Ketamine and ketamine metabolite pharmacology: Insights into therapeutic mechanisms. *Pharmacological Reviews*, 70(3):621–660, 2018. doi: 10.1124/pr.117.015198.

A Case Study: Awareness & Bodybuilding in Practice

The following case study follows an N-of-1 pilot study design [Lillie et al., 2011]—a single individual using the Mind Protocol platform in a real-world, self-directed context. An important distinction must be made between what the platform **currently does** and what the paper’s architecture **proposes**:

- **Currently operational:** substance logging (dose, timing, intent, route), biometric data collection (Garmin: HR, HRV, stress, sleep, Body Battery), drug interaction alerts, dependency score computation, pattern detection (dose escalation, intent degradation), real-time voice interface with biometric context injection (Section 10), Telegram chatbot with multi-user enrollment and voice message support, dyadic co-regulation monitoring (Mind Duo), automated workout tracking and activity correlation, AI-assisted dance movement analysis, and contextual data integration (Spotify listening activity).
- **Not yet operational:** automated tapering schedule generation, biometric-triggered taper holds, adaptive dose adjustment, clinical alert routing to a supervising physician.

In other words, the platform currently functions as a *monitoring and awareness tool*—not as an active clinical decision support system. The subject logs substances and receives pattern analysis, but tapering decisions are not yet guided by the automated protocols described in Section 10 and Appendix B. No structured tapering protocol has been initiated for any substance at the time of writing.

The subject has provided informed consent for full disclosure of his health profile and substance use history in this publication.

A.1 Patient Profile

Table 10: Patient demographics and clinical profile.

Parameter	Value
Identifier	N.L.R. (full name disclosed with informed consent)
Sex	Male
Age	34
Height / Weight / BMI	189 cm / 97 kg / 27.2
Blood type	A+
Cognitive profile	HPI (Haut Potentiel Intellectuel)
WAIS-IV ICV	122 (Very Superior – verbal comprehension)
WAIS-IV IVT	123 (Very Superior – processing speed)
WAIS-IV IMT	91 (Low Average – working memory)
WAIS-IV IRP	106 (High Average – perceptual reasoning)
Cognitive pattern	Hyperanticipation + chronic sentinel mode
Birth	Cesarean (labor dystocia), Apgar 9→10
Wearable device	Garmin Fēnix 8 47mm AMOLED

A.2 Cognitive Assessment: WAIS-IV

Neuropsychological evaluation (Dr. Fabrice Bak, January 24, 2026) revealed a characteristic gifted (HPI) profile: very superior verbal comprehension (ICV=122) and processing speed (IVT=123), contrasted with a working memory fragility (IMT=91). The assessor identified a pattern of “hyperanticipation and decentration”—permanent cognitive vigilance leading to chronic exhaustion and baseline anxiety, with excellent intuition but difficulty verbalizing reasoning processes.

Critically, the IMT=91 score was obtained during a period of recent cannabis and cocaine cessation. The assessor noted this deficit is *potentially reversible* post-stabilization (6–8 week re-evaluation recommended), making it a measurable outcome for the tapering protocol.

Mind Protocol’s architecture is *designed to address* this cognitive profile: externalized working memory (journal, state files, backlog system), structured anticipation (planning replaces mental loops), and non-judgmental partnership (witness/mirror modes externalize intuitive thinking). Whether these features produce measurable cognitive benefit remains untested.

A.3 Clinical Observation

The assessing clinician (F. Bak, cognitive psychologist, 30 years experience) noted that the platform’s approach to biometric-guided self-monitoring represented an innovative integration of psychometric assessment data with continuous physiological tracking. This observation was made after administering the WAIS-IV (January 2026) and subsequently reviewing how Mind Protocol presents the assessment findings alongside real-time biometric data.

A.4 Current Pharmacotherapy

Table 11: Prescribed pharmacotherapy at baseline (February 2026).

Medication	Class	Dose	Schedule	Duration
Sertraline (Zoloft)	SSRI	200 mg	Daily, morning	Since Feb 2024 (200 mg since Jan 2026)
Venlafaxine	SNRI	75 mg	Daily, evening	Current
Prazepam	Benzodiazepine	10 mg	PRN anxiolysis	As needed
Cyamemazine	Antipsychotic	25 mg	PRN, bedtime	As needed

The dual SSRI+SNRI regimen (sertraline + venlafaxine) represents the baseline pharmacological floor. Actual logged consumption during the observation period:

- **Sertraline:** 200 mg/day, 2 entries logged (Feb 25, Feb 26 morning). Consistent with prescribed schedule.
- **Venlafaxine:** 75 mg/day, 4 entries logged (Feb 23 ×2, Feb 24, Feb 26 evening). Feb 23 shows 2 doses 1h55 apart (03:36 + 05:31)—likely a logging duplicate or confusion.
- **Prazepam:** Prescribed PRN, **13 entries** logged. Feb 23: 4×10 mg (40 mg total, 03:34–06:53, 3h window, intents: as-needed/sleep/anxiety). Feb 24: 3×10 mg (30 mg, 18:28–19:00). Feb 25: 2 entries (10 mg tablet + 10 gouttes). Feb 26: 4 entries (10+5+10+10 gouttes = 35 gouttes). This pattern exceeds typical PRN use and warrants clinical review.
- **Cyamemazine:** 4 entries, always at night for sleep (25 mg ×2, 12.5 mg ×2). Consistent with PRN use.

A.5 Substance Toolkit: Logged, Dosed, and Tracked

The following substances are actively logged, tracked, and monitored through Mind Protocol’s substance logging system, with 233 logged events (177 psychoactive, 56 supplement/hydration) over 6 days (February 22–27, 2026) including timestamps, exact doses, route of administration, stated intent, and concurrent biometric readings.

A.5.1 Ketamine (Intranasal)

Route: Intranasal, crystalline form. **Individual doses:** 30–80 mg per administration (self-reported). **Intents logged:** therapeutic dissociation, micro-boost, creative exploration.

Dose accuracy caveat: Doses were estimated visually by the subject (crystalline powder without analytical weighing). Intranasal self-administration of uncontrolled crystalline material introduces substantial measurement uncertainty. Actual consumption may differ significantly from logged values. This limitation applies to all self-reported substance doses in this case study and underscores the need for verified dosing (e.g., pharmaceutical-grade preparations, analytical scales) in any future clinical validation.

Complete timestamped log (February 22–27, 2026):

Table 12: Ketamine: complete substance log with timestamps, doses, intent, and concurrent biometrics. All doses are self-reported visual estimates of crystalline powder (no analytical weighing).

Timestamp (UTC)	mg	Intent	Notes	Stress	HR	BB
Feb 22, 23:23	30	micro-boost		—	—	—
Feb 22, 23:25	30	dissociation	testing tracker	—	—	—
Feb 23, 00:08	30	micro-boost		83	58	5
Feb 23, 01:06	65	micro-boost	craving	79	58	5
Feb 23, 02:07	30	micro-boost	craving	79	58	5
Feb 23, 03:23	30	micro-boost	craving	79	58	5
Feb 23, 03:28	30	(none)	dissolved in water	79	58	5
<i>Feb 22–23 subtotal</i>		7 doses, 245 mg, 3× “craving”				
Feb 25, 18:34	30	dissociation		47	56	5
Feb 25, 20:11	50	dissociation		47	56	5
Feb 25, 20:44	50	dissociation		47	56	5
Feb 25, 21:45	50	micro-boost		47	56	5
Feb 25, 22:28	50	dissociation		47	56	5
Feb 25, 23:28	50	dissociation		29	58	5
<i>Feb 25 subtotal</i>		6 doses, 280 mg, dose ↑ 30→50 mg				
Feb 26, 00:21	50	dissociation		45	58	5
Feb 26, 01:04	50	dissociation		36	58	5
Feb 26, 02:19	80	micro-boost	<i>plus que prévu*</i>	36	58	5
Feb 26, 02:50	50	dissociation		36	58	5
Feb 26, 03:11	50	dissociation		36	58	5
Feb 26, 03:41	50	dissociation		36	58	5
Feb 26, 04:34	50	micro-boost		36	58	5
Feb 26, 05:09	30	micro-boost		73	58	5
Feb 26, 05:53	50	micro-boost		73	58	5
Feb 26, 06:52	30	micro-boost		73	58	5
Feb 26, 12:17	30	micro-boost		46	58	5
<i>Feb 26 subtotal</i>		11 doses, 520 mg, peak 80 mg single dose				
Total	1,045	24 doses over 4 active days, mean 261 mg/day				

**plus que prévu* = “more than planned” (French). All subject notes preserved in original language.

Analysis: The log reveals a clear **dose escalation**: daily totals increased from 60 mg (Feb 22) to 520 mg (Feb 26)—an 8.7-fold increase over 4 days. Equally significant is the **intent degradation**: on Feb 22, intents were specific (“micro-boost”, “dissociation”); by Feb 23, notes shifted to “craving” (3 of 5 entries); by Feb 26, the subject noted “plus que prévu” (more than planned) at the 80 mg peak dose. The Body Battery remained at 5/100 throughout most of the observation period, indicating severe energy depletion. Biometric data was collected concurrently (stress, HR) but was *not used to modulate consumption*—the subject continued escalating despite high stress readings (73–83).

Current status: Active dependency detected by pattern analysis (dose escalation + intent degradation flagged). No structured tapering protocol has been initiated. The subject continues unsupervised use. This gap between detection and clinical action illustrates the current limitation: the platform identifies dependency patterns but does not yet translate them into supervised tapering interventions.

A.5.2 LSD (Oral, Carton)

Complete log (5 entries):

Table 13: LSD: complete timestamped log.

Timestamp	Dose	Intent	Notes	Stress	HR	BB
Feb 23, 00:46	1 carton (~100 µg)	creative		79	58	5
Feb 23, 03:33	1 carton (~100 µg)	creative		79	58	5
Feb 25, 20:12	1 carton (~100 µg)	creative	engineering session	47	56	5
Feb 26, 09:32	1 carton (~100 µg)	therapeutic		46	58	5
Feb 26, 12:17	0.1 carton (~10 µg)	microdose		46	58	5
Feb 27, 18:15	1 carton (~100 µg)	creative		—	—	—
Feb 27, 20:17	1 carton (~100 µg)	creative		—	—	—

6 full doses + 1 microdose over 5 days. Notable: 2 full doses on Feb 23 (00:46 + 03:33, 2.8h apart) concurrent with ketamine (245 mg total that night) and prazepam (40 mg). This polypharmacy pattern represents significant serotonergic risk given the concurrent SSRI/SNRI regimen. Feb 27 shows 2 additional full doses in the evening (18:15 + 20:17). No dependency pattern detected (irregular timing, consistent intent specificity).

A.5.3 THC (Vaporized)

25 entries, Feb 22–27. 1–2 chambers, 22% THC strain, 230°C vaporized. Intent: relax (12), focus (8), creative (2), sleep (1), none (2). Daily frequency: Feb 22: 1, Feb 23: 3, Feb 24: 6, Feb 25: 1, Feb 26: 9, Feb 27: 5. Peak consumption on Feb 24 (6 chambers, stress 94–98, BB declining 93→18) and Feb 26 (9 chambers).

Current status: Possible light dependency (logging only, no tapering initiated). Intent pattern shows predominant “relax” usage (55%), suggesting anxiolytic self-medication.

A.5.4 CBD (Sublingual, Tablet, Vaporized)

Doses: 1–20 mg, variable routes. **Intents logged:** anxiolysis, inflammation reduction. CBD is used as a low-risk anxiolytic tool. No dependency risk. The platform tracks its interaction with the SSRI/SNRI regimen (CBD inhibits CYP2D6, potentially increasing sertraline plasma levels—flagged by the interaction engine).

A.5.5 Nicotine (Vaporized Salt)

63 entries, Feb 22–27. 20% nicotine salt, 1–5 puffs per session, 22W. Daily frequency: Feb 22: 1, Feb 23: 7, Feb 24: 24, Feb 25: 10, Feb 26: 7, Feb 27: 14. Intent: focus (45, 71%), break (5, 8%), **craving (4, 6%)**, none (9, 14%). Peak: 24 sessions on Feb 24 (stress 94–98 sustained). Total estimated: ~170 puffs over 6 days.

Current status: Active dependency. The “craving” intent (4 entries) and empty intent fields (5 entries) indicate compulsive rather than intentional use. Feb 24 shows 23 sessions in 16 hours (one every 42 minutes on average). No tapering initiated.

A.6 Biometric–Substance Temporal Overview

Figure 3 presents an integrated view of the 6-day observation period, overlaying daily psychoactive substance events with key biometric signals. The figure reveals an inverse relationship between substance load and physiological recovery: peak consumption days (Feb 23: 27 events, Feb 26: 42 events) coincide with maximum stress (76/100), minimum Body Battery (5/100), and elevated resting HR, while lower-consumption days (Feb 22, Feb 27) show better autonomic recovery. This temporal pattern does not establish causation—the relationship is likely bidirectional (substance use ↔ stress ↔ poor recovery ↔ more substance use).




Figure 3: Six-day biometric–substance temporal overview. Psychoactive events are scaled $\times 2$ for visual comparison (actual counts: 4, 27, 48, 32, 42, 39). Body Battery oscillates between full recovery (100) and complete depletion (5), tracking inversely with substance load. Stress peaks at 76/100 on Feb 23, the highest polypharmacy day (ketamine + LSD + prazepam concurrent).

A.7 Co-Interaction Risk Assessment

The subject’s concurrent substance profile presents significant pharmacological co-interaction risks that must be explicitly evaluated. During the observation period, the following substances were used concurrently or within overlapping pharmacokinetic windows:

Table 14: Active co-interaction risks in the case study subject’s substance profile.

Substance A	Substance B	Severity	Risk
Sertraline	Ketamine	Moderate	Serotonergic potentiation
Sertraline	LSD	Moderate	Serotonergic interaction (blunting or amplification)
Venlafaxine	Ketamine	Moderate	Dual serotonergic + dissociative
Venlafaxine	LSD	Moderate	Serotonin syndrome risk
Prazepam	Ketamine	High	Additive CNS/respiratory depression
Prazepam	THC	Moderate	Additive sedation
CBD	Sertraline	Moderate	CYP2D6 inhibition $\rightarrow$ $\uparrow$ sertraline levels
Ketamine	THC	Moderate	Additive dissociation/sedation
Ketamine	Nicotine	Low	Cardiovascular stress

Critical observation: The subject was simultaneously taking two serotonergic medications (sertraline + venlafaxine) while using ketamine (up to 520 mg/day) and occasional LSD—creating a multi-layered interaction profile with compounding risks. While Mind Protocol’s interaction engine flags these pairwise, the *combinatorial* risk of 4+ concurrent serotonergic/dissociative substances exceeds what pairwise matrices can capture. No clinical evidence exists for the safety of this specific polypharmacy profile, and the interaction risk should be considered a serious limitation of the self-directed approach documented here. Clinical supervision is essential.

A.8 Pharmacological Analysis: The Night of February 22–23

Between 23:00 on February 22 and 07:00 on February 23 (8 hours), the subject consumed:

- **Ketamine:** 7 doses, 245 mg total (intranasal)
- **LSD:** 2 full doses ($\sim 200 \mu\text{g}$ total, oral)
- **Prazepam:** 4 doses, 40 mg total (sublingual)
- **Venlafaxine:** 2 doses, 150 mg total
- **THC:** 3 chambers (vaporized)
- **CBD:** 1 chamber (vaporized)
- **Nicotine:** 8 sessions, ~ 22 puffs total
- **Cyamemazine:** 2 doses, 37.5 mg total

This represents **8 psychoactive substances in 8 hours**, with the following simultaneous pharmacological actions on the CNS:

Table 15: Pharmacological profile of the Feb 22–23 polypharmacy night.

Substance	Primary target	Net effect	Risk context
Sertraline (200 mg, AM)	SERT inhibition	$\uparrow$ 5-HT	Still active ($t_{1/2}=26\text{h}$)
Venlafaxine (150 mg)	SERT + NET inhibition	$\uparrow$ 5-HT, NE	Serotonergic load
LSD (200 μg)	5-HT _{2A} agonism	$\uparrow$ 5-HT _{2A}	Triple serotonergic
Ketamine (245 mg)	NMDA antagonism	Dissociation	Respiratory risk + 5-HT
Prazepam (40 mg)	GABA-A positive mod.	CNS depression	+ ketamine = resp. risk
Cyamemazine (37.5 mg)	D ₂ /5-HT ₂ antagonism	Sedation	Additive CNS depression
THC (3 chambers)	CB ₁ agonism	Sedation/dissociation	Additive with ketamine
Nicotine (22 puffs)	nAChR agonism	Sympathetic activation	Cardiovascular stress

Critical risks identified: (1) Triple serotonergic load (sertraline + venlafaxine + LSD) creating serotonin syndrome risk; (2) Triple CNS depression (prazepam + ketamine + cyamemazine) creating respiratory depression risk; (3) Dissociative potentiation (ketamine + THC + LSD); (4) $4\times$ the prescribed prazepam daily dose in 3 hours. The Garmin stress score remained at 79 throughout this period, while Body Battery was at 5/100 (minimum), suggesting severe physiological depletion that the subject continued to override pharmacologically.

This night exemplifies the gap between the platform’s monitoring capability and clinical action. The interaction engine would flag multiple critical/high-severity pairs, but no automated intervention occurred. In a fully deployed system, this pattern should trigger an immediate clinical alert.

A.9 Temporal Pattern Analysis

Of the 181 substance events, **72%** (130 events) occurred between 22:00 and 10:00, indicating predominantly nocturnal consumption. This pattern is clinically significant: nocturnal substance use (a) disrupts sleep architecture, (b) occurs when clinical supervision is unavailable, (c) may indicate insomnia-driven self-medication, and (d) correlates with the sustained low Body Battery readings (5/100 for 80% of the observation period).

A.10 Body Pillar: Movement and Physical Practices

A.10.1 Running

Frequency: ~ 9 km/week (P57 in 30–34M cohort). **Mechanism:** Endorphin release, BDNF upregulation, cardiovascular conditioning. **Biometric signature:** HR max tracked via Garmin, post-run HRV recovery measured, Body Battery drain/recharge quantified.

Running serves as the primary endogenous tool for mood regulation. The platform records correlations between running days and next-day biometric readings (HRV and stress baselines); whether these reflect a causal dose–response relationship has not been assessed.

A.10.2 Dance as Integrated Cognitive-Motor Assessment

Mechanism: Rhythmic full-body movement recruits motor cortex, cerebellum, and basal ganglia simultaneously; engages proprioceptive feedback loops; combines cardiovascular exertion with coordination, musicality, and foreign-language cognitive load. **Biometric signature:** HR elevated to training zone, stress reduction post-session, acute HRV improvement.

Video analysis: Şımarık (Tarkan, Turkish pop, 4:03). **Method:** An automated analysis pipeline (`dance_analysis.py`) extracts 20 frames at 12s intervals via `ffmpeg`, transcribes the audio track via OpenAI Whisper (with language detection), and submits the frame sequence to a vision-language model (Claude) for structured movement analysis across the three A&B dimensions. The pipeline produces JSON and Markdown reports for each session; all three analyses referenced below were generated by this tool.

Frame-by-frame analysis of a recorded session reveals the A&B thesis in its most literal form: the body as a physical vector transmitting an awareness space proposition.

Dominant movement patterns:

- **Squat pattern (60% of frames):** Persistent wide stance with center of gravity below hip level. Quadriceps, adductors, and glutes under sustained isometric and eccentric load throughout—equivalent to a 4-minute bodyweight squat endurance protocol.
- **Dynamic lunge pattern (25%):** Forward, reverse, and lateral lunges in continuous motion. Peak amplitude at 2:12 approaches a near-split lateral position under active muscular control—not passive flexibility, but loaded mobility.
- **Hip rotation and torsion (15%):** Oblique engagement through torso-pelvis dissociation. The “şımarık” (spoiled/playful) character of the song is expressed through hip-dominant movements—the body communicates the song’s emotional register directly.

Multi-channel cognitive load: The session is performed to Turkish-language music. The subject simultaneously: (1) decodes foreign phonemes (Turkish vowel harmony, the soft “ğ”, agglutinative morphology); (2) maintains choreographic sequence memory; (3) synchronizes movement to syncopated rhythm; (4) projects emotional expression outward; (5) manages cardiovascular effort. This represents concurrent engagement of linguistic, motor, musical, proprioceptive, and expressive awareness channels.

Proprioceptive feedback: Multiple frames capture the subject touching abdomen or thigh mid-movement—self-monitoring gestures that maintain body awareness during high-intensity output. Barefoot practice on hardwood floor provides unmediated ground contact, engaging plantar proprioceptors that are suppressed by footwear.

Biometric integration: The Garmin wearable (visible in frames) records the session in real time. The same device that tracks substance tapering safety captures this dance session’s cardiovascular signature—illustrating that body optimization and consciousness practice produce data on the same biometric substrate.

Energy system: The choreography exhibits natural interval structure—high-intensity phases (deep squats, explosive transitions) alternating with relative recovery (higher stance, reduced range of motion)—producing a cardiovascular training effect without deliberate periodization. This mirrors the organic intensity modulation observed in traditional folk dance forms.

Expand Yourself (Naâman, French reggae, 4:01). The second session uses a French reggae track whose theme—personal expansion and mutual connection (“*Nous nous regardons, nous nous rayonnons*”)—invites radically different movement vocabulary. The choreography is *jump-dominant*: explosive vertical movements, wide-legged landings, and dynamic weight transfers produce moderate-to-high cardiovascular demand (contrast: Şımarık’s squat-dominant, moderate demand). Primary muscle recruitment shifts toward calves and hip flexors alongside quadriceps and glutes. The reggae off-beat requires internalization of syncopated rhythm patterns—a

different cognitive channel than Turkish agglutinative phonology, but equally demanding. Linguistically, the subject processes French lyrics containing neologisms (“Expandiose”) and reflexive constructions (“nous nous regardons”) while maintaining jump timing, spatial awareness, and musical phrase tracking. The session’s *message*: joyful embodiment of expansion, communicated through arms extended outward, grounded positions connecting to roots, and fluid transitions expressing musical flow.

Soul Plan (Naâman, French ambient reggae, 3:53). The third session inverts the energetic profile. “Soul Plan” is ambient, meditative—nearly all transcribed audio is tagged “musique d’ambiance” by Whisper, with a single spoken phrase: “*Ressentis-vous ?*” (Do you feel it?). The choreography responds with *stability-dominant* movement: wide stances, slight knee bends, controlled transitions. Cardiovascular demand is low; the primary training effect is balance and lower-body isometric strength. The cognitive state cultivated is meditative—the repetitive ambient music creates a rhythmic anchor, and the body maintains grounded positions resembling standing meditation postures. Cognitive load is minimal (French vocabulary difficulty: easy; memorization load: low), freeing attentional resources for proprioceptive monitoring—the body’s position and balance become the primary focus of attention.

Comparative A&B profile. Table 16 summarizes the three sessions across A&B dimensions. Together they illustrate, in this single case, that dance with deliberate song selection can target different cognitive-motor profiles: Şımarık maximizes cognitive load (linguistic complexity, multi-channel processing), Expand Yourself maximizes cardiovascular demand (explosive power, jump-dominant), and Soul Plan maximizes meditative proprioceptive focus. All three sessions score 7/10 on the Strong × Smart index, but via different pathways—consistent with the hypothesis that integrated cognitive-motor tasks produce measurably different biometric signatures depending on the balance of physical and cognitive demands.

Table 16: Comparative A&B profile of three dance sessions (same subject, same week).

Dimension	Şımarık (Tarkan)	Expand Yourself (Naâman)	Soul Plan (Naâman)
Genre / language	Turkish pop	French reggae	French ambient reggae
Duration	4:03	4:01	3:53
Dominant pattern	Squat 60%	Jump-dominant	Stability-dominant
Cardio demand	Moderate	Moderate-High	Low
Primary muscles	Quads, glutes, core	Quads, glutes, calves, core	Quads, hamstrings, glutes
Flexibility focus	Hip, thoracic	Hip, ankle, spine	Hip, ankle
Linguistic load	High (Turkish)	Moderate (French + neologisms)	Low (French, repetitive)
Cognitive state	Multi-channel processing (5)	Rhythmic entrainment	Meditative / proprioceptive
A&B pillar emphasis	Mind	Body	Awareness
Strong × Smart	7/10	7/10	7/10

This three-session corpus—captured within a single week—illustrates how deliberate song selection can target different cognitive-motor profiles within a single movement practice. The wearable (Garmin, visible in all sessions) records each session’s distinct physiological signature, enabling data-driven comparison of training modalities over time.

A.10.3 Yin-Yang Yoga

Mechanism: Yin (parasympathetic activation, vagal tone, flexibility) + Yang (sympathetic engagement, strength, energy). **Biometric signature:** Real-time HRV monitored during practice; stress levels tracked pre/post.

The yin-yang pairing is deliberate: yin yoga targets the autonomic imbalance identified in the WAIS-IV assessment (chronic sentinel mode = sympathetic dominance), while yang yoga builds the physical resilience that supports tapering. The platform quantifies each session’s autonomic impact.

A.10.4 Breathwork and Meditation

Mechanism: Direct ANS regulation via controlled breathing; default mode network modulation via meditation. **Biometric signature:** HRV biofeedback during practice, consistent with Garland et al. [2014].

Mind Protocol includes 30+ guided practice routines, each with biometric tracking. The PREVENT trial evidence [Kuyken et al., 2015a] for MBCT as a tapering support informed these features. The platform includes guided practices inspired by mindfulness-based approaches, though fidelity to the structured MBCT protocol used in the PREVENT trial has not been assessed.

A.11 Mind Pillar: Cognitive Enhancement Stack

Table 17: Active supplement stack (started January 31, 2026).

Supplement	Dose	Mechanism	Target
Lion’s Mane	420–1260 mg	NGF/BDNF stimulation [Saitsu et al., 2019]	Cognitive recovery
Griffonia (5-HTP)	900 mg	Serotonin precursor [Javelle et al., 2020]	Mood support
Saffron extract	1000 mg	Monoamine modulation [Hausenblas et al., 2013]	Antidepressant adjuvant
Magnesium	Standard	NMDA modulation, GABA [Boyle et al., 2017]	Sleep, stress, recovery
Omega-3 (PiLeJe)	Standard	Anti-inflammatory [Grosso et al., 2014]	Neuroprotection
Dynabiane probiotic	Standard	Gut-brain axis [Nikolova et al., 2021]	Mood, inflammation

The supplement stack targets the neurobiological substrates that will support eventual SSRI/SNRI tapering: serotonin precursors (griffonia), BDNF upregulation (Lion’s Mane), and NMDA modulation (magnesium). The 2–4 week onset of Lion’s Mane effects is tracked via cognitive task performance and biometric trends.

A.12 Data Visualizations

A.12.1 Substance Use Timeline

Figure 4 displays all 233 logged events across the 6-day observation period, revealing consumption patterns, nocturnal clustering, and escalation trajectories that are not apparent from tabular data alone.




Figure 4: Substance use timeline (Feb 22–27, 2026). Each dot represents one logged administration. Gray bands = nighttime (22:00–06:00). Note the nocturnal clustering and the escalation of ketamine events ($y=4$) from Feb 22 to Feb 26. Red dots indicate “craving” or absent intent—a dependency signal.

A.12.2 Stress Trajectory with Substance Events

Figure 5 overlays the Garmin stress readings captured at each substance log event, revealing the relationship between consumption intensity and physiological stress.




Figure 5: Garmin stress score trajectory overlaid with substance events (Feb 22–26). Stress peaked at 96–98 on Feb 24 (sustained for 12+ hours despite heavy nicotine + THC use), dropped to 19 during a sleep period, then oscillated 29–73 during the Feb 25–26 ketamine escalation. The stress score did not decrease with increasing ketamine consumption, suggesting tolerance or ineffective self-medication.

A.12.3 Dependency Score: Worked Example (Ketamine)

Figure 6 illustrates the dependency scoring algorithm applied to the subject’s actual ketamine data.




Figure 6: Dependency score computation for ketamine using actual logged data. Raw score (73/90) × risk class modifier (1.1) = 80/100 (critical). Each component is derived from the 24 timestamped entries in Table 12.

A.13 Biometric Baseline (February 2026)

Table 18: Biometric baseline from Garmin Fēnix 8 47mm AMOLED.

Metric	Value	Interpretation
Resting HR	52 bpm	Athletic range
HRV (RMSSD)	44 ms	Low-normal for age
Stress	14/100	Very low (recovery)
Body Battery	93/100	Excellent reserves
ANS Mode	Recovery	Parasympathetic dominant
Sleep duration	8.5 hr (P74)	Above median
Daily steps	6,500 (P32)	Below median
Floors climbed	14/day (P75)	Above median

The biometric profile reveals a paradox observed in this particular subject: excellent physiological recovery capacity (resting HR 52, Body Battery 93) coexisting with chronic psychological hypervigilance. The HRV of 44 ms is low-normal for a 34-year-old male—published norms for short-term RMSSD in healthy adults range from 19–75 ms [Nunan et al., 2010, Shaffer and Ginsberg, 2017]—likely reflecting the impact of the dual SSRI/SNRI regimen and ongoing substance use on autonomic flexibility. This baseline would serve as the reference point for future tapering decisions. In the proposed protocol (Appendix B), a dose reduction that drops HRV below personal baseline or elevates stress above personal threshold would trigger a taper hold—but this automated feedback loop is not yet active. Currently, biometric data is collected and displayed but not used to drive tapering decisions.

A.14 Extended Biometric Baseline (22 Days)

While the single-day snapshot above captures a moment of excellent recovery, a 22-day longitudinal view (February 15–March 8, 2026) reveals a more nuanced picture. The extended baseline was computed from daily Garmin Connect API pulls, with the latest reading per day used to avoid intra-day duplicate inflation.

Table 19: Extended biometric baseline for the primary subject over 22 days. Values are mean $\pm$ SD.

Metric	Mean	SD	Note
Resting HR (bpm)	55.9	—	Elevated vs. single-day (52)
HRV RMSSD (ms)	37.7	—	Below 44 ms baseline; declining trend
Stress (0–100)	45.6	—	Moderate-high; peaks to 98 during case study
Sleep duration (h)	8.7	—	Adequate; irregular timing
Daily steps	5,661	—	Below population median (P32)

Notably, the 22-day mean HRV (37.7 ms) is lower than the snapshot value (44 ms), suggesting that the snapshot captured a recovery peak rather than the true baseline—a common pitfall in single-point biometric reporting [Laborde et al., 2017]. The elevated mean stress (45.6/100) reflects the chronic hypervigilance already identified in the clinical assessment.

A.15 Three-Participant Biometric Comparison

To contextualize the primary subject’s biometric profile, data from two additional participants was collected over the same 22-day period (February 15–March 8, 2026) using the Mind Protocol platform. All participants wore Garmin wearable devices and transmitted data via the same API pipeline. Participants B and C provided informed consent for anonymized data inclusion.

A.15.1 Descriptive Statistics

Table 20: Three-participant biometric comparison over 22 days. Values are mean $\pm$ SD. Participant A is the primary case study subject (polymedicated, substance-tracking); B is a co-regulation partner (no psychiatric medication); C is an elite endurance athlete (no psychiatric medication). HRV was unavailable for B and C due to device model limitations (Venu 2S/3S).

ID	Profile	<i>n</i> days	HR _{rest} (bpm)	Stress (0–100)	BB _{max}	Steps
A	Polymedicated, substance-tracking	21	55.9 $\pm$ 3.3	45.6 $\pm$ 15.1	54.2 $\pm$ 35.4	5,661 $\pm$ 4,297
B	Partner, no psych. meds	22	55.4 $\pm$ 2.5	27.0 $\pm$ 11.0	77.1 $\pm$ 23.2	6,221 $\pm$ 4,233
C	Elite athlete, no psych. meds	12–17	58.2 $\pm$ 9.0	23.8 $\pm$ 23.7	—	7,872 $\pm$ 4,433

A.15.2 Between-Participant Comparisons

Given the small sample and non-normal distributions, non-parametric Mann-Whitney *U* tests were used for between-participant comparisons, with Cohen’s *d* for effect size estimation (Table 21).

Table 21: Between-participant statistical comparisons (Mann-Whitney *U*). Effect sizes: $|d| > 0.8$ = large, > 0.5 = medium.

Metric	Comparison	<i>U</i>	<i>p</i>	Sig.	Cohen’s <i>d</i>
Stress (avg)	A vs. B	404	<0.001	***	1.40
Stress (avg)	A vs. C	290	0.001	**	1.10
HR (resting)	A vs. B	259	0.50	ns	0.18
HR (resting)	A vs. C	112	0.60	ns	−0.33
Body Battery (max)	A vs. B	142	0.03	*	−0.76

A.15.3 Within-Participant Correlations

The relationship between daily stress and Body Battery was assessed using Spearman's rank correlation:

- **Participant A:** $r_s = -0.63$, $p = 0.002$ ($n = 21$)
- **Participant B:** $r_s = -0.62$, $p = 0.002$ ($n = 22$)

Both participants show a significant inverse relationship between daily stress and energy reserves, consistent with the physiological expectation that sustained sympathetic activation depletes Body Battery.

A.15.4 Interpretation

Four findings emerge from this comparison:

1. **Stress–HR decoupling:** Participant A's mean stress (45.6/100) was significantly elevated relative to both B (27.0, $d = 1.40$, $p < 0.001$) and C (23.8, $d = 1.10$, $p = 0.001$), despite *indistinguishable* resting heart rates (~ 55 – 58 bpm, $p > 0.50$ for both comparisons). This decoupling suggests that Garmin's HRV-derived stress metric captures autonomic load not reflected in cardiac output alone—consistent with the [Alvares et al. \[2016\]](#) finding that HRV variability patterns differentiate psychiatric from non-psychiatric populations independently of heart rate.
2. **Energy reserve deficit:** Participant A's Body Battery maximum (54.2 ± 35.4) was significantly lower than B's (77.1 ± 23.2 , $d = -0.76$, $p = 0.03$), with higher variability (CV = 65% vs. 30%), indicating irregular recovery patterns.
3. **Conserved stress–battery relationship:** The nearly identical Spearman correlations (A: $r_s = -0.63$; B: $r_s = -0.62$) suggest that the stress–recovery coupling is preserved across individuals despite large differences in absolute stress levels. The mechanism operates similarly; the operating point differs.
4. **HRV data gap:** HRV was available only for Participant A (Garmin Fēnix 8). Future cohort expansion should standardize on HRV-capable devices to enable cross-participant autonomic comparison.

Limitations of this comparison: This $N = 3$ comparison uses observational data without randomization, blinding, or matching. Participants differ in age, sex (not reported for B/C to preserve anonymity), medication status, lifestyle, and device model. The significant stress difference could reflect any combination of polymedication effects, substance use, psychological profile, or device calibration differences. No causal inference is possible. A properly powered comparison ($N \geq 30$, matched controls) is proposed in Section 13.

A.16 Consolidated Pilot Study Findings

The following quantitative findings emerge from the 7-day observation period and 22-day biometric baseline. These are descriptive observations from an uncontrolled N-of-1 pilot and do not support causal inference.

1. **Substance logging volume:** 236 timestamped events across 20 unique substances over 7 days (mean 33.7 events/day). Of these, approximately 75% were psychoactive substances and 25% were supplements or hydration entries.
2. **Dependency detection:** The platform's pattern analysis identified active dependency in two substances: ketamine (8.7-fold dose escalation over 4 days, intent degradation from specific to “craving”) and nicotine (63 sessions, 4 “craving” entries, compulsive 42-minute average interval on peak day). DSM-5 mapping confirmed severe nicotine SUD (7/11 criteria) and mild cannabis SUD (3/11 criteria).
3. **Biometric–substance correlation:** Peak substance consumption days (Feb 23: 27 psychoactive events; Feb 26: 42 events) coincided with maximum stress (76/100), minimum Body Battery (5/100), and elevated resting HR (58–59 bpm vs. 51 bpm baseline). The

temporal pattern suggests a bidirectional relationship: substance use $\leftrightarrow$ autonomic disturbance $\leftrightarrow$ compensatory use.

4. **HRV baseline erosion:** The 22-day mean HRV (37.7 ms RMSSD) was 14% below the single-day recovery snapshot (44 ms), consistent with chronic autonomic strain. This highlights the importance of longitudinal baselines over point measurements.
5. **Stress–HR decoupling ($N = 3$):** The primary subject’s mean stress (45.6/100) was significantly elevated relative to both controls (B: 27.0, $p < 0.001$, $d = 1.40$; C: 23.8, $p = 0.001$, $d = 1.10$), despite indistinguishable resting heart rates (~ 55 – 58 bpm, $p > 0.50$). Body Battery was also significantly lower (A: 54.2 vs. B: 77.1, $p = 0.03$, $d = -0.76$). This decoupling suggests the HRV-derived stress metric captures autonomic load beyond cardiac output.
6. **Polypharmacy risk events:** At least 3 concurrent-use episodes involved substances with known serotonergic interaction risk (SSRI + LSD, SSRI + ketamine), flagged by the interaction engine but not actioned.
7. **Predictive tracker:** A 6-signal prediction engine was implemented and validated with 30 automated tests. Accuracy data will accumulate as the system enters daily use.

These findings demonstrate the platform’s capacity for detailed substance monitoring and pattern detection. The key gap remains the translation from detection to intervention: dependency patterns were identified but no structured tapering protocol was initiated during the observation period.

A.17 Tapering Roadmap

The following is a *proposed* long-term tapering sequence, informed by the evidence reviewed in this paper. **None of these phases have been initiated at the time of writing.** The subject is currently in a pre-tapering phase (monitoring and logging only):

1. **Phase 1 (Current):** Stabilize ketamine dependency via monitored tapering. Track dose–HRV correlation to establish minimum effective dose. Target: transition from dependency to intentional tool use (PRN, not daily).
2. **Phase 2:** Nicotine tapering guided by stress biometrics and craving intent tracking. Replace with breathwork when possible (same anxiolytic effect, superior HRV profile).
3. **Phase 3:** Evaluate THC dependency vs. tool use status via biometric drift analysis. If tool: maintain as PRN. If dependency: initiate gradual taper.
4. **Phase 4:** Sertraline hyperbolic taper (per Horowitz protocol), supported by established supplement stack, MBCT practices, and continuous HRV monitoring. Minimum 6–12 months.
5. **Phase 5:** Venlafaxine taper (last due to SNRI withdrawal severity). Supported by all preceding tools and practices.
6. **Phase 6:** PRN-only medication profile. All substances available as tools, none required as maintenance. Zero dependencies. Verified by: no dose escalation, no intent degradation, no compulsive timing, stable HRV baseline.

A.18 Proposed Outcome Metrics

These metrics are proposed for longitudinal self-tracking in this individual case. Their validity as clinical endpoints has not been established.

Progress toward zero-dependency is quantified through:

- Number of substances showing dependency signals (target: 0)
- Resting HRV trajectory (target: progressive improvement toward age-appropriate norms)
- Working memory re-assessment (WAIS-IV IMT re-test at 6–8 weeks post-stabilization; target: ≥ 100)

- Intent specificity ratio: proportion of substance logs with specific functional intents vs. vague/craving intents (target: >90%)
- PRN medication frequency: trending down over months
- Body Battery and sleep quality: maintained or improved during each tapering phase

B Technical Appendix: Mind Protocol Implementation Parameters

This appendix documents the exact parameters extracted from Mind Protocol’s source code (`server.py`), enabling independent replication and audit. All values are from the production codebase as of February 2026.

B.1 Dependency Scoring Algorithm

The composite dependency score $D \in [0, 100]$ is computed as:

$$D = \min \left(100, \left(\underbrace{S_{\text{freq}}}_{\text{frequency}} + \underbrace{S_{\text{trend}}}_{\text{freq. trend}} + \underbrace{S_{\text{dose}}}_{\text{dose esc.}} + \underbrace{S_{\text{intent}}}_{\text{intent shift}} + \underbrace{S_{\text{reg}}}_{\text{regularity}} \right) \times R_{\text{class}} \right) \quad (2)$$

where:

Table 22: Dependency scoring components and ranges.

Component	Range	Description
S_{freq}	0–30	Raw frequency of substance use (logs/period)
S_{trend}	0–15	Trend direction of use frequency (increasing = higher)
S_{dose}	0–20	Dose escalation over time (dose at end vs. start)
S_{intent}	0–15	Intent degradation (specific $\rightarrow$ craving/habit)
S_{reg}	0–10	Regularity of timing (fixed intervals = higher risk)
R_{class}	0.3–1.3	Risk class modifier (supplements: 0.3; opioids: 1.3)

Maximum raw score = $90 \times 1.3 = 117$, clamped to 100. Risk thresholds: <20 = low, $20\text{--}40$ = moderate, $40\text{--}70$ = high, >70 = critical.

B.2 Pharmacokinetic Duration Model

The `PK_DURATION_MIN` map specifies minimum pharmacokinetic durations (in minutes) used to enforce safe intervals between dose changes. Values are derived from published clinical pharmacokinetic data [Rowland and Tozer, 2011, Hiemke et al., 2018, DeVane et al., 2002] and represent approximate duration of clinically relevant effects (not terminal half-lives):

Table 23: Pharmacokinetic duration parameters (selected substances).

Substance	Min	Substance	Min	Substance	Min
Nicotine	30	Caffeine	300	LSD	720
Ketamine	120	THC	240	Psilocybin	360
Alcohol	180	CBD	360	MDMA	360
Cocaine	60	Prazepam	720	Sertraline	1,440
Amphetamine	480	Venlafaxine	720	Cyamemazine	720

B.3 Drug Interaction Matrix

The `SUBSTANCE_INTERACTIONS` matrix encodes 47 pairwise drug–drug interactions with severity levels. Selected examples:

Table 24: Selected entries from the 47-interaction matrix.

Substance A	Substance B	Severity	Mechanism
Sertraline	MDMA	Critical	Serotonin syndrome risk
Sertraline	LSD	Moderate	Serotonergic interaction
SSRI	MAO inhibitor	Critical	Hypertensive crisis
Prazepam	Alcohol	High	CNS depression potentiation
Venlafaxine	Ketamine	Moderate	Serotonergic + dissociative
CBD	Sertraline	Moderate	CYP2D6 inhibition
THC	Cyamemazine	Moderate	Additive sedation
Ketamine	Alcohol	High	CNS/respiratory depression

B.4 Tapering Protocol Specifications

The `TAPERING_PROTOCOLS` module defines 21 substance-specific tapering protocols. Each specifies:

Table 25: Tapering protocol parameters by risk class.

Risk Class	Step (%)	Interval (d)	Hold trigger	Substances
Critical	5–10	7–14	Any threshold breach	BZDs, barbiturates
High	10–15	14–28	≥2 markers	SSRIs, SNRIs, APs, opioids
Moderate	15–20	7–14	≥2 markers	Gabapentin, pregabalin
Standard	20–25	3–7	≥3 markers	Caffeine, nicotine
Minimal	25–50	1–3	Not monitored	Supplements

B.5 Biometric Validation Thresholds

The `BiometricValidation` module defines per-substance deviation thresholds from personal baselines that trigger taper holds or alerts:

Table 26: Per-substance biometric withdrawal detection thresholds.

Substance	Stress Δ	HRV Δ	Sleep Δ
Sertraline	+10	-8	-10
Venlafaxine	+15	-10	-15
Prazepam	+25	-15	-20
Cyamemazine	+15	-10	-15
Ketamine	+20	-12	-15
THC	+15	-8	-10
Nicotine	+10	-5	-8

Where Δ represents the deviation from the individual's 7-day rolling baseline. Positive stress Δ = increase above baseline; negative HRV/sleep Δ = decrease below baseline. When a threshold is breached, the protocol triggers a hold (pause current taper step) and generates a clinical alert. Two consecutive breach-free intervals are required before resuming the taper.

B.6 Surveillance Symptom Categories

73 withdrawal symptoms are monitored across 8 categories:

1. **Neurological** (12): headache, dizziness, paresthesia, brain zaps, tremor, ...
2. **Gastrointestinal** (8): nausea, vomiting, diarrhea, appetite changes, ...
3. **Cardiovascular** (6): palpitations, tachycardia, blood pressure changes, ...
4. **Psychological** (14): anxiety, irritability, insomnia, vivid dreams, depersonalization, ...
5. **Sensory** (8): visual disturbances, tinnitus, hypersensitivity, ...
6. **Motor** (7): akathisia, myoclonus, gait instability, ...
7. **Cognitive** (9): concentration difficulty, memory impairment, confusion, ...
8. **Autonomic** (9): sweating, temperature dysregulation, dry mouth, ...

Symptoms are tracked via daily patient self-report and correlated with biometric data streams. The structured tracking addresses the finding by [Henssler et al. \[2024\]](#) that structured assessment detects significantly more withdrawal cases than unstructured clinical evaluation.

B.7 Pseudo-Code: Taper Step Decision

The core decision algorithm for each taper step evaluation:

```

function evaluate_taper_step(patient, substance):
    baseline = get_rolling_baseline(patient, days=7)
    current = get_current_biometrics(patient)
    thresholds = BIOMETRIC_THRESHOLDS[substance]

    stress_delta = current.stress - baseline.stress
    hrv_delta = current.hrv - baseline.hrv
    sleep_delta = current.sleep_score - baseline.sleep_score

    breaches = 0
    if stress_delta > thresholds.stress_increase: breaches += 1
    if hrv_delta < thresholds.hrv_decrease: breaches += 1
    if sleep_delta < thresholds.sleep_decrease: breaches += 1

    protocol = TAPERING_PROTOCOLS[substance]

    if breaches >= protocol.hold_trigger:
        return HOLD // pause taper, alert clinician

```

```
1700     elif dependency_score(patient, substance) < 20:
1701         return COMPLETE // taper finished
1702     else:
1703         next_dose = current_dose * (1 - protocol.step_pct)
1704         return REDUCE(next_dose, protocol.interval_days)
```

1705 C Venice Convention: Design Heuristic Framework

1706 The Venice Convention is the project’s internal set of design heuristics governing data handling,
1707 interaction style, and architectural trade-offs. It is presented here as contextual documentation
1708 of engineering choices—not as empirically validated principles. The framework draws on value-
1709 sensitive design methodology [Friedman et al., 2013].

1710 C.1 Dual-Layer Architecture Rationale

1711 Each architectural decision has an *experiential* dimension (user autonomy, privacy, sense of
1712 agency) and a *substrate* dimension (data structures, protocols, and enforcement mechanisms).
1713 For example, biometric co-regulation is simultaneously a user-facing feature for relational sup-
1714 port and a Pearson-correlation computation over time-aligned stress timeseries in append-only
1715 JSONL files. The convention requires that neither dimension be sacrificed to the other.

1716 C.2 Epistemic Scope Constraints

1717 The system’s AI components operate under explicit epistemic boundaries: reporting observable
1718 behavioral patterns (e.g., “dose escalation detected over 4 days”) rather than claiming causal
1719 insight into the user’s psychological state. This design constraint limits the system to non-
1720 directive information display, consistent with the FDA’s CDS non-device pathway [U.S. Food
1721 and Drug Administration, 2023].

1722 C.3 Art. 9 Special Category Firewall

1723 A deterministic post-filter screens extracted insights against GDPR Art. 9 keyword categories
1724 (health conditions, political affiliation, sexual orientation). Matched insights are dropped before
1725 persistence. Raw conversation transcripts are never stored; persisted memory uses paraphrased
1726 summaries under 150 characters. Ephemeral `tempfile.mkdtemp` storage ensures intermediate
1727 processing artifacts do not survive beyond the current session.