
# Drug Misuse Exploration

003 Team Exploratory // Datafest 2021 //

- Analyzed by
  Andre Williams
  - Rebecca Martin
  - Cindy Luna Miranda


### **Problem Statement**

Patients with a history of <u>illegal drug use</u>, <u>chronic pain</u>, <u>anxiety</u>, <u>tobacco use</u>, and <u>cannabis use</u> are more likely to misuse prescribed medication.




□ Variability Summary for OP_NMU				
	Mean	Std Dev	Observations	
OP_NMU	0.111104	0.314266	29873	
Illegal_USE[No]	0.064957	0.246455	24293	
Illegal_USE[Yes]	0.312007	0.463354	5580	

The mean jumps 5x for opioid misuse if have used illicit drugs.



Approximately 12k people die each year in the U.S. from prescribed opioids alone, according to CDC graph (above).

## Predicting Future Misuse



Parameter Estimates						
Term	Estimate	Std Error	ChiSquare	Prob>ChiSq		
Intercept	1.41350251	0.0395218	1279.1	<.0001		
lllegal_USE[No]	0.67934596	0.0233395	847.22	<.0001		
CAN_USE[0]	0.16481463	0.0245915	44.92	<.0001		
PAIN_CHRONICYR[0]	0.13015619	0.020739	39.39	<.0001		
MENT_ANYANX[0]	0.26901635	0.0237361	128.45	<.0001		
TOBACCO_FREQ[Don't know]	0.12121738	0.0793437	2.33	0.1266		
TOBACCO_FREQ[Everyday]	-0.1817144	0.0414859	19.19	<.0001		
TOBACCO_FREQ[Not at all]	0.58686063	0.0367199	255.43	<.0001		
MENT_DEP[0]	-0.0142221	0.0326904	0.19	0.6635		

When running a logistic model, we found the following predictors to be significant when determining future prescribed drug misuse.

- History of illegal drug use
- Cannabis use
- Chronic Pain
- Mental Anxiety
- Tobacco use

#### Recommendations:

- Alternative medication to prescribe for chronic pain or those suffering from anxiety
- Observing those prescribed opioids for trends in misuse