

An Analysis of Demographic Patterns and Frequent Combinations of Drug Misuse in Canada

Team Name: *Bad Boys*

Team Members: *Haoluan Chen, Dawei Dong, Yujie Li, Tu Wu*

Table of
Contents

Thank You!

Department of Statistical Science,
University of Toronto

A diagram with a large light blue circle in the center. Inside this circle, the text "Table of Contents" is written in a large, bold, black font. Surrounding the central circle are three smaller, light gray circles. One is at the top, one at the bottom left, and one at the bottom right. Each of these smaller circles contains text in a bold, black font. The top circle contains "Introduction". The bottom left circle contains "Methods & Results". The bottom right circle contains "Conclusion & Discussion". The entire diagram is set against a dark blue background.

Table of Contents

Introduction

**Methods
&
Results**

**Conclusion
&
Discussion**

Introduction



Background

**Research
Question**

Background

- Prescription drug misuse is a serious problem
- Data collected from Survey of Non-Medical Use of Prescription Drugs (NMURx) Program
- Variables dictionary
 - Survey administration variables
 - Demographics of respondent (i.e. age, gender, income, residence, etc.)
 - Prescription Pain Relievers
 - Prescription and Non-Prescription Codeine
 - Other Prescription Drugs- (i.e. benzodiazepines, stimulants, etc.)
 - End of Survey Questions- (i.e. other risk factors, illicit drug use, etc.)
 - DAST-10 Questionnaire
 - Summary Variables- derived variables from dataset to aid in analysis

Research Question

1. To explore the **demographic** and **behavioral** pattern of general and specific drugs misuse
2. To investigate the **combinations** of drugs that tend to be misused together

Methods



**Generalized
Linear
Mixed
Model**

**Generalized
Linear
Model**

**Descriptive
Statistics**

Generalized Linear Mixed Model

Response Variable: Misuse or not (0 or 1)

Predictors: Demographic Variables and Behavioral Variables.

**Fixed
Effect**

**Random
Effect**


Fixed Effect

Table 1: The Summary Table of Coefficients

Coefficient Parameter	Estimated	95% Credible Interval	
	Median odds ratio	Lower Bound (0.025 quantile)	Upper Bound (0.975 quantile)
(Intercept)	1.039	0.779	1.385
Age	0.984	0.982	0.986
Sex (Female)	0.795	0.743	0.852
Income	0.954	0.941	0.967
Crack Cocaine (In last week)	2.876	1.984	4.168
Crack Cocaine (In last 30 days)	3.698	2.498	5.473
Crack Cocaine (In last 12 months)	3.811	2.475	5.868
Crack Cocaine (Yes, during my lifetime)	1.739	1.475	2.050
Tobacco Use (Occasionally)	1.091	0.959	1.240
Tobacco Use (Not at all)	0.780	0.715	0.851
Alcohol Use (4 to 5 times a week)	0.908	0.782	1.054
Alcohol Use (2 to 3 times a week)	0.889	0.782	1.010
Alcohol Use (Once a week)	0.841	0.731	0.967
Alcohol Use (2 to 3 times a month)	0.744	0.648	0.855
Alcohol Use (Once a month)	0.760	0.644	0.898
Alcohol Use (Less than once a month)	0.634	0.554	0.724
Alcohol Use (Never)	0.712	0.621	0.815
DAST_3	0.607	0.563	0.656
DAST_5	1.365	1.230	1.514
DAST_8	1.788	1.551	2.062

DAST_3 : Are you able to stop using drugs when you want to? (yes)
DAST_5: Do you ever feel bad or guilty about your drug use? (yes)
DAST_8: Have you engaged in illegal activities in order to obtain drugs? (yes)

Random Effect



Generalized Linear Model

1. Prescription Codeine product

+ effect:

- crack use(within last week),
- tobacco/alcohol use frequency,
- engaged in illegal activities to obtain drugs

- effect:

- age,
- gender(female),
- higher income level,
- able to stop using drugs if want to

2. Non-prescription Codeine product

+ effect:

- crack use(within last week),
- tobacco/alcohol use frequency,
- engaged in illegal activities to obtain drugs

- effect:

- age,
- gender(female),
- higher income level,
- able to stop using drugs if want to

3. THC

+ effect:

- alcohol use frequency,
- engaged in illegal activities to obtain drugs
- ever feel guilty

- effect:

- age,
- higher education

4. Stimulant

+ effect:


- ever feel guilty

- effect:

- age

Descriptive Statistics


Heatmap of proportion of misuse



**Top-8
combination**

Top-8 Combination

CH - Methadone and Tapentadol
FN - Oxymorphone and Stimulants
HM - Tapentadol and Sufentanil
FM - Oxymorphone and Sufentanil
BN - Buprenorphine and Stimulants
BH - Buprenorphine and Tapentadol
FH - Oxymorphone and Tapentadol
CM - Methadone and Sufentanil



Discussion

Application:

- Help in regulations or physician questionnaires

Future work:

- Demographics for multiple drug misuse
- Add weight parameter if time permitted



Thank You For Watching!

