

# **Solving Non-Linear Real Arithmetic Formulas with Virtual Substitution**

Author: Aklima Zaman  
Supervision: Erika Ábrahám

Theory of Hybrid Systems - Informatik 2 - RWTH-Aachen

Satisfiability Checking Seminar, Winter-16/17


# Outline

- Motivation
- Real Arithmetic Formula
- Virtual Substitution
  - Sign Invariant Regions
  - Compute Zeros
  - Compute Test Candidates
  - Virtual Substitution Rules

# Motivation

- Other related methods
  - interval constraint propagation
  - cylindrical algebraic decomposition
- Virtual substitution
  - applicable only to sub-language
  - eliminates quantified variables up to degree 4

# Flow Chart of Virtual Substitution



## Real Arithmetic Formula

- Real arithmetic (RA) formula has the following syntax:

**terms:**  $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$

**constraints:**  $c := t < t$

**formulas:**  $\varphi := c \mid \neg\varphi \mid \varphi \wedge \varphi \mid \exists x \cdot \varphi$

## Real Arithmetic Formula

- Real arithmetic (RA) formula has the following syntax:

**terms:**  $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$

**constraints:**  $c := t < t$

**formulas:**  $\varphi := c \mid \neg \varphi \mid \varphi \wedge \varphi \mid \exists x \cdot \varphi$

## Real Arithmetic Formula

- Real arithmetic (RA) formula has the following syntax:

**terms:**  $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$

**constraints:**  $c := t < t$

**formulas:**  $\varphi := c \mid \neg \varphi \mid \varphi \wedge \varphi \mid \exists x. \varphi$

## Real Arithmetic Formula

- Real arithmetic (RA) formula has the following syntax:

**terms:**  $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$

**constraints:**  $c := t < t$

**formulas:**  $\varphi := c \mid \neg\varphi \mid \varphi \wedge \varphi \mid \exists x. \varphi$



## Real Arithmetic Formula

- Real arithmetic (RA) formula has the following syntax:

**terms:**  $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$

**constraints:**  $c := t < t$

**formulas:**  $\varphi := c \mid \neg\varphi \mid \varphi \wedge \varphi \mid \exists x \cdot \varphi$

- Polynomial  $p(x) \in \mathbb{Z}[x_1 \dots, x_n][x]$  normal form:

$$p(x) = a_d x^d + a_{d-1} x^{d-1} + \dots + a_0 x^0$$

Example:

$$\varphi = \underbrace{((x^2 + 2x + 4z))}_{p_1} \leq 0 \vee \underbrace{(yx^2 + 6y^3x + 4z)}_{p_2} = 0$$

## Real Arithmetic Formula

- Real arithmetic (RA) formula has the following syntax:

**terms:**  $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$

**constraints:**  $c := t < t$

**formulas:**  $\varphi := c \mid \neg \varphi \mid \varphi \wedge \varphi \mid \exists x \cdot \varphi$

- Polynomial  $p(x) \in \mathbb{Z}[x_1 \dots, x_n][x]$  normal form:

$$p(x) = a_d x^d + a_{d-1} x^{d-1} + \dots + a_0 x^0$$

Example:

$$\varphi = \underbrace{((x^2 + 2x + 4z) \leq 0)}_{p_1} \vee \underbrace{(yx^2 + 6y^3x + 4z = 0)}_{p_2}$$

## Real Arithmetic Formula

- Real arithmetic (RA) formula has the following syntax:

**terms:**  $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$

**constraints:**  $c := t < t$

**formulas:**  $\varphi := c \mid \neg \varphi \mid \varphi \wedge \varphi \mid \exists x \cdot \varphi$

- Polynomial  $p(x) \in \mathbb{Z}[x_1 \dots, x_n][x]$  normal form:

$$p(x) = a_d x^d + a_{d-1} x^{d-1} + \dots + a_0 x^0$$

Example:

$$\varphi = \underbrace{((x^2 + 2x + 4z))}_{p_1} \leq 0 \vee \underbrace{(yx^2 + 6y^3x + 4z)}_{p_2} = 0$$

## Real Arithmetic Formula

- Real arithmetic (RA) formula has the following syntax:

**terms:**  $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$

**constraints:**  $c := t < t$

**formulas:**  $\varphi := c \mid \neg \varphi \mid \varphi \wedge \varphi \mid \exists x \cdot \varphi$


- Polynomial  $p(x) \in \mathbb{Z}[x_1 \dots, x_n][x]$  normal form:

$$p(x) = a_d x^d + a_{d-1} x^{d-1} + \dots + a_0 x^0$$

Example:

$$\varphi = \underbrace{((x^2 + 2x + 4z) \leq 0)}_{p_1} \vee \underbrace{(yx^2 + 6y^3x + 4z) = 0)}_{p_2}$$

# Flow Chart of Virtual Substitution



## Virtual Substitution

- Quantifier elimination procedure:


$$\exists x_1 \dots \exists x_n \cdot \varphi \equiv \exists x_1 \dots \exists x_{n-1} \cdot \psi$$

where  $\varphi, \psi$  quantifier free.

- Quantifier elimination by virtual substitution:


$$\exists x_1 \dots \exists x_n \cdot \varphi \equiv \exists x_1 \dots \exists x_{n-1} \cdot \bigvee_{t \in T} (\varphi[t//x] \wedge S_t)$$

## Sign Invariant Regions



## Compute Zeros

$$p(x) = ax^2 + bx + c \quad \text{side condition: } a = 0 \wedge b = 0$$




constant polynomial  $\Rightarrow$  constant zero or non zeros



## Compute Zeros


$$p(x) = ax^2 + bx + c \quad \text{side condition: } a = 0 \wedge b \neq 0$$



$$\xi = -b/c$$

## Compute Zeros




$$p(x) = ax^2 + bx + c \quad \text{side condition: } a \neq 0 \wedge b^2 - 4ac \geq 0$$



$$\xi_1 = \frac{-b - \sqrt{b^2 - 4ac}}{2a}, \quad \xi_2 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$

# Compute Test Candidates

Possible solution intervals for  $x$  on  $p \sim 0$ :



Constraints	$-\infty$	$x_1$	$x_1 + \epsilon$	$x_2$	$x_2 + \epsilon$	$x_3$	$x_3 + \epsilon$
$p = 0$	-	✓	-	✓	-	✓	-
$p > 0, p < 0, p \neq 0$	✓	-	✓	-	✓	-	✓
$p \geq 0, p \leq 0$	✓	✓	-	✓	-	✓	-


## Example

$$\varphi := ((\underbrace{xy - 1}_{p_1} = 0) \wedge \underbrace{y^2 - 1}_{p_2} < 0)$$

Elimination of  $y$ :

	constraints	test candidates
1.	from all constraints	$-\infty$
2.	$p_1 = 0$	$1/x$ if $x \neq 0$
3.	$p_2 < 0$	$1 + \epsilon$
4.	$p_2 < 0$	$-1 + \epsilon$

# Example



## Example

$$\varphi := ((\underbrace{xy - 1}_{p_1} = 0) \wedge \underbrace{y^2 - 1}_{p_2} < 0)$$

Elimination of  $y$ :

	constraints	test candidates
1.	from all constraints	$-\infty$
2.	$p_1 = 0$	$1/x$ if $x \neq 0$
3.	$p_2 < 0$	$1 + \epsilon$
4.	$p_2 < 0$	$-1 + \epsilon$

$$\begin{aligned} \exists x \cdot \exists y \cdot \varphi &\leftrightarrow \exists x \cdot \begin{aligned} &(\varphi[-\infty/y]) && \vee \\ &(\varphi[\frac{1}{x}/y] \quad \wedge x \neq 0) && \vee \\ &(\varphi[1 + \epsilon/y]) && \vee \\ &(\varphi[-1 + \epsilon/y]) \end{aligned} \end{aligned}$$

## Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \quad \text{with } u, q, r, s \text{ polynomials.}$$

## Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \quad \text{with } u, q, r, s \text{ polynomials.}$$

Assume,  $p(x) = 0$  and test candidate is  $\frac{u+q\sqrt{r}}{s}$



## Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \text{ with } u, q, r, s \text{ polynomials.}$$

Assume,  $p(x) = 0$  and test candidate is  $\frac{u+q\sqrt{r}}{s}$

①  $(p(x) = 0)[\frac{u+q\sqrt{r}}{s} // x]$  to be computed.

## Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \text{ with } u, q, r, s \text{ polynomials.}$$

Assume,  $p(x) = 0$  and test candidate is  $\frac{u+q\sqrt{r}}{s}$

- 1  $(p(x) = 0)[\frac{u+q\sqrt{r}}{s} // x]$  to be computed.
- 2 Transform the result to  $\frac{u' + q'\sqrt{r}}{s'} = 0$  where  $u', q', s'$  are polynomials.

## Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \text{ with } u, q, r, s \text{ polynomials.}$$

Assume,  $p(x) = 0$  and test candidate is  $\frac{u+q\sqrt{r}}{s}$

- 1  $(p(x) = 0)[\frac{u+q\sqrt{r}}{s} // x]$  to be computed.
- 2 Transform the result to  $\frac{u' + q'\sqrt{r}}{s'} = 0$  where  $u', q', s'$  are polynomials.
- 3  $\frac{u' + q'\sqrt{r}}{s'} = 0$

## Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \quad \text{with } u, q, r, s \text{ polynomials.}$$

Assume,  $p(x) = 0$  and test candidate is  $\frac{u+q\sqrt{r}}{s}$

- ①  $(p(x) = 0)[\frac{u+q\sqrt{r}}{s} // x]$  to be computed.
- ② Transform the result to  $\frac{u' + q'\sqrt{r}}{s'} = 0$  where  $u', q', s'$  are polynomials.
- ③  $\frac{u' + q'\sqrt{r}}{s'} = 0$   
 $\iff u' + q'\sqrt{r} = 0$

## Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \quad \text{with } u, q, r, s \text{ polynomials.}$$

Assume,  $p(x) = 0$  and test candidate is  $\frac{u+q\sqrt{r}}{s}$

- ①  $(p(x) = 0)[\frac{u+q\sqrt{r}}{s} // x]$  to be computed.
- ② Transform the result to  $\frac{u' + q'\sqrt{r}}{s'} = 0$  where  $u', q', s'$  are polynomials.
- ③  $\frac{u' + q'\sqrt{r}}{s'} = 0$   
 $\iff u' + q'\sqrt{r} = 0$   
 $\iff u'q' \leq 0 \wedge |u'| = |q'\sqrt{r}|$

## Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \quad \text{with } u, q, r, s \text{ polynomials.}$$

Assume,  $p(x) = 0$  and test candidate is  $\frac{u+q\sqrt{r}}{s}$

- ①  $(p(x) = 0)[\frac{u+q\sqrt{r}}{s} // x]$  to be computed.
- ② Transform the result to  $\frac{u' + q'\sqrt{r}}{s'} = 0$  where  $u', q', s'$  are polynomials.
- ③  $\frac{u' + q'\sqrt{r}}{s'} = 0$   
 $\iff u' + q'\sqrt{r} = 0$   
 $\iff u'q' \leq 0 \wedge |u'| = |q'\sqrt{r}|$   
 $\iff u'q' \leq 0 \wedge u'^2 - q'^2\sqrt{r} = 0$


## Substitution of Infinitesimal Expressions

Assume  $p(x) < 0$  and test candidate is  $e + \epsilon$

## Substitution of Infinitesimal Expressions

Assume  $p(x) < 0$  and test candidate is  $e + \epsilon$

$$\begin{aligned}
 (p < 0)[e + \epsilon/x] &= \underbrace{((p < 0)[e/x])}_{\text{Case 1}} && \vee \\
 &\quad \underbrace{((p = 0)[e/x] \wedge (p' < 0)[e/x])}_{\text{Case 2}} && \vee \\
 &\quad \underbrace{((p = 0)[e/x] \wedge (p' = 0)[e/x] \wedge (p'' < 0)[e/x])}_{\text{Case 3}} && \vee
 \end{aligned}$$






## Substitution of Infinitesimal Expressions

Assume  $p(x) < 0$  and test candidate is  $e + \epsilon$


$$\begin{aligned}
 (p < 0)[e + \epsilon/x] &= \underbrace{((p < 0)[e/x])}_{\text{Case 1}} && \vee \\
 &\quad \underbrace{((p = 0)[e/x] \wedge (p' < 0)[e/x])}_{\text{Case 2}} && \vee \\
 &\quad \underbrace{((p = 0)[e/x] \wedge (p' = 0)[e/x] \wedge (p'' < 0)[e/x])}_{\text{Case 3}} && \vee
 \end{aligned}$$



## Substitution of Infinitesimal Expressions

Assume  $p(x) < 0$  and test candidate is  $e + \epsilon$

$$\begin{aligned}
 (p < 0)[e + \epsilon/x] &= \underbrace{((p < 0)[e/x])}_{\text{Case 1}} && \vee \\
 &\underbrace{((p = 0)[e/x] \wedge (p' < 0)[e/x])}_{\text{Case 2}} && \vee \\
 &\underbrace{((p = 0)[e/x] \wedge (p' = 0)[e/x] \wedge (p'' < 0)[e/x])}_{\text{Case 3}} && \vee
 \end{aligned}$$



## Substitution of a Minus Infinity

Assume  $(p(x) = ax^2 + bx + c) < 0$  and test candidate is  $-\infty$

$$p(x) < 0[-\infty/x] = \underbrace{a < 0}_{\text{Case 1}} \quad \vee$$


Case 1

$$\underbrace{a = 0 \wedge b > 0}_{\text{Case 2}} \quad \vee$$

Case 2

$$\underbrace{a = 0 \wedge b = 0 \wedge c < 0}_{\text{Case 3}}$$

Case 3



## Substitution of a Minus Infinity

Assume  $(p(x) = ax^2 + bx + c) < 0$  and test candidate is  $-\infty$

$$p(x) < 0[-\infty/x] = \underbrace{a < 0}_{\text{Case 1}} \quad \vee$$


Case 1

$$\underbrace{a = 0 \wedge b > 0}_{\text{Case 2}} \quad \vee$$

Case 2

$$\underbrace{a = 0 \wedge b = 0 \wedge c < 0}_{\text{Case 3}}$$

Case 3




## Substitution of a Minus Infinity

Assume  $(p(x) = ax^2 + bx + c) < 0$  and test candidate is  $-\infty$

$$p(x) < 0[-\infty/x] = \underbrace{a < 0}_{\text{Case 1}} \quad \vee$$

$$\underbrace{a = 0 \wedge b > 0}_{\text{Case 2}} \quad \vee$$

$$\underbrace{a = 0 \wedge b = 0 \wedge c < 0}_{\text{Case 3}}$$



## Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

Elimination of  $y$ :

1. Test candidate:  $-\infty$

$$\begin{aligned} \exists x \cdot \quad & ( \quad (xy - 1 = 0)[- \infty / y] \\ & \wedge \quad (y^2 - 1 < 0)[- \infty / y] ) \end{aligned}$$

## Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

Elimination of  $y$ :

1. Test candidate:  $-\infty$

$$\begin{aligned} \exists x \cdot \quad & ( \quad (xy - 1 = 0)[- \infty / y] \\ & \wedge \quad (y^2 - 1 < 0)[- \infty / y] ) \end{aligned}$$

$$\Leftrightarrow \exists x \cdot \quad ( \quad (x = 0 \wedge -1 = 0)$$

## Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

Elimination of  $y$ :

1. Test candidate:  $-\infty$

$$\begin{aligned} \exists x \cdot & \quad ( \quad (xy - 1 = 0)[- \infty / y] \\ & \quad \wedge \quad (y^2 - 1 < 0)[- \infty / y] ) \end{aligned}$$

$$\begin{aligned} \Leftrightarrow \exists x \cdot & \quad ( \quad (x = 0 \wedge -1 = 0) \\ & \quad \wedge \quad (1 < 0 \vee (1 = 0 \wedge 0 > 0) \vee (1 = 0 \wedge 0 = 0 \wedge -1 < 0))) \end{aligned}$$



## Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

Elimination of  $y$ :

1. Test candidate:  $-\infty$

$$\begin{aligned} \exists x \cdot & \quad ( \quad (xy - 1 = 0)[- \infty / y] \\ & \quad \wedge \quad (y^2 - 1 < 0)[- \infty / y] ) \end{aligned}$$

$$\begin{aligned} \Leftrightarrow \exists x \cdot & \quad ( \quad (x = 0 \wedge -1 = 0) \\ & \quad \wedge \quad (1 < 0 \vee (1 = 0 \wedge 0 > 0) \vee (1 = 0 \wedge 0 = 0 \wedge -1 < 0))) \end{aligned}$$

$$\Leftrightarrow \exists x \cdot \quad (\mathbf{false})$$

## Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

Elimination of  $y$ :

2. Test candidate:  $\frac{1}{x}$  if  $x \neq 0$

$$\begin{aligned} \exists x \cdot & \quad ( \quad (xy - 1 = 0)[\frac{1}{x}/y] \\ & \quad \wedge \quad (y^2 - 1 < 0)[\frac{1}{x}/y] \\ & \quad \wedge \quad x \neq 0 ) \end{aligned}$$

## Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

Elimination of  $y$ :

2. Test candidate:  $\frac{1}{x}$  if  $x \neq 0$

$$\begin{aligned} \exists x \cdot & \left( (xy - 1 = 0) \left[ \frac{1}{x}/y \right] \right. \\ & \wedge (y^2 - 1 < 0) \left[ \frac{1}{x}/y \right] \\ & \left. \wedge x \neq 0 \right) \end{aligned}$$

$$\Leftrightarrow \exists x \cdot \left( (0 = 0) \right)$$

## Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

Elimination of  $y$ :

2. Test candidate:  $\frac{1}{x}$  if  $x \neq 0$

$$\begin{aligned} \exists x \cdot & \left( (xy - 1 = 0)[\tfrac{1}{x}/y] \right. \\ & \wedge (y^2 - 1 < 0)[\tfrac{1}{x}/y] \\ & \left. \wedge x \neq 0 \right) \\ \Leftrightarrow \exists x \cdot & \left( (0 = 0) \right. \\ & \left. \wedge ((1 > 0) \wedge 1 - x^2 < 0 \vee (1 < 0 \wedge 1 - x^2 < 0)) \right) \end{aligned}$$

## Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

Elimination of  $y$ :

2. Test candidate:  $\frac{1}{x}$  if  $x \neq 0$

$$\begin{aligned} \exists x \cdot & \quad ( \quad (xy - 1 = 0)[\frac{1}{x}/y] \\ & \quad \wedge \quad (y^2 - 1 < 0)[\frac{1}{x}/y] \\ & \quad \wedge \quad x \neq 0 ) \end{aligned}$$

$$\begin{aligned} \Leftrightarrow \exists x \cdot & \quad ( \quad (0 = 0) \\ & \quad \wedge \quad ((1 > 0) \wedge 1 - x^2 < 0 \vee (1 < 0 \wedge 1 - x^2 < 0)) \\ & \quad \wedge \quad x \neq 0 ) \end{aligned}$$

## Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

Elimination of  $y$ :

2. Test candidate:  $\frac{1}{x}$  if  $x \neq 0$

$$\begin{aligned} \exists x \cdot & \left( (xy - 1 = 0) \left[ \frac{1}{x}/y \right] \right. \\ & \wedge (y^2 - 1 < 0) \left[ \frac{1}{x}/y \right] \\ & \left. \wedge x \neq 0 \right) \end{aligned}$$

$$\begin{aligned} \Leftrightarrow \exists x \cdot & \left( (0 = 0) \right. \\ & \wedge ((1 > 0) \wedge 1 - x^2 < 0 \vee (1 < 0 \wedge 1 - x^2 < 0)) \\ & \left. \wedge x \neq 0 \right) \end{aligned}$$

$$\begin{aligned} \Leftrightarrow \exists x \cdot & \left( (1 - x^2 < 0) \right. \\ & \left. \wedge x \neq 0 \right) \end{aligned}$$

## Example

$$\exists x. (1 - x^2 < 0 \wedge x \neq 0)$$

Elimination of  $x$ :

1. Test candidate:  $-\infty$

$$(1 - x_2 < 0)[- \infty / x]$$

$$= (-1 < 0 \vee (-1 = 0 \wedge 0 > 0)) \vee (-1 = 0 \wedge 0 = 0 \wedge 1 < 0))$$

$$= \text{true}$$

## Example

$$\exists x. ( \text{true} \wedge x \neq 0 )$$

Elimination of  $x$ :

1. Test candidate:  $-\infty$

$$\begin{aligned} & (x \neq 0)[- \infty / x] \\ &= (1 \neq 0 \vee 0 \neq 0) \\ &= \text{true} \end{aligned}$$



## Example


$$\exists x. ( \text{true} \wedge \text{true} )$$

Elimination of  $x$ :

1. Test candidate:  $-\infty$

$$\begin{aligned} & (x \neq 0)[-\infty/x] \\ = & (1 \neq 0 \vee 0 \neq 0) \\ = & \text{true} \end{aligned}$$

# Flow Chart of Virtual Substitution



## Example: Search Tree

$$\varphi := ((\underbrace{xy - 1}_{p_1} = 0) \wedge \underbrace{y^2 - 1}_{p_2} < 0)$$

