

Solving Non-Linear Real Arithmetic Formulas with Virtual Substitution

Author: Aklima Zaman
Supervision: Erika Ábrahám

Theory of Hybrid Systems - Informatik 2 - RWTH-Aachen

Satisfiability Checking Seminar, Winter-16/17


Outline

- Motivation
- Real Arithmetic Formula
- Virtual Substitution
 - Sign Invariant Regions
 - Compute Zeros
 - Compute Test Candidates
 - Virtual Substitution Rules

Motivation

- Other related methods
 - interval constraint propagation
 - cylindrical algebraic decomposition
- Virtual substitution
 - applicable only to sub-language
 - eliminates quantified variables up to degree 4

Flow Chart of Virtual Substitution



Real Arithmetic Formula

- Real arithmetic (RA) formula has the following syntax:
polynomials: $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$
constraints: $c := t < t$
formulas: $\varphi := c \mid \neg \varphi \mid \varphi \wedge \varphi \mid \exists x. \varphi$

Real Arithmetic Formula

- Real arithmetic (RA) formula has the following syntax:

polynomials: $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$

constraints: $c := t < t$

formulas: $\varphi := c \mid \neg \varphi \mid \varphi \wedge \varphi \mid \exists x. \varphi$

Real Arithmetic Formula

- Real arithmetic (RA) formula has the following syntax:
polynomials: $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$
constraints: $c := t < t$
formulas: $\varphi := c \mid \neg \varphi \mid \varphi \wedge \varphi \mid \exists x \cdot \varphi$

Real Arithmetic Formula

- Real arithmetic (RA) formula has the following syntax:
polynomials: $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$
constraints: $c := t < t$
formulas: $\varphi := c \mid \neg \varphi \mid \varphi \wedge \varphi \mid \exists x. \varphi$

Real Arithmetic Formula

- Real arithmetic (RA) formula has the following syntax:
polynomials: $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$
constraints: $c := t < t$
formulas: $\varphi := c \mid \neg \varphi \mid \varphi \wedge \varphi \mid \exists x \cdot \varphi$
- Polynomial $p(x) \in \mathbb{Z}[x_1 \dots, x_n][x]$ normal form:

$$p(x) = a_d x^d + a_{d-1} x^{d-1} + \dots + a_0 x^0$$

Example:

$$\varphi = \underbrace{((x^2 + 2x + 4z) \leq 0)}_{p_1} \vee \underbrace{(yx^2 + 6y^3x + 4z) = 0)}_{p_2}$$

Real Arithmetic Formula

- Real arithmetic (RA) formula has the following syntax:
polynomials: $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$
constraints: $c := t < t$
formulas: $\varphi := c \mid \neg \varphi \mid \varphi \wedge \varphi \mid \exists x \cdot \varphi$
- Polynomial $p(x) \in \mathbb{Z}[x_1 \dots, x_n][x]$ normal form:

$$p(x) = a_d x^d + a_{d-1} x^{d-1} + \dots + a_0 x^0$$

Example:

$$\varphi = \underbrace{((x^2 + 2x + 4z) \leq 0)}_{p_1} \vee \underbrace{(yx^2 + 6y^3x + 4z) = 0)}_{p_2}$$

Real Arithmetic Formula

- Real arithmetic (RA) formula has the following syntax:
polynomials: $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$
constraints: $c := t < t$
formulas: $\varphi := c \mid \neg \varphi \mid \varphi \wedge \varphi \mid \exists x. \varphi$
- Polynomial $p(x) \in \mathbb{Z}[x_1 \dots, x_n][x]$ normal form:

$$p(x) = a_d x^d + a_{d-1} x^{d-1} + \dots + a_0 x^0$$

Example:

$$\varphi = \underbrace{((x^2 + 2x + 4z))}_{p_1} \leq 0 \vee \underbrace{(yx^2 + 6y^3x + 4z)}_{p_2} = 0$$

Real Arithmetic Formula


- Real arithmetic (RA) formula has the following syntax:
polynomials: $t := 0 \mid 1 \mid x \mid t + t \mid t - t \mid t \cdot t$
constraints: $c := t < t$
formulas: $\varphi := c \mid \neg \varphi \mid \varphi \wedge \varphi \mid \exists x \cdot \varphi$
- Polynomial $p(x) \in \mathbb{Z}[x_1 \dots, x_n][x]$ normal form:

$$p(x) = a_d x^d + a_{d-1} x^{d-1} + \dots + a_0 x^0$$

Example:

$$\varphi = \underbrace{((x^2 + 2x + 4z) \leq 0)}_{p_1} \vee \underbrace{(yx^2 + 6y^3x + 4z) = 0)}_{p_2}$$

Flow Chart of Virtual Substitution



Virtual Substitution

- Quantifier elimination procedure:


$$\exists x_1 \dots \exists x_n \cdot \varphi \equiv \exists x_1 \dots \exists x_{n-1} \cdot \psi$$

where φ, ψ quantifier free.

- Quantifier elimination by virtual substitution:


$$\exists x_1 \dots \exists x_n \cdot \varphi \equiv \exists x_1 \dots \exists x_{n-1} \cdot \bigvee_{t \in T} (\varphi[t//x] \wedge S_t)$$

Sign Invariant Regions



Compute Zeros


$$p(x) = ax^2 + bx + c \quad \text{side condition: } a = 0 \wedge b = 0$$



constant polynomial \Rightarrow constant zero or non zeros

Compute Zeros


$$p(x) = ax^2 + bx + c \quad \text{side condition: } a = 0 \wedge b \neq 0$$



$$x = -c/b$$

Compute Zeros




$$p(x) = ax^2 + bx + c \quad \text{side condition: } a \neq 0 \wedge b^2 - 4ac \geq 0$$



$$x_1 = \frac{-b - \sqrt{b^2 - 4ac}}{2a}, \quad x_2 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$

Compute Test Candidates

Possible solution intervals for x on $p \sim 0$:



Constraints	$-\infty$	x_1	$x_1 + \epsilon$	x_2	$x_2 + \epsilon$	x_3	$x_3 + \epsilon$
$p = 0$	-	✓	-	✓	-	✓	-
$p > 0, p < 0, p \neq 0$	✓	-	✓	-	✓	-	✓
$p \geq 0, p \leq 0$	✓	✓	-	✓	-	✓	-


Example

$$\varphi := ((\underbrace{xy - 1}_{p_1} = 0) \wedge \underbrace{y^2 - 1}_{p_2} < 0)$$

Elimination of y :

	constraints	test candidates
1.	from all constraints	$-\infty$
2.	$p_1 = 0$	$1/x$ if $x \neq 0$
3.	$p_2 < 0$	$1 + \epsilon$
4.	$p_2 < 0$	$-1 + \epsilon$

Example



Example

$$\varphi := ((\underbrace{xy - 1}_{p_1} = 0) \wedge \underbrace{y^2 - 1}_{p_2} < 0)$$

Elimination of y :

	constraints	test candidates
1.	from all constraints	$-\infty$
2.	$p_1 = 0$	$1/x$ if $x \neq 0$
3.	$p_2 < 0$	$1 + \epsilon$
4.	$p_2 < 0$	$-1 + \epsilon$

$$\begin{aligned} \exists x \cdot \exists y \cdot \varphi &\leftrightarrow \exists x \cdot \left(\begin{aligned} &(\varphi[-\infty//y]) && \vee \\ &(\varphi[\frac{1}{x}//y] \quad \wedge x \neq 0) && \vee \\ &(\varphi[1 + \epsilon//y]) && \vee \\ &(\varphi[-1 + \epsilon//y]) \end{aligned} \right. \end{aligned}$$


Substitution of a Minus Infinity

Assume $(p(x) = ax^2 + bx + c) < 0$ and test candidate is $-\infty$

$$p(x) < 0[-\infty//x] = \underbrace{a < 0}_{\text{Case 1}} \quad \vee$$

$$\underbrace{a = 0 \wedge b > 0}_{\text{Case 2}} \quad \vee$$

$$\underbrace{a = 0 \wedge b = 0 \wedge c < 0}_{\text{Case 3}}$$



Substitution of a Minus Infinity

Assume $(p(x) = ax^2 + bx + c) < 0$ and test candidate is $-\infty$

$$p(x) < 0[-\infty//x] = \underbrace{a < 0} \quad \vee$$


Case 1

$$\underbrace{a = 0 \wedge b > 0} \quad \vee$$

Case 2

$$\underbrace{a = 0 \wedge b = 0 \wedge c < 0}$$

Case 3




Substitution of a Minus Infinity

Assume $(p(x) = ax^2 + bx + c) < 0$ and test candidate is $-\infty$

$$p(x) < 0[-\infty//x] = \underbrace{a < 0}_{\text{Case 1}} \quad \vee$$

$$\underbrace{a = 0 \wedge b > 0}_{\text{Case 2}} \quad \vee$$

$$\underbrace{a = 0 \wedge b = 0 \wedge c < 0}_{\text{Case 3}}$$



Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

$$p(x) < 0[-\infty//x] = (a < 0) \vee (a = 0 \wedge b > 0) \\ \vee (a = 0 \wedge b > 0 \wedge c < 0)$$

Elimination of y : Test candidate: $-\infty$

$$\exists x \cdot ((xy - 1 = 0)[- \infty // y] \\ \wedge (y^2 - 1 < 0)[- \infty // y])$$

Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

$$p(x) < 0[-\infty//x] = (a < 0) \vee (a = 0 \wedge b > 0) \\ \vee (a = 0 \wedge b > 0 \wedge c < 0)$$

Elimination of y : Test candidate: $-\infty$

$$\exists x \cdot ((xy - 1 = 0)[- \infty // y] \\ \wedge (y^2 - 1 < 0)[- \infty // y])$$

$$\Leftrightarrow \exists x \cdot ((x = 0 \wedge -1 = 0)$$

Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

$$p(x) < 0[-\infty//x] = (a < 0) \vee (a = 0 \wedge b > 0) \\ \vee (a = 0 \wedge b > 0 \wedge c < 0)$$

Elimination of y : Test candidate: $-\infty$

$$\exists x \cdot ((xy - 1 = 0)[- \infty // y] \\ \wedge (y^2 - 1 < 0)[- \infty // y])$$

$$\Leftrightarrow \exists x \cdot ((x = 0 \wedge -1 = 0) \\ \wedge (1 < 0 \vee (1 = 0 \wedge 0 > 0) \\ \vee (1 = 0 \wedge 0 = 0 \wedge -1 < 0)))$$

Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

$$p(x) < 0[-\infty//x] = (a < 0) \vee (a = 0 \wedge b > 0) \\ \vee (a = 0 \wedge b > 0 \wedge c < 0)$$

Elimination of y : Test candidate: $-\infty$


$$\exists x \cdot ((xy - 1 = 0)[- \infty // y] \\ \wedge (y^2 - 1 < 0)[- \infty // y])$$

$$\Leftrightarrow \exists x \cdot ((x = 0 \wedge -1 = 0) \\ \wedge (1 < 0 \vee (1 = 0 \wedge 0 > 0) \\ \vee (1 = 0 \wedge 0 = 0 \wedge -1 < 0)))$$

$$\Leftrightarrow \exists x \cdot (\mathbf{false})$$

Example: Search Tree



$$\varphi := ((\underbrace{xy - 1}_{p_1} = 0) \wedge \underbrace{y^2 - 1}_{p_2} < 0)$$



Outlook

- Implemented in SMT-RAT and Redlog
- Strengths:
 - Better than Fourier Motzkin
 - Efficient than cylindrical algebraic decomposition
- Weaknesses:
 - Unable to handle the formulas of degree > 4
 - Incomplete
 - Exponentially Complex

Reference

-  V. Weispfenning, *Quantifier elimination for real algebra - the quadratic case and beyond*. Appl. Algebra Eng. Commun. Comput, 1997.
-  R. Loss, V. Weispfenning, *Applying linear quantifier elimination*. The computer Journal 36 (1993), pp. 450-462.

Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \quad \text{with } u, q, r, s \text{ polynomials.}$$

Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \quad \text{with } u, q, r, s \text{ polynomials.}$$

Assume, $p(x) = 0$ and test candidate is $\frac{u+q\sqrt{r}}{s}$

Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \quad \text{with } u, q, r, s \text{ polynomials.}$$

Assume, $p(x) = 0$ and test candidate is $\frac{u+q\sqrt{r}}{s}$

① $(p(x) = 0)[\frac{u+q\sqrt{r}}{s} // x]$ to be computed.

Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \quad \text{with } u, q, r, s \text{ polynomials.}$$

Assume, $p(x) = 0$ and test candidate is $\frac{u+q\sqrt{r}}{s}$

- 1 $(p(x) = 0)[\frac{u+q\sqrt{r}}{s} // x]$ to be computed.
- 2 Transform the result to $\frac{u' + q'\sqrt{r}}{s'} = 0$ where u', q', s' are polynomials.

Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \quad \text{with } u, q, r, s \text{ polynomials.}$$

Assume, $p(x) = 0$ and test candidate is $\frac{u+q\sqrt{r}}{s}$

- 1 $(p(x) = 0)[\frac{u+q\sqrt{r}}{s} // x]$ to be computed.
- 2 Transform the result to $\frac{u' + q'\sqrt{r}}{s'} = 0$ where u', q', s' are polynomials.
- 3 $\frac{u' + q'\sqrt{r}}{s'} = 0$

Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \quad \text{with } u, q, r, s \text{ polynomials.}$$

Assume, $p(x) = 0$ and test candidate is $\frac{u+q\sqrt{r}}{s}$

- ① $(p(x) = 0)[\frac{u+q\sqrt{r}}{s} // x]$ to be computed.
- ② Transform the result to $\frac{u' + q'\sqrt{r}}{s'} = 0$ where u', q', s' are polynomials.
- ③ $\frac{u' + q'\sqrt{r}}{s'} = 0$
 $\iff u' + q'\sqrt{r} = 0$

Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \quad \text{with } u, q, r, s \text{ polynomials.}$$

Assume, $p(x) = 0$ and test candidate is $\frac{u+q\sqrt{r}}{s}$

- 1 $(p(x) = 0)[\frac{u+q\sqrt{r}}{s} // x]$ to be computed.
- 2 Transform the result to $\frac{u' + q'\sqrt{r}}{s'} = 0$ where u', q', s' are polynomials.
- 3 $\frac{u' + q'\sqrt{r}}{s'} = 0$
 $\iff u' + q'\sqrt{r} = 0$
 $\iff u'q' \leq 0 \wedge |u'| = |q'\sqrt{r}|$

Substitution of Square Root Expression

A square root expression has following form:

$$k = \frac{u + q\sqrt{r}}{s} \quad \text{with } u, q, r, s \text{ polynomials.}$$

Assume, $p(x) = 0$ and test candidate is $\frac{u+q\sqrt{r}}{s}$

- ① $(p(x) = 0)[\frac{u+q\sqrt{r}}{s} // x]$ to be computed.
- ② Transform the result to $\frac{u' + q'\sqrt{r}}{s'} = 0$ where u', q', s' are polynomials.
- ③ $\frac{u' + q'\sqrt{r}}{s'} = 0$

$$\iff u' + q'\sqrt{r} = 0$$

$$\iff u'q' \leq 0 \wedge |u'| = |q'\sqrt{r}|$$

$$\iff u'q' \leq 0 \wedge u'^2 - q'^2r = 0$$


Substitution of Infinitesimal Expressions

Assume $p(x) < 0$ and test candidate is $e + \epsilon$

Substitution of Infinitesimal Expressions

Assume $p(x) < 0$ and test candidate is $e + \epsilon$


$$\begin{aligned}
 (p < 0)[e + \epsilon // x] &= \underbrace{((p < 0)[e // x])}_{\text{Case 1}} \\
 &\quad \underbrace{((p = 0)[e // x] \wedge (p' < 0)[e // x])}_{\text{Case 2}} \\
 &\quad \underbrace{((p = 0)[e // x] \wedge (p' = 0)[e // x] \wedge (p'' < 0)[e // x])}_{\text{Case 3}}
 \end{aligned}$$



Substitution of Infinitesimal Expressions

Assume $p(x) < 0$ and test candidate is $e + \epsilon$


$$\begin{aligned}
 (p < 0)[e + \epsilon // x] &= \underbrace{((p < 0)[e // x])}_{\text{Case 1}} \\
 &\quad \underbrace{((p = 0)[e // x] \wedge (p' < 0)[e // x])}_{\text{Case 2}} \\
 &\quad \underbrace{((p = 0)[e // x] \wedge (p' = 0)[e // x] \wedge (p'' < 0)[e // x])}_{\text{Case 3}}
 \end{aligned}$$



Substitution of Infinitesimal Expressions

Assume $p(x) < 0$ and test candidate is $e + \epsilon$

$$\begin{aligned}
 (p < 0)[e + \epsilon // x] &= \underbrace{((p < 0)[e // x])}_{\text{Case 1}} \\
 &\quad \underbrace{((p = 0)[e // x] \wedge (p' < 0)[e // x])}_{\text{Case 2}} \\
 &\quad \underbrace{((p = 0)[e // x] \wedge (p' = 0)[e // x] \wedge (p'' < 0)[e // x])}_{\text{Case 3}}
 \end{aligned}$$



Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

Elimination of y :

2. Test candidate: $\frac{1}{x}$ if $x \neq 0$

$$\begin{aligned} \exists x \cdot & \quad (\quad (xy - 1 = 0)[\frac{1}{x} // y] \\ & \quad \wedge \quad (y^2 - 1 < 0)[\frac{1}{x} // y] \\ & \quad \wedge \quad x \neq 0) \end{aligned}$$

Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

Elimination of y :

2. Test candidate: $\frac{1}{x}$ if $x \neq 0$

$$\begin{aligned} \exists x \cdot & \left(\begin{array}{l} (xy - 1 = 0)[\frac{1}{x} // y] \\ \wedge \quad (y^2 - 1 < 0)[\frac{1}{x} // y] \\ \wedge \quad x \neq 0 \end{array} \right) \\ \Leftrightarrow \exists x \cdot & \left(\begin{array}{l} (0 = 0) \end{array} \right) \end{aligned}$$

Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

Elimination of y :

2. Test candidate: $\frac{1}{x}$ if $x \neq 0$

$$\begin{aligned} \exists x \cdot & \quad (\quad (xy - 1 = 0)[\frac{1}{x} // y] \\ & \quad \wedge \quad (y^2 - 1 < 0)[\frac{1}{x} // y] \\ & \quad \wedge \quad x \neq 0) \end{aligned}$$

$$\begin{aligned} \Leftrightarrow \exists x \cdot & \quad (\quad (0 = 0) \\ & \quad \wedge \quad ((1 > 0) \wedge 1 - x^2 < 0 \vee (1 < 0 \wedge 1 - x^2 < 0))) \end{aligned}$$

Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

Elimination of y :

2. Test candidate: $\frac{1}{x}$ if $x \neq 0$

$$\begin{aligned} \exists x \cdot & \quad (\quad (xy - 1 = 0)[\frac{1}{x} // y] \\ & \quad \wedge \quad (y^2 - 1 < 0)[\frac{1}{x} // y] \\ & \quad \wedge \quad x \neq 0) \end{aligned}$$

$$\begin{aligned} \Leftrightarrow \exists x \cdot & \quad (\quad (0 = 0) \\ & \quad \wedge \quad ((1 > 0) \wedge 1 - x^2 < 0 \vee (1 < 0 \wedge 1 - x^2 < 0)) \\ & \quad \wedge \quad x \neq 0) \end{aligned}$$

Example

$$\exists x \cdot \exists y \cdot ((xy - 1 = 0) \wedge y^2 - 1 < 0)$$

Elimination of y :

2. Test candidate: $\frac{1}{x}$ if $x \neq 0$

$$\begin{aligned} \exists x \cdot & \left((xy - 1 = 0) \left[\frac{1}{x} // y \right] \right. \\ & \wedge (y^2 - 1 < 0) \left[\frac{1}{x} // y \right] \\ & \left. \wedge x \neq 0 \right) \end{aligned}$$

$$\begin{aligned} \Leftrightarrow \exists x \cdot & \left((0 = 0) \right. \\ & \wedge ((1 > 0) \wedge 1 - x^2 < 0 \vee (1 < 0 \wedge 1 - x^2 < 0)) \\ & \left. \wedge x \neq 0 \right) \end{aligned}$$

$$\begin{aligned} \Leftrightarrow \exists x \cdot & \left((1 - x^2 < 0) \right. \\ & \left. \wedge x \neq 0 \right) \end{aligned}$$

Example

$$\exists x. (1 - x^2 < 0 \wedge x \neq 0)$$

Elimination of x :

1. Test candidate: $-\infty$

$$(1 - x_2 < 0)[- \infty // x]$$

$$= (-1 < 0 \vee (-1 = 0 \wedge 0 > 0)) \vee (-1 = 0 \wedge 0 = 0 \wedge 1 < 0))$$

$$= \text{true}$$

Example

$$\exists x. (\text{true} \wedge x \neq 0)$$

Elimination of x :

1. Test candidate: $-\infty$

$$\begin{aligned} & (x \neq 0)[- \infty // x] \\ &= (1 \neq 0 \vee 0 \neq 0) \\ &= \text{true} \end{aligned}$$

Example

$$\exists x. (\text{true} \wedge \text{true})$$

Elimination of x :

1. Test candidate: $-\infty$

$$\begin{aligned} & (x \neq 0)[- \infty // x] \\ &= (1 \neq 0 \vee 0 \neq 0) \\ &= \text{true} \end{aligned}$$