

Quantum Vacuum Catalyzation in Topological Excitonic Systems:

A Theoretical Framework for Zero-Point Restructured Energy Barriers

Mineral Hoiland
Applied Vacuum Dynamics Labs

August 2026

Abstract

We propose Quantum Vacuum Catalyzation (QVC), a theoretical framework in which zero-point fluctuations of phononic and electromagnetic fields reshape barrier-crossing rates in strongly correlated flat-band systems without contributing net energy. The host is a Topological Excitonic Vacuum Crystal (TEVC), realized in magic-angle twisted bilayer graphene (MATBG) and related moiré heterostructures, whose hopfion cores furnish a toroidal Casimir cavity of radii $(r, R) = (8, 50)$ nm. The vacuum energy of this cavity is obtained from the two-dimensional spectral form $E_{\text{vac}} = (1/4\pi^2)(\hbar c_s/r) \sum_n K_0(nx)$, which evaluates to $F = 0.58068$ and $E_{\text{vac}} = 0.1287$ meV. The condensate amplitude then obeys the self-consistent map $\delta = 1 + \alpha(\ln \delta + \ell)$ with drive $\alpha = \lambda E_{\text{vac}}/\Delta_0$. That map undergoes a saddle-node fold at finite $\Delta_c = 0.109$ meV: unit vacuum participation lies beyond the fold, while the working point $\lambda^* = 0.7628$ remains on the physical branch with static gap $\Delta^* = 0.2324$ meV. Independently, off-diagonal coupling among collective modes produces a dark (catalyzon) eigenvalue $\lambda_{\text{min}} = -g_0$ and a Mexican-hat barrier reduction of 30% at $g_0^* = 0.0982$ meV. Truncated Gross-Pitaevskii evolution over 1 ps raises the live gap to 0.6305 meV, showing that the time-dependent sector is distinct from the static fold. Fractional Hopf charge $Q_H = 1/N$ on lens spaces implies a Hall staircase of height $\delta\sigma_{xy} = e^2/(2Nh)$. A many-body Schwinger rate $\Gamma = \omega_0 \alpha^* \exp(-\pi/\alpha^*)$ yields a thermal-quantum crossover at $T^* = 118.7$ mK. We construct the topological Ginzburg-Landau functional, analyze vortex renormalization in the phase sector, and propose THz and cavity-QED tests of the mechanism.

1 Introduction

The zero-point energy of a quantized field arises as a mathematical consequence of canonical quantization: for each mode ω , the ground state contributes $\frac{1}{2}\hbar\omega$. While the sum over all modes is formally divergent, differences in vacuum energy between distinct boundary configurations are finite and observable, as demonstrated by the Casimir attraction between parallel conducting plates[1, 2]. Quantum Vacuum Catalyzation (QVC) addresses a different question: can zero-point fluctuations of electromagnetic and phononic fields dynamically reshape the effective energy landscape governing barrier-crossing processes in strongly-correlated many-body systems, yielding observable consequences in transport, spectroscopy, and pair-production phenomena? We argue the answer is affirmative, and the mechanism is vacuum *catalyzation*—a coherent coupling between zero-point modes and collective condensate modes that reduces energy barriers for processes such as exciton-pair creation, topological defect nucleation, and Schwinger-type tunneling without contributing net energy to the system.

1.1 Vacuum as Catalyst, Not Energy Source

A central clarification: QVC does not claim net energy extraction from the vacuum. Energy conservation is strictly respected; in every prediction, the energy required for barrier crossing or pair production is supplied by external sources (electric fields, laser radiation, chemical potential). The vacuum plays a role analogous to a chemical catalyst in an enzymatic reaction: it reshapes the topography of the effective potential so that transition states lie at lower free energy, without being consumed and without contributing energy. Consequently, condensed-matter processes whose rates are exponentially suppressed in the absence of vacuum catalyzation become accessible at laboratory scales—a transformation of rates, not energetics.

We note that Andolina *et al.*[3] established a rigorous no-go theorem showing that gauge invariance and the diamagnetic $\mathbf{A}^2$ response forbid equilibrium photon condensation in nonrelativistic electron systems coupled to cavity photons. QVC is not in conflict with this result for three structural reasons. First, the vacuum mode mediating catalyzation is phononic—an acoustic displacement field $\phi(\mathbf{r}, \tau)$ —not photonic; no diamagnetic $\mathbf{A}^2$ cancellation exists for lattice displacements because the coupling is scalar (deformation potential), not minimal-substitution gauge coupling. Second, the self-consistent fixed point requires a *driven* parametric channel (dynamical Casimir emission, Section 20), not an equilibrium vacuum instability; the no-go theorem applies to ground-state photon number, whereas QVC operates in a steady state sustained by external work. Third, the vacuum does not supply net energy to the condensate; it restructures the barrier landscape while the drive provides the work crossing those barriers, consistent with the catalytic (non-energetic) role emphasized above. That cavity-vacuum coupling can nonetheless modify topological transport within fully gauge-invariant formulations has been demonstrated experimentally by Appugliese *et al.*[4], who observed breakdown of integer quantum Hall protection under strong vacuum fields—providing direct evidence that vacuum-matter coupling produces measurable condensed-matter consequences beyond the scope of the no-go assumptions.

1.2 The TEVC Platform

The framework applies specifically to Topological Excitonic Vacuum Crystals (TEVCs), a class of materials exemplified by magic-angle twisted bilayer graphene (MATBG) near $\theta \approx 1.1^\circ$ [5, 6] and transition-metal dichalcogenide (TMD) heterostructures such as $\text{MoSe}_2/\text{WSe}_2$ [7, 8]. A TEVC exhibits three defining properties simultaneously:

Definition 1.1 (Topochirality). *A crystal is **topochiral** if it satisfies:*

1. *A moiré superlattice with period $\lambda_M \gg a_{\text{atomic}}$, providing a Casimir cavity at the correct geometric scale;*
2. *Non-trivial band topology characterized by non-zero Berry curvature $\Omega(\mathbf{k})$ and quantized band Chern number $C \neq 0$;*
3. *Broken time-reversal or sublattice (chirality) symmetry, induced by Kekulé bond order[9] or spin-orbit coupling, selecting definite handedness for hopfion textures mediating vacuum catalyzation.*

No single property suffices. The superlattice supplies the Casimir cavity; topology underwrites quantized Hall response; broken chirality selects hopfion handedness. For MATBG, two graphene sheets twisted by $\theta \approx 1.1^\circ$ develop a moiré superlattice with period $\lambda_M = a/[2\sin(\theta/2)] \approx 13$ nm. Near this angle, the lowest two moiré bands become nearly flat (bandwidth $W \lesssim 10$ meV). Because

the Coulomb scale U at this superlattice period is tens of meV, the flat-band limit $U/W \rightarrow \infty$ places MATBG firmly in the strongly-correlated regime[10, 11].

1.3 Three Physical Scales

Three energy scales interact non-trivially in a TEVC:

1. **Exchange energy** $J \sim 0.3$ meV, set by Coulomb interactions of flat-band electrons. In the absence of vacuum coupling, the excitonic gap is $\Delta_{\text{ex}} = 2J$.
2. **Phonon energy** $\hbar\omega_{\text{ph}} = \hbar c_s/a_H = 8.75$ meV, where $c_s = 1.06 \times 10^5$ m/s follows from $\hbar c_s = 0.07$ meV $\cdot \mu\text{m}$ and $a_H = \hbar/\sqrt{2m^*\Delta_0} = 8$ nm is the healing length of the excitonic condensate (with $m^* = 0.99 m_e$ from $\kappa_{\text{GL}} = \Delta_0 a_H^2$; Section 10.2).
3. **Casimir energy** $E_{\text{Cas}}(r, R)$ of a toroidal vortex core with minor radius $r \sim a_H$ and major radius $R \sim \xi$ (coherence length). This energy determines the self-consistent zero-point field amplitude A_{ZPF} coupling into the gap equation.

The Casimir energy is neither a small perturbation (because moiré flat bands quench kinetic energy that would ordinarily dominate) nor a dominant effect (because the moiré period remains well-separated from the atomic scale). It sits precisely in the regime where self-consistent feedback between vacuum and condensate becomes physically relevant.

1.4 Organization

The paper is organized as follows. Sections 2–5 introduce the TEVC platform, the self-consistent gap equation and its fold, the toroidal vacuum energy, and the participation bridge $\alpha = \lambda E_{\text{vac}}/\Delta_0$. Section 6 treats the phase sector as a two-dimensional vortex renormalization-group problem. Sections 7–13 develop the catalyzon, the topological Ginzburg–Landau functional, the matched coefficient dictionary, and time-dependent dynamics. Sections 14–17 collect experimental signatures, numerical predictions, and implications. The remaining sections treat cavity QED, Schwinger pair production, dynamical Casimir saturation, Berry phase, kinematics, and the topological and analogue-gravity extensions. Conventions and provenance are recorded in Section 10 and Appendix A.

2 The TEVC Platform and Symmetry-Breaking Cascade

The TEVC ground state is reached not by a single mean-field transition but by a sequential cascade of symmetry breaking. Starting from the enlarged $U(4) \times U(4)$ symmetry of the flat-band manifold (spin $\otimes$ valley), three stages occur:

$$U(4) \times U(4) \xrightarrow{T_1 \sim 30 \text{ K}} G_1 \text{ (valley polarization, } P_v) \quad (1)$$

$$G_1 \xrightarrow{T_2 \sim 6 \text{ K}} G_2 \text{ (Kekulé order, } \Delta_{KK'}) \quad (2)$$

$$G_2 \xrightarrow{T_{\text{BEC}} \sim 1 \text{ K}} G_{\text{BEC}} \text{ (excitonic condensation, } \Delta_{\text{ex}}) \quad (3)$$

Once $P_v \neq 0$ and $\Delta_{KK'} \neq 0$, the two act as a trilinear source for the excitonic order parameter through the Landau coupling

$$\mathcal{F}_{\text{tri}} = \delta \Delta_{\text{ex}}^* \Delta_{KK'} P_v + \text{h.c.}, \quad (4)$$




Figure 1: **TEVC multilayer structure and moiré superlattice.** (Left) Five-layer twisted graphene stack showing alternating twist angles $\pm 1.1^\circ$ with interlayer spacing $d_\perp = 0.34$ nm. Each layer contains hexagonal graphene lattice (colored points) with moiré unit cell overlay (purple hexagon, $\lambda_M \approx 13$ nm). Gray lines indicate AA-site Josephson coupling. (Center) Moiré superlattice pattern from interference of twisted layers showing AA-site maxima (red circles) where interlayer coupling is strongest. Moiré unit cell (purple) has period $\lambda_M = a/(2 \sin(\theta/2)) \approx 13$ nm for twist angle $\theta = 1.1^\circ$. (Right) Cross-sectional view showing layer stack ($L_z = 1.36$ nm total thickness), Josephson coupling J between adjacent layers (green arrows), excitonic condensate clouds (cyan), valley polarization ($C = +1$), Kekulé order, and Chern-Simons level $k = 2N = 10$.




Figure 2: **Topochiral symmetry-breaking cascade.** The TEVC platform emerges through sequential symmetry breaking from $U(4) \times U(4)$ parent symmetry at $T \sim 30$ K down to excitonic condensation at $T_{BEC} \sim 1$ K. Each stage corresponds to a distinct energy scale. Zero-point field coupling enters non-thermally at the lowest temperature scale, dynamically locking the topochiral order.




Figure 3: **Computed Bistritzer–MacDonald band structure at $\theta = 1.1^\circ$.** Moiré bands along the high-symmetry path Γ – K – M – Γ in the moiré Brillouin zone, showing the nearly-flat lowest bands with bandwidth $W \lesssim 10$ meV that place the system in the strong-coupling regime $U/W \gg 1$. Computed using the continuum BM model with interlayer coupling $w_{AA} = 79.7$ meV, $w_{AB} = 97.5$ meV on a 7-shell reciprocal lattice expansion. The flat-band manifold is the substrate on which the QVC gap equation operates.

making the emergence of the excitonic condensate inductive rather than independent. The final step is not thermodynamic but dynamical: as the excitonic condensate stiffens, its toroidal Casimir cavity (see Section 4) reaches the critical aspect ratio for zero-point field self-consistency, and the TEVC ground state locks in.

3 Self-Consistent Gap Equation and Gap Collapse

3.1 Path Integral Formulation

We begin with the Euclidean action for the excitonic condensate field $\Psi(\mathbf{r}, \tau)$ coupled linearly to a phonon field $\phi(\mathbf{r}, \tau)$:

$$S[\Psi, \phi] = S_\Psi[\Psi] + S_\phi[\phi] + S_{\Psi\phi}[\Psi, \phi], \quad (5)$$

with

$$S_\Psi = \int d\tau d^2r \left[\Psi^* \partial_\tau \Psi + \frac{\hbar^2}{2m^*} |\nabla \Psi|^2 + \alpha |\Psi|^2 + \frac{\beta}{2} |\Psi|^4 \right], \quad (6)$$

$$S_\phi = \frac{1}{2} \int d\tau d^2r \left[(\partial_\tau \phi)^2 + (c_s \nabla \phi)^2 \right], \quad (7)$$

$$S_{\Psi\phi} = g \int d\tau d^2r \phi(\mathbf{r}, \tau) |\Psi(\mathbf{r}, \tau)|^2. \quad (8)$$

The coupling g encodes how phonon density displacement modulates excitonic condensate energy through the deformation potential and Coulomb exchange corrections. Because S_ϕ is quadratic in ϕ , we integrate out the phonon field exactly. The Gaussian integration yields an effective action:

$$S_{\text{eff}}[\Psi] = S_\Psi[\Psi] - \frac{g^2}{2} \int d\tau d\tau' d^2r d^2r' |\Psi(\mathbf{r}, \tau)|^2 D(\mathbf{r} - \mathbf{r}', \tau - \tau') |\Psi(\mathbf{r}', \tau')|^2, \quad (9)$$

where D is the phonon propagator

$$D(\mathbf{q}, \omega_n) = \frac{1}{\omega_n^2 + c_s^2 q^2}. \quad (10)$$

Self-consistent gap equation. After integrating out the phonon and imposing a static, translationally invariant condensate, the working amplitude equation on a cavity of fixed healing length $a_H = r$ is

$$\delta = 1 + \alpha (\ln \delta + \ell), \quad \ell = \ln \frac{\Delta_0 a_H}{2\hbar c_s} + \gamma_E, \quad \alpha = \lambda \frac{E_{\text{vac}}}{\Delta_0}, \quad (11)$$

or, equivalently,

$$\Delta = 2J + \hbar c_s A_{\text{ZPF}} \left[\ln \left(\frac{\Delta a_H}{2\hbar c_s} \right) + \gamma_E \right]. \quad (12)$$

The logarithm is the regulated remnant of the two-dimensional phonon loop derived below; $\gamma_E \approx 0.5772$ is the Euler–Mascheroni constant. Fold algebra of this map is exact: $\alpha_c = \delta_c$ and $\delta_c(1 - \ln \delta_c - \ell) = 1$, with square-root branching $\delta - \delta_c \sim \pm \sqrt{\alpha_c - \alpha}$ as $\alpha \rightarrow \alpha_c^-$. Numerical solution yields two real roots below α_c , no physical root at unit participation, a physical root at λ^* , and a log-log slope 0.536 against the analytic 1/2 on $t_A = (\alpha_c - \alpha)/\alpha_c$ (Appendix A.1). Finite $\Delta_c = 0.109 \text{ meV}$ distinguishes the saddle-node from BKT vanishing of the gap.

Phonon loop and infrared regularization. The Euclidean propagator admits analytic continuation to $D^R(\mathbf{q}, \omega) = 2\omega_q/(\omega^2 - \omega_q^2 + i\eta)$ with $\omega_q = c_s |\mathbf{q}|$. In the static limit this produces an attractive kernel $V_{\text{eff}}(\mathbf{q}) = -g_0^2/(c_s^2 q^2)$. Integrating over the two-dimensional Brillouin zone between an infrared momentum q_{IR} and the ultraviolet cutoff $\Lambda = 1/a_H = 125 \mu\text{m}^{-1}$ gives

$$\Sigma = -\frac{g_0^2}{2\pi c_s^2} \int_{q_{\text{IR}}}^{1/a_H} \frac{dq}{q} = -\frac{g_0^2}{2\pi c_s^2} \ln \left(\frac{\Lambda}{q_{\text{IR}}} \right). \quad (13)$$

A gap-scale infrared cutoff $q_{\text{IR}} = 2\hbar c_s/(\Delta a_H^2)$ evaluates to $3.65 \times 10^3 \mu\text{m}^{-1}$ and therefore lies above Λ . The kinematically consistent gap momentum $q_{\text{gap}} = \Delta_0/(\hbar c_s) = 8.57 \mu\text{m}^{-1}$ restores $\Lambda > q_{\text{IR}}$ but yields $\ln(\Lambda/q_{\text{gap}}) = +2.68$, opposite in sign to $\ln(\Delta_0 a_H/(2\hbar c_s)) = -3.37$ that enters $\ell = -2.7958 < 0$. Attractive one-loop Σ at the consistent cutoff would therefore *enhance* the gap, whereas Eq. (12) with $\ell < 0$ *suppresses* it. We therefore treat (12) as a regulated self-consistent equation whose logarithm is motivated by the phonon loop but whose sign and infrared endpoint are fixed by the frozen-cavity evaluation of E_{vac} (Section 4). Retardation remains $O(\eta_{\text{ret}})$ with $\eta_{\text{ret}} = 2\Delta_0\xi/(\hbar c_s)$ (Section 29).

3.2 Static Approximation and Retardation

Retardation is governed by the causal-propagator identity

$$\eta_{\text{ret}} = \frac{2\Delta_0\xi}{\hbar c_s}. \quad (14)$$

On the Ginzburg–Landau healing length $a_H = 8 \text{ nm}$ this is $\eta_{\text{ret}}(a_H) = 0.137$; on the phonon healing $\xi_{\text{ph}} = \hbar c_s/\Delta_0$ it is the identity $\eta_{\text{ret}}(\xi_{\text{ph}}) = 2$ (Section 10). These lengths are not interchangeable. The leading Matsubara contribution to S_{eff} is the static $\omega_n = 0$ term, $V_{\text{eff}}(\mathbf{q}) = -g^2/(c_s^2 q^2)$. Equation (12) is the self-consistent closure of that kernel on a cavity of fixed a_H ; it is not obtained by a BCS decoupling of the quartic. The gap appears linearly on the left and under a logarithm on the right.

3.3 Gap Collapse: Fold Bifurcation and Vortex-RG Sector

The structure of Eq. (12) is closely related to self-consistent gap equations that appear in the BKT problem[12, 13]. Written in polar form $\Psi = \rho e^{i\theta}$, the theory has two sectors. In the amplitude ρ , the mean-field map $\delta = 1 + \alpha(\ln \delta + \ell)$ undergoes a *saddle-node* (fold) bifurcation: the physical and unstable branches merge at finite δ_c , with square-root scaling $\delta - \delta_c \sim \pm\sqrt{\alpha_c - \alpha}$ as $\alpha \rightarrow \alpha_c^-$. In the phase θ , vortex renormalization-group flow of the stiffness κ implies a Berezinskii–Kosterlitz–Thouless essential singularity

$$\Delta(A_{\text{ZPF}}) \sim \exp\left(-\frac{C}{\sqrt{A_{\text{ZPF}}^c - A_{\text{ZPF}}}}\right), \quad A_{\text{ZPF}} \rightarrow A_{\text{ZPF}}^c, \quad (15)$$

whenever the physical drive reaches the KT surface before the fold (Section 6, Theorem 6.1, hypothesis H-VRG1). Experimental discrimination between square-root fold softening and essential-singularity collapse is a key test of which infrared surface is realized.

3.4 Static fold, catalyzon, and Gross–Pitaevskii dynamics

The static fold, the catalyzon Mexican-hat, and truncated Gross–Pitaevskii evolution probe distinct operators and are not combined as additive percentages of one baseline. They share the same vacuum E_{vac} and participation λ^* . On the amplitude map (11), unit λ lies beyond the fold, λ^* sits on the physical branch at $\Delta^* = 0.2324 \text{ meV}$, and the branches merge at $\Delta_c = 0.109 \text{ meV}$. The catalyzon channel is the democratic spectrum of $G = g_0(J - I)$ at $g_0^* = \lambda^* E_{\text{vac}}$, giving $\Delta_{\text{cat}} = \Delta_0 - g_0^*$ and a well-depth ratio $(\Delta_{\text{cat}}/\Delta_0)^2$ that corresponds to a 30% barrier reduction at fixed Track A quartic (Section 10.5). The GPE channel uses an attractive V_{ZPF} : over 1 ps the live gap *increases* to 0.6305 meV and the corresponding well *deepens*. Increasing α on the static map suppresses δ

because $\ell < 0$; that is opposite in sign to BCS pairing, in which an attractive kernel opens a gap. The Landau potential of the catalyzon channel,

$$U_{\text{eff}}(|\Psi|) = \alpha_{\text{GL}}|\Psi|^2 + \frac{\beta}{2}|\Psi|^4, \quad (16)$$

is shallower when $|\alpha_{\text{GL}}|$ tracks Δ_{cat} and deeper under the GPE shift $-g_0^*\rho^2$. Both evaluations are reported; they are not the same operator.




Figure 4: **Self-consistent gap map.** Numerical physical and unstable branches of $\delta = 1 + \alpha(\ln \delta + \ell)$ terminate in a saddle-node fold at finite δ_c with square-root scaling $\delta - \delta_c \sim \pm \sqrt{\alpha_c - \alpha}$. The dotted curve is the vortex-sector essential-singularity form of hypothesis H-VRG1 (Section 6).

3.5 Catalyzon-Mediated Quasiparticle Pairing and Modified BCS Structure

The structure of Eq. (12) departs from conventional BCS superconductivity[14] in ways that illuminate a broader principle: *pairing of composite quasiparticles in flat-band systems generically deviates from bare-electron BCS predictions.* Recent work on rhombohedral trilayer graphene (RTG) demonstrates that superconductivity near intervalley coherent (IVC) phase boundaries exhibits $T_c \propto \epsilon_D \exp(-2/\rho_{\text{qp}}U)$ —a factor-of-2 suppression relative to standard BCS—because Cooper pairs form from *IVC quasiparticles* rather than bare electrons[15]. The QVC framework predicts an analogous phenomenon mediated by the catalyzon.

3.5.1 Catalyzon Pairing Kernel

In a TEVC, excitonic quasiparticles are vacuum-dressed through the catalyzon mechanism (Section 8). The pairing interaction between two catalyzon quasiparticles separated by wavevector $\mathbf{q}$ is




Figure 5: **Gap phase diagram (schematic).** For $A_{\text{ZPF}} < A_c$ the mean-field amplitude map admits a gapped physical branch; beyond the fold there is no real gapped solution of that map. The vortex-RG sector implies an essential singularity if the KT surface is reached first (Section 6).




Figure 6: **Numerically integrated vortex RG flow.** Kosterlitz renormalization trajectories in the (K^{-1}, y) plane computed by RK4 integration of the flow equations $dK^{-1}/d\ell = 4\pi^3 y^2$, $dy/d\ell = (2 - \pi K)y$. Horizontal axis is inverse stiffness K^{-1} ; vertical axis is vortex fugacity y . Trajectories with $K_0^{-1} < K_c^{-1}$ (blue) flow to the bound-pair phase; those starting left of the separatrix (orange) flow to the free-vortex plasma. Vertical dashed lines mark the bosonic critical stiffness $K_c^{\text{boson}} = 2/\pi$ and the anyon-corrected K_c^{anyon} for $N = 5$ layers. At the physical operating point ($T^* = 118.7$ mK, $K_0 = 46.7$), the system lies deep in the bound phase: $K \gg K_c$, confirming that BKT vortex unbinding does not preempt the saddle-node fold (Section 3).




Figure 7: **Frozen- α Kosterlitz flow at T^*** . Fugacity $y(\ell)$ at fixed drive α . Ultraviolet, working-point, and near-fold stations remain in the bound phase ($y \rightarrow 0$).




Figure 8: **Competing-scale scan at frozen α and T^*** . Kosterlitz $K(\alpha)$ along the physical branch for the primary stiffness closure (healing-disk plus geometric floor, $\kappa_{UV} \approx 0.477$ meV, $K(\alpha_c) \approx 10.57$) and the conventional-only variant. Both remain above $K_c = 2/\pi$ in the bare data.

not the bare Coulomb exchange $V(\mathbf{q})$ but the *vacuum-renormalized kernel*:

$$\mathcal{K}_{\text{cat}}(\mathbf{q}, \omega) = V(\mathbf{q}) + \sum_{i=1}^5 \frac{|g_i(\mathbf{q})|^2}{\omega^2 - \omega_i^2(\mathbf{q})} \cdot \left(1 + \frac{\lambda_{\text{min}}}{\hbar\omega_0}\right), \quad (17)$$

where $g_i(\mathbf{q})$ are the coupling vertices between excitonic quasiparticles and the i -th collective mode, $\omega_i(\mathbf{q})$ are the five mode dispersions, and the factor $(1 + \lambda_{\text{min}}/\hbar\omega_0)$ encodes the catalyzon dressing. At the resonance wavevector q^* where all five modes become degenerate at $\omega_0 = 2\Delta/\hbar$, the kernel develops a pole:

$$\mathcal{K}_{\text{cat}}(\mathbf{q}^*, \omega \rightarrow \omega_0) \sim \frac{G_{\text{eff}}}{\omega - \omega_0 + i\Gamma_{\text{cat}}}, \quad (18)$$

where $G_{\text{eff}} = \sum_i |g_i|^2 / (2\omega_0)$ is the effective coupling and Γ_{cat} is the catalyzon lifetime.

3.5.2 Modified Gap Equation: Band-Edge Suppression

The linearized self-consistency equation for catalyzon-mediated pairing near the excitonic band edge takes the form:

$$1 = \rho_{\text{cat}} \mathcal{K}_{\text{eff}} \int_0^{\epsilon_D} \frac{d\epsilon}{\sqrt{\epsilon(\epsilon + 2\Delta_{\text{cat}})}} \tanh\left(\frac{\sqrt{\epsilon(\epsilon + 2\Delta_{\text{cat}})}}{2k_B T}\right), \quad (19)$$

where $\rho_{\text{cat}} = m^*/(2\pi\hbar^2) \cdot (\Delta_{\text{cat}}/\sqrt{\Delta_{\text{cat}}^2 + \mu^2})$ is the catalyzon density of states and $\mathcal{K}_{\text{eff}}$ is the angle-averaged pairing kernel. Two regimes emerge:

Away from band edge ($|\mu| \gg \Delta_{\text{cat}}$): Standard BCS recovered with $\rho_{\text{cat}} \rightarrow \rho_0$ and $T_c \propto \epsilon_D \exp(-1/\rho_0 \mathcal{K}_{\text{eff}})$. **Near band edge** ($|\mu| \lesssim \Delta_{\text{cat}}$): The integration range is effectively halved because no catalyzon states exist within the gap Δ_{cat} , yielding

$$\boxed{T_c^{\text{QVC}} \propto \epsilon_D \exp\left(-\frac{2}{\rho_{\text{cat}} \mathcal{K}_{\text{eff}}}\right)}. \quad (20)$$

The factor of 2 in the exponent represents a *universal suppression* for quasiparticle pairing near a gap edge—identical in mathematical origin to that found in RTG IVC-superconductivity[15], but here arising from catalyzon band-edge physics rather than intervalley coherence.

3.5.3 Anomalous Coherence Length

The Ginzburg-Landau coherence length for catalyzon-mediated pairing near the band edge is

$$\xi_{\text{QVC}} \approx \frac{1}{4} \frac{v_{\text{cat}}}{\sqrt{|\mu| T_c^{\text{QVC}}}}, \quad (21)$$

where $v_{\text{cat}} = \partial\omega_{\text{cat}}/\partial k|_{k^*}$ is the catalyzon group velocity. For MATBG parameters with $v_{\text{cat}} \sim c_s \approx 10^4$ m/s, $|\mu| \sim \Delta_{\text{cat}} = 0.502$ meV, and $T_c^{\text{QVC}} \sim 1$ K, this yields $\xi_{\text{QVC}} \sim 40$ –80 nm—consistent with the healing length $a_H \approx 8$ nm and hopfion radius $R \approx 50$ nm, and *two orders of magnitude shorter* than the naive BCS prediction $\xi_{\text{BCS}} = \hbar v_F / (k_B T_c) \sim 10^4$ nm based on the bare graphene Fermi velocity $v_F \approx 10^6$ m/s. This dramatic coherence length suppression is a *hallmark prediction* of catalyzon-mediated pairing: because the relevant velocity scale is the collective-mode group velocity $v_{\text{cat}} \ll v_F$, the Cooper pair size is set by the collective excitation wavelength rather than the electronic mean free path.

3.5.4 Catalyzon vs. BCS: Summary of Distinctions

The catalyzon pairing mechanism differs from standard BCS in three measurable ways:

1. **Transition temperature:** $T_c^{\text{QVC}} \propto \exp(-2/\rho\mathcal{K})$ vs. $T_c^{\text{BCS}} \propto \exp(-1/\rho V)$ —factor-of-2 suppression from band-edge restriction.
2. **Coherence length:** $\xi_{\text{QVC}} \propto v_{\text{cat}}/\sqrt{\mu T_c}$ vs. $\xi_{\text{BCS}} \propto v_F/T_c$ —anomalously short due to collective velocity replacement.
3. **Catalytic nature:** Unlike BCS where phonons provide binding energy, the catalyzon reshapes the effective barrier without contributing net energy—barrier reduction rather than condensation energy.

4 Toroidal Casimir Geometry and Zero-Point Amplitude




Figure 9: **Toroidal hopfion geometry (3D visualization).** Hopfion vortex core has minor radius $r \approx a_H \approx 8$ nm (healing length) and major ring radius $R \approx \xi \approx 50$ nm (coherence length). Aspect ratio $x = 2\pi r/R \approx 1.0$ places the system in the fat-torus regime where the proximity force approximation breaks down and full Bessel mode sum is required, yielding $\mathcal{F}(x) = 0.58068$ and $E_{\text{vac}} = 0.1287$ meV.




Figure 10: **3D hopfion torus geometry.** The vortex core centerline (red major circle, radius $R \approx \xi \approx 50$ nm) defines the toroidal Casimir cavity. The minor radius $r \approx a_H \approx 8$ nm sets the cavity gap. Purple curves represent Hopf fibration field lines carrying topological charge $Q_H = 1/5$. The aspect ratio $x = 2\pi r/R \approx 1.0$ places the system in the fat-torus regime requiring full Bessel mode sum.

The zero-point amplitude A_{ZPF} in Eq. (12) is not a free parameter; it is fixed by the Casimir energy of the vortex-ring structure of each hopfion in the TEVC. Each hopfion has a vortex core of topological radius $r \sim a_H$ (minor) and a ring radius R (major). The Casimir energy of the resulting toroidal cavity determines the vacuum stress inside the core and, through that, the effective amplitude of the zero-point field felt by the condensate.

4.1 Toroidal spectral form

The vacuum energy used throughout is the two-dimensional toroidal spectral form

$$E_{\text{vac}} = \frac{1}{4\pi^2} \cdot \frac{\hbar c_s}{r} \cdot \mathcal{F}(x), \quad \mathcal{F}(x) = \sum_{n=1}^{\infty} K_0(nx), \quad x = \frac{2\pi r}{R}, \quad (22)$$

with K_0 the modified Bessel function of the second kind. For $r = 8$ nm and $R = 50$ nm one has $x = 2\pi r/R = 1.0053$, in the fat-torus regime where a proximity-force plate approximation does not apply. Equation (22) is a phononic spectral sum on the torus, not a three-dimensional electromagnetic scattering-matrix Casimir energy.

4.2 Numerical Calculation

We evaluate Eq. (22) by summing $K_0(nx)$ (asymptotic expansion for $z > 0.5$, Abramowitz–Stegun 9.8.1–9.8.6). For $x = 1.0053$ the sum converges to

$$\mathcal{F}(x = 1.0053) = 0.58068. \quad (23)$$

With $\hbar c_s = 0.07 \text{ meV} \cdot \mu\text{m}$ this yields

$$E_{\text{vac}} = 0.1287 \text{ meV}. \quad (24)$$

The corresponding geometric amplitude is $A_{\text{geom}} = E_{\text{vac}}/(\hbar c_s) = 1.839 \mu\text{m}^{-1}$.




Figure 11: **Casimir mode sum convergence.** Partial sum $\mathcal{F}_N(x) = \sum_{n=1}^N K_0(nx)$ as a function of the number of winding modes N , for the fat-torus aspect ratio $x = 2\pi r/R = 1.0053$. The sum saturates by $N = 14$ to $\mathcal{F} = 0.58068$, demonstrating rapid convergence driven by the exponential decay of $K_0(z)$ at large argument. The final vacuum energy $E_{\text{vac}} = 0.1287$ meV is stable to better than 10^{-6} beyond 14 modes.




Figure 12: **Spectral form factor $F(x)$ for the fat-torus geometry.** The form factor $F(x) = \sum_{n=1}^{\infty} K_0(nx)$ evaluated at the locked aspect ratio $x = 2\pi r/R = 1.0053$ (vertical marker). The rapid exponential suppression of $K_0(nx)$ for $n \geq 2$ ensures that only the lowest winding modes contribute appreciably. At the physical aspect ratio, $F = 0.58068$, yielding $E_{\text{vac}} = (1/4\pi^2)(\hbar c_s/r) F = 0.1287 \text{ meV}$.

4.3 Zero-Point Amplitude

The geometric amplitude associated with (22) is

$$A_{\text{geom}} = \frac{E_{\text{vac}}}{\hbar c_s} = 1.839 \mu\text{m}^{-1}, \quad (25)$$

or $A_{\text{geom}} R \approx 0.0919$ on $R = 50 \text{ nm}$. This is the amplitude that enters $\alpha = \lambda E_{\text{vac}}/\Delta_0$. Jacobi ϑ_3 is a distinct regularized spectral object and is not identified with $F(x)$.

4.4 Self-Consistency Loop

The cavity geometry is held at $a_H = r$. There is no running-healing loop $\Delta \rightarrow a_H(\Delta) \rightarrow E_{\text{vac}}$: E_{vac} is computed from the frozen torus, and participation λ enters only through $\alpha = \lambda E_{\text{vac}}/\Delta_0$ (Section 5). At $\lambda = 1$ the map lies beyond the fold. The working static root is

$$\boxed{\Delta^* = 0.2324 \text{ meV} \quad (\lambda^* = 0.7628),} \quad (26)$$

a 61.3% reduction from $\Delta_{\text{ex}} = 2J = 0.600 \text{ meV}$, with maximum physical-branch reduction 81.8% at $\Delta_c = 0.109 \text{ meV}$.

Self-Consistency Loop for Gap Determination




Figure 13: **Self-consistency on the frozen cavity.** Geometry (r, R) and $a_H = r$ are held fixed. E_{vac} is the Bessel spectral form. Participation λ maps E_{vac} into α ; unit λ lies beyond the fold ($\alpha/\alpha_c \approx 1.18$). Static physics at the working point uses $\lambda^* = 0.7628$ and $\Delta^* = 0.2324$ meV.

5 Derived α - E_{vac} Bridge and Participation λ

The fat-torus freeze $(r, R) = (8, 50)$ nm, the spectral form

$$E_{\text{vac}} = \frac{1}{4\pi^2} \frac{\hbar c_s}{r} F\left(\frac{2\pi r}{R}\right), \quad F(x) = \sum_{n=1}^{\infty} K_0(nx), \quad (27)$$

and package $\hbar c_s = 0.07$ meV $\cdot \mu\text{m}$ fix the vacuum energy independently of a running healing length. Frozen-cavity geometry sets $a_H = r = \text{const}$, so

$$\ell(r) = \ln \frac{\Delta_0 r}{2\hbar c_s} + \gamma_E = -2.7958. \quad (28)$$

Identifying $A = E_{\text{vac}}/(\hbar c_s)$ and $\alpha = \hbar c_s A/\Delta_0$ yields the boxed bridge

$$\alpha = \lambda \frac{E_{\text{vac}}}{\Delta_0}. \quad (29)$$

The gap map $\delta = 1 + \alpha(\ln \delta + \ell)$ has a saddle-node fold at $\alpha_c = \delta_c = 0.1818$, hence $\Delta_c = 0.109$ meV. Unit participation ($\lambda = 1$) gives $\alpha/\alpha_c \approx 1.180 > 1$: there is no physical static Δ^* at $\lambda = 1$. A physical static gap exists for $\lambda \leq \lambda_{\text{fold}} = \alpha_c \Delta_0 / E_{\text{vac}} = 0.8475$. The recommended working point is

$$\lambda^* = 0.90 \lambda_{\text{fold}} = 0.7628, \quad \Delta^* = 0.2324 \text{ meV} \quad (30)$$

(61.3% reduction from $\Delta_0 = 0.600$ meV). The deepest static reduction on the physical branch is 81.8% at the fold. Running healing $a_H = \hbar c_s / \Delta$ is a distinct theory from the frozen-cavity freeze used here.

Catalyzon hybridization is a parallel channel: $\Delta_{\text{cat}} = \Delta_{\text{ex}} - g_0$ with democratic $\lambda_{\text{min}} = -g_0$ (multiplicity $N - 1$). Fold, catalyzon, and truncated GPE dynamics are not additive percentages of a single baseline.




Figure 14: **Bridge fold with operating point.** Gap map physical and unstable branches $\delta_{\pm}(\alpha/\alpha_c)$ with locked $\ell = -2.7958$. Gold square: recommended working point $\lambda^* = 0.7628$ ($\Delta^* = 0.2324$ meV, $\alpha^*/\alpha_c = 0.900$). Red circle: fold critical point ($\alpha_c = 0.1818$, $\delta_c = 0.1818$). Arrow indicates the direction of increasing vacuum participation λ . The gap reduction from Δ_0 to Δ^* is 61.3%; maximum physical-branch reduction at the fold is 81.8%.




Figure 15: **Gap reduction as a function of participation λ .** Physical-branch gap $\Delta(\lambda)$ computed from the bridge relation $\alpha = \lambda E_{\text{vac}}/\Delta_0$ inserted into the gap map. The gap decreases monotonically with λ , dropping from $\Delta_0 = 0.600$ meV at $\lambda = 0$ to $\Delta_c = 0.109$ meV at $\lambda_{\text{fold}} = 0.8475$. Vertical marker at $\lambda^* = 0.7628$ shows the locked working point.

6 Vortex sector, competing infrared scales, and BKT implications

The excitonic order parameter admits a polar decomposition before any renormalization-group analysis,

$$\Psi(\mathbf{r}) = \rho(\mathbf{r}) e^{i\theta(\mathbf{r})}. \quad (31)$$

The locked gap map of Section 3 lives in the amplitude ρ . The vortex sector of the theory lives in the phase θ . Together the two sectors furnish a competing-scale structure: a saddle-node fold in ρ , and a Kosterlitz–Thouless unbinding surface in the stiffness of θ . When a physical drive reaches the KT surface first, the implied gap closing is a Berezinskii–Kosterlitz–Thouless essential singularity. That implication is part of the theory; it is stated as hypothesis H-VRG1 and is evaluated below by the computational suite C1–C6.

Option A* freezes the toroidal cavity, so the vacuum amplitude—and therefore α —does not run with the KT scale $\ell = \ln(L/a_c)$. The suite therefore scans frozen α along the locked physical branch $\delta_+(\alpha)$. Kosterlitz’s coupling is $K = \kappa/T_{\text{eff}}$, with critical value $K_c = 2/\pi$. No bare transition temperature is inserted by hand, and no chain-rule β_α is integrated.

Defect vocabulary. A **hopfion** is a map $S^3 \rightarrow S^2$ with fractional Hopf charge $Q_H = 1/N$, realized here as a toroidal Casimir cavity. A **2D phase vortex** is a point defect in the moiré plane with $\oint \nabla\theta \cdot d\mathbf{l} = 2\pi n$; this is the BKT defect of the present section. A **Chern–Simons fluxon** is a 2π winding that binds flux $\Phi_{\text{CS}} = 2\pi/N$ at level $k = 2N$. The three objects are distinct.

6.1 Two-sector decomposition

Substituting Eq. (31) into the bulk TQGL energy of Section 9 separates a Higgs potential for ρ from a phase stiffness for θ ,

$$\mathcal{F}_{\text{bulk}}[\rho, \theta] = \int d^2r \left[\alpha_{\text{GL}} \rho^2 + \frac{\beta}{2} \rho^4 + \kappa_{\text{GL}} (|\nabla\rho|^2 + \rho^2 |\nabla\theta|^2) \right], \quad (32)$$

up to the Chern–Simons, Berry, and Skyrme terms retained in the parent functional. Here κ_{GL} is the Ginzburg–Landau gradient weight written κ in Section 9, and α_{GL} is the Landau mass, distinct from the gap-map coupling α of Section 3. Equation (32) captures the algebra of the split. The fold of the amplitude map at $\alpha_c = \delta_c = 0.1818$, $\Delta_c = 0.109 \text{ meV}$ is a locked input. Vortex unbinding is the content of the phase sector.


The live-gap estimator $\Delta_{\text{live}}[\psi] = \Delta_0 \sqrt{n_w[\psi]/n_{\text{ref}}}$ of Section 13 is an amplitude proxy. A field whose modulus remains finite while θ disorders still returns a nonzero Δ_{live} . Amplitude and phase diagnostics therefore complement one another: a 1 ps trajectory with $\Delta_{\text{live}}(1 \text{ ps}) = 0.6305 \text{ meV}$ constrains ρ , while $g(r)$ and the stiffness flow constrain θ .

6.2 C1: phase stiffness

The 2D phase stiffness (superfluid density) extracted from Eq. (32) is not κ_{GL} itself. Matching $\kappa_{\text{GL}} |\nabla\Psi|^2$ to the conventional $(\rho_s/2) |\nabla\theta|^2$ with $\Psi = \sqrt{n_0} \rho e^{i\theta}$ gives

$$\kappa \equiv \rho_s = 2n_0 \kappa_{\text{GL}} \langle \rho^2 \rangle + \kappa_{\text{geom}}, \quad (33)$$

where $\kappa_{\text{GL}} = \Delta_0 a_H^2$ follows from the truncated GPE kinetic match and $\langle \rho^2 \rangle = \delta_+(\alpha)^2$ is read from the locked physical branch through the live-gap identification. The density unit is the healing-disk ansatz $n_0 = 1/(\pi a_H^2)$, which converts dimensionless $|\psi|^2 \sim 1$ into a 2D density and makes the



Live-gap estimator $\Delta_0 \sqrt{n_w/n_{\text{ref}}}$ reads ρ , not vortex unbinding. Chern–Simons is **TRUNCATED** in the GPE and in the default XY–KT scan.

Figure 16: **Two-sector split**. The locked fold is an amplitude phenomenon. Vortex RG is a phase phenomenon. The live-gap estimator reads ρ . Chern–Simons is truncated from the default XY–KT scan, matching the GPE truncation of Section 13.

conventional piece at unit amplitude equal to $2\Delta_0/\pi$, independent of a_H . The geometric weight κ_{geom} is a frozen Peotta–Törmä floor [16],

$$\kappa_{\text{geom}} = \frac{\Delta_0}{2\pi}, \quad (34)$$

treated as an ansatz: it is not a Brillouin-zone integral of the TEVC quantum metric, and it need not vanish with δ . From here on in this section, κ means the stiffness (33). A uniform phase-twist check on the same conventional piece agrees with $2n_0\kappa_{\text{GL}}\langle\rho^2\rangle$ to machine precision.

On this primary vortex-RG closure—healing-disk n_0 plus the frozen geometric floor (34)—one finds $\kappa_{\text{UV}} \approx 0.477 \text{ meV}$ at $\alpha = 0$ and $\kappa(\alpha_c) \approx 0.108 \text{ meV}$ at the fold. With $T_{\text{eff}} = T^* = 118.7 \text{ mK}$ the Kosterlitz couplings are $K(\alpha = 0, T^*) \approx 46.7$ and $K(\alpha_c, T^*) \approx 10.57$, both far above $K_c = 2/\pi \approx 0.637$. A Qi–Wu–Zhang toy metric (Section 11.6) is a second labeled C1 estimator ($\kappa_{\text{UV}}^{\text{toy}} = 0.425 \text{ meV}$, $K_{\text{fold}}(T^*) = 5.44$); both miss KT at T^* before the fold. A conventional-only variant with $\kappa_{\text{geom}} = 0$ is retained as a labeled alternative; its ultraviolet stiffness is $2\Delta_0/\pi \approx 0.382 \text{ meV}$. The energy of an isolated 2D phase vortex in a system of size L with core cutoff a_c is

$$E_v = \pi\kappa \ln(L/a_c) + E_{\text{core}}. \quad (35)$$

The gap Δ sets a scale for ρ and for $a_c \sim \hbar c_s/\Delta$. BKT physics is a statement about whether κ/T_{eff} flows through the universal jump

$$\kappa(T_{\text{BKT}}^-) = \frac{2T_{\text{BKT}}}{\pi}. \quad (36)$$

A finite Δ_c at the fold is compatible with a still-superfluid phase, with a plasma of 2D phase vortices, or with a rounded quantum crossover; the amplitude map and the stiffness flow decide jointly.

6.3 Coulomb gas and Villain action

The effective theory for 2D phase vortices is the XY/Villain model [13, 17, 18], equivalently the integer Coulomb gas

$$\frac{H}{T_{\text{eff}}} = 2\pi K \sum_{i < j} n_i n_j \ln \frac{|r_i - r_j|}{a_c} + \frac{E_{\text{core}}}{T_{\text{eff}}} \sum_i n_i^2, \quad (37)$$

with spin-wave piece $S_{\text{sw}}[\theta] = \frac{K}{2} \int (\nabla \theta)^2$. We adopt the Kosterlitz normalization $K = \kappa/T_{\text{eff}}$ (not the José–Kadanoff $\tilde{K} = \pi\kappa/T_{\text{eff}}$), so that the one-loop flow

$$\frac{dK^{-1}}{d\ell} = 4\pi^3 y^2, \quad \frac{dy}{d\ell} = (2 - \pi K)y \quad (38)$$

has critical point $K_c = 2/\pi$. Equivalently, $\pi\kappa/T_{\text{eff}} = \pi K$ equals 2 on the universal jump (36). Equation (38) is the vortex-RG sector of the theory. Its derivation from the microscopic TQGL functional proceeds through a Villain reduction, which we state as an ansatz: the phase sector is governed by (37)–(38) once amplitude fluctuations have been integrated at the scale a_c . The flow equations are derived as an effective theory; the Villain reduction from the full TQGL functional is treated as an ansatz. The fugacity is the textbook form $y = \exp(-E_{\text{core}}/T_{\text{eff}})$.

6.4 C2: winding-one core

The defect that enters (35) is a winding-1 radial Gross–Pitaevskii vortex in the moiré plane, not a hopfion and not a Chern–Simons fluxon. Integrating the truncated GP energy on a disk and subtracting the infrared logarithm at the healing-length cutoff $a_c = \xi = 8\text{ nm}$ yields $E_{\text{core}}/\kappa \approx 0.343$, with the cutoff itself an ansatz. That ratio sets the bare fugacity along the branch, $y = \exp[-(E_{\text{core}}/\kappa)K]$. Because E_{core}/κ is $\mathcal{O}(1)$ rather than $\mathcal{O}(10)$, finite y can matter even when the bare coupling still lies well above $2/\pi$: the infrared plasma is a separatrix phenomenon, not merely a $K < K_c$ crossing of the ultraviolet data.

6.5 Competing infrared scales

Two surfaces compete in the (α, K, y) space of a driven TEVC.

Theorem 6.1 (Competing-scale criterion). *On the locked amplitude theory the fold surface is*

$$\alpha = \alpha_c, \quad \Delta = \Delta_c > 0. \quad (39)$$

On the Villain target (38) the KT surface is

$$K = 2/\pi, \quad y \rightarrow 0, \quad (40)$$

with $\Delta \rightarrow 0$ only if Δ tracks a positive power of κ after unbinding. Hypothesis H-VRG1 holds if and only if the physical drive path hits the KT surface before the fold. If $\kappa(A)$ remains larger than κ_{BKT} at $\alpha = \alpha_c$ and finite- y flow stays bound, the fold wins and the gap map retains a finite Δ_c .

Whether a given protocol satisfies it is a property of $\kappa(\alpha)$, E_{core}/κ , and T_{eff} .

Hypothesis 1 (H-VRG1). *Including a 2D phase-vortex fugacity in the RG of $\theta = \arg \Psi$ converts the locked saddle-node fold into a BKT essential singularity $\Delta \sim \exp(-C/\sqrt{A_c - A})$ with continuous gap vanishing, whenever the drive path of Theorem 6.1 reaches the KT surface first.*

H-VRG1 is a conjecture. The C1 stiffness is a computed closure with ansatz inputs, not a Brillouin-zone theorem, so the competing-scale diagram below is the implication of that closure.

6.6 Chern–Simons statistics: truncation or anyon plasma

The parent functional contains a Chern–Simons term at level $k = 2N$, which attaches flux $\Phi_{\text{CS}} = 2\pi/N$ to a 2π winding. Ordinary XY–KT assumes bosonic integer 2D phase vortices. An anyonic plasma changes the Coulomb coupling.

Two consistent options exist. The first, adopted for the default scan and matching the GPE of Section 13, is to *truncate* Chern–Simons, Berry, and Skyrme from the vortex sector: the flow (38) then applies to ordinary 2D phase vortices, and CS fluxons are outside the calculation. The second is to restore level $k = 2N$ and treat 2D phase vortices as anyons with statistical angle

$$\theta_{\text{stat}} = \frac{\pi}{N} \quad (N = 5 \Rightarrow \theta_{\text{stat}} = \pi/5). \quad (41)$$

A Wen–Zee-inspired shift of the KT criterion [19, 20] then reads

$$K_c^{\text{anyon}} = \frac{2}{\pi} \left(1 - \frac{\theta_{\text{stat}}}{\pi}\right)^2 = \frac{2}{\pi} \left(1 - \frac{1}{N}\right)^2, \quad (42)$$

which for $N = 5$ lowers K_c from $2/\pi \approx 0.637$ to ≈ 0.407 . A lower K_c moves the KT surface toward weaker stiffness. Equation (42) remains conjectural unless the anyon–Coulomb–gas algebra is derived from the TQGL Chern–Simons term; it is displayed so that the truncation is explicit, and it is never mixed into the truncated default.

6.7 C3: competing-scale scan

On the primary closure—healing-disk n_0 plus the frozen geometric floor—at $T^* = 118.7$ mK, Kostertiz flow at frozen α remains bound all the way to the fold. The infrared coupling at α_c is $K(\alpha_c) \approx 10.57 \gg 2/\pi$: the fold is reached first. Restoring the labeled anyon threshold (42) does not change the winner on this $\kappa(\alpha)$. The same $E_{\text{core}}/\kappa \approx 0.34$ that would be dangerous near $K \sim 1$ is harmless at $K \sim 10$, because $y = \exp(-0.34K)$ is then exponentially small.

The conventional-only variant ($\kappa_{\text{geom}} = 0$) at the same T^* is different. Bare K still stays above $2/\pi$ through α_c , so there is no ultraviolet $K < K_c$ crossing. Finite- y flow, however, hits the plasma separatrix at $\alpha/\alpha_c \approx 0.945$: KT before the fold, as an infrared phenomenon. The computed core ratio is large enough for that separatrix even though the geometric floor is what had kept the primary branch bound.

Raising T_{eff} opens a temperature window for the vortex sector even on the primary closure. Below ~ 50 mK both closures remain bound out to α_c . At T^* only conventional-only unbinds. At $T_{\text{eff}} = 0.5$ K the primary branch already plasmas at $\alpha/\alpha_c \approx 0.819$; at 1 K the plasma station has moved in to $\alpha/\alpha_c \approx 0.302$. That window is the physically interesting content of the sector: whether a laboratory protocol sits on the fold side or the KT side of Theorem 6.1 is a joint question of stiffness closure and temperature, not an algebraic identity.

6.8 C4: vacuum driving and the essential-singularity implication

On the primary T^* scan the flow stays bound, so the locked finite Δ_c stands: there is no vortex-driven continuous $\Delta \rightarrow 0$ on that closure. When a protocol does pass Theorem 6.1—as the conventional-only variant does at T^* —increasing the vacuum amplitude A (equivalently α) walks the system along $\delta_+(\alpha) \searrow \delta_c$ while κ and E_{core} fall and y rises. Unbinding occurs if the finite- y separatrix, or $K = 2/\pi$, is reached while α is still less than α_c . Identifying the KT correlation length with a gap,

$$\Delta \sim \hbar c_s / \xi_{\text{KT}}, \quad (43)$$




Figure 17: **Kosterlitz flow at frozen α and T^* .** Primary-closure trajectories at $\alpha/\alpha_c = 0.30$, at λ^* , and near the fold all run to $y \rightarrow 0$ with $K \gg 2/\pi$ (bound). A conventional-only station at the finite- y onset $\alpha/\alpha_c \approx 0.945$ runs to a plasma. Option A* holds α fixed; there is no β_α .




Figure 18: **Competing-scale diagram at T^* .** Kosterlitz $K = \kappa/T^*$ along the locked physical branch for the primary C1 closure and the conventional-only variant. The bosonic line $K = 2/\pi$ and the labeled anyon line lie far below the primary curve; the fold is reached first. Conventional-only unbinding is a finite- y infrared plasma at $\alpha/\alpha_c \approx 0.945$, marked, not a bare $K < 2/\pi$ crossing.




Figure 19: **Temperature window of the vortex sector.** First α/α_c at which frozen- α flow reaches an infrared plasma, or $\alpha/\alpha_c = 1$ if the fold is reached first. The Schwinger crossover T^* is marked. Raising T_{eff} to 0.5–1 K lets even the primary closure unbind by finite y .

then produces the essential singularity

$$\Delta(A) \sim \exp(-C/\sqrt{A_c - A}) \quad (44)$$

whenever Δ tracks $\sqrt{\kappa}$ or an equivalent stiffness collapse. Equation (43) is a conjectural identification; Eq. (44) is the BKT implication of the vortex sector (H-VRG1). It is a physical prediction for a protocol that passes Theorem 6.1, not a replacement of the locked amplitude fold.

6.9 Finite-size phase correlations

Define the 2D phase correlator

$$g(r) = \langle e^{i[\theta(\mathbf{r}) - \theta(0)]} \rangle. \quad (45)$$

In a Villain spin-wave treatment (vortices omitted) $g(r) \sim r^{-\eta}$ with $\eta = 1/(2\pi K)$; at $K = 2/\pi$ one has $\eta = 1/4$. In a plasma, $g(r)$ is exponential. An effective exponent on a torus of size L ,

$$\eta_{\text{eff}}(L) = -\frac{\ln g(L/2)}{\ln(L/(2a))}, \quad (46)$$

approaches 1/4 at a putative KT point and rises when vortices proliferate. On the primary T^* branch, $K \sim 10$ –47 implies a spin-wave η of order 10^{-3} – 10^{-2} , far below 1/4, consistent with a still-bound phase. Figure 20 shows that family together with the algebraic KT target and an exponential plasma toy; Fig. 21 shows η_{eff} along the physical branch. A modest $L = 16$ Metropolis XY run remains a program item for sampling vortices rather than spin waves.

6.10 C5: fold versus essential singularity

On the locked physical branch, $\delta - \delta_c \sim \sqrt{t_A}$ with $t_A = (\alpha_c - \alpha)/\alpha_c$ is a theorem of the amplitude map. A Gaussian comparison of $\frac{1}{2} \ln t_A$ against $1/\sqrt{t_A}$ on $\ln(\delta - \delta_c)$ over $\gtrsim 1.5$ decades in t_A returns a free slope 0.542 (versus the theorem value 1/2) and prefers the square root by $\Delta\ell \approx +431$. That comparison is a consistency check on fold data. Vortex-implied Δ from Eq. (43) is not available on the primary T^* scan, which never unbinds, so the essential-singularity curve in Fig. 22 is the H-VRG1 target overlaid on a map that does not contain vortices. The preference for $\sqrt{t_A}$ confirms the locked fold; it is not a statement that a KT surface cannot be hit on a different closure or at a higher T_{eff} .




Figure 20: **Phase correlator $g(r)$** . Villain spin-wave decay at the primary λ^* stiffness, the algebraic KT target $\eta = 1/4$, and an exponential plasma toy. Vortices are not sampled in the spin-wave curves.




Figure 21: **Spin-wave $\eta = 1/(2\pi K(\alpha))$ along the primary T^* branch**. With $K(\alpha_c) \approx 10.57$ the exponent remains far below the KT target $\eta = 1/4$. The line $\eta = 1/4$ is the would-be unbinding marker.




Figure 22: **Amplitude-map scaling on the locked branch**. Physical $\delta(t_A)$ follows $\delta_c + B\sqrt{t_A}$. The dotted essential singularity is the H-VRG1 target of the vortex sector; vortex-implied Δ is not constructed on the primary T^* scan.

6.11 C6: thermal versus quantum unbinding

Classical KT flow requires $T > 0$. The TEVC condensation window is $\lesssim 1$ K; the Schwinger crossover is $T^* = 118.7$ mK; the live GPE of Section 13 is $T = 0$. At strictly zero temperature the thermal fugacity $y = e^{-E_{\text{core}}/T}$ vanishes and textbook KT does not run. Quantum phase slips, the 2D Bose–Hubbard/Villain dual [21], or dissipative Ambegaokar–Halperin–Nelson–Siggia (AHNS) dynamics [22, 23] are then the relevant mechanisms.

Hypothesis 2 (H-VRG2). *Phonon drag in a Gao–Khalaf-type moiré electron–phonon bath places 2D phase vortices in an overdamped regime $\hbar/\tau_v \gtrsim \pi\kappa$, converting a universal jump into a rounded crossover rather than a textbook thermal KT transition.*

Two labeled mobility estimates are compared with the primary $\kappa(\alpha)$ at T^* . The phonon amplitude already present in the truncated GPE is $\hbar/\tau_v \sim g_{\text{ph}} = 0.1\Delta_0$. An Eliashberg-type OOM with $\lambda_{\text{ep}} = 1$ is $\hbar/\tau_v \sim 2\pi\lambda_{\text{ep}}k_B T$. Along the primary branch both give $(\hbar/\tau_v)/(\pi\kappa) \sim 0.04\text{--}0.19 < 1$: the vortices are underdamped. A sharp stiffness jump could therefore survive if a KT surface were hit. H-VRG2 is not forced by these estimates; it remains a conjecture pending a microscopically computed vortex mobility.

6.12 Observables, acceptance, and closures

Table 1: Fold versus vortex-RG observables, as tested by suite C1–C6. Acceptance of H-VRG1 on a given protocol requires the essential-singularity comparison on vortex-inclusive data over $\gtrsim 1.5$ decades in $t_A = (A_c - A)/A_c$, together with $\eta \rightarrow 1/4$ in $g(r)$.

Observable	Fold (established)	Vortex RG (conjectured)	Present suite
$\Delta(A)$	ends at Δ_c ; $\sqrt{t_A}$	$\Delta \rightarrow 0$; $e^{-C/\sqrt{t_A}}$	C4 bound at T^* ; C5 on fold data
κ	need not jump	universal jump (36)	C1 two-sector $\kappa(\alpha)$
$g(r)$	amplitude death first	algebraic $\rightarrow$ exponential	spin-wave at primary K
vortex density	not required	proliferates above A_c	C2 core; C3 plasma flag
mobility	—	jump vs AHNS crossover	C6 underdamped at T^*
specific heat	mean-field anomaly possible	no latent heat	program

The vortex-RG sector is closed by the following statements. The Villain reduction from the TQGL functional is an ansatz. The C1 identity (33) is derived given the healing-disk n_0 and the frozen geometric floor, both of which are ansatzes. Hypothesis H-VRG1 is a conjecture: if the drive path satisfies Theorem 6.1, the gap closing takes the essential-singularity form (44). On the primary closure at T^* the fold is reached first; conventional-only stiffness, and the primary closure at $T_{\text{eff}} \sim 0.5\text{--}1$ K, populate the KT side of the diagram as finite- y plasmas. Chern–Simons remains truncated from the default scan unless Eq. (42) is restored as a labeled option. The locked amplitude fold is the mean-field backbone of the static theory and is independent of whether a given protocol realizes H-VRG1.

7 Phononic Band Structure and Resonance Condition

7.1 Moiré Phonon Dispersion

For MATBG, the moiré superlattice with period $\lambda_M \approx 13$ nm creates a reduced Brillouin zone with wavevector $K_M = 4\pi/(3\lambda_M) \approx 0.32$ nm $^{-1}$. Phonon dispersion includes contributions from

in-plane acoustic phonons (longitudinal $\omega_{\text{LA}}(q) = c_{\text{LA}}q$, transverse $\omega_{\text{TA}}(q) = c_{\text{TA}}q$), out-of-plane flexural modes ($\omega_{\text{ZA}}(q) = \alpha_{\text{ZA}}q^2$), and interlayer shear modes ($\omega_S \sim 10\text{--}30 \text{ cm}^{-1}$). Moiré folding creates replicas at $q \pm nK_M$, and interlayer coupling produces flat phonon bands near the Brillouin zone boundaries through hybridization.

7.2 Resonance Condition

The resonance condition $\omega_0 = 2\Delta/\hbar$ emerges rigorously from analytic continuation of the retarded self-energy. The one-loop Matsubara self-energy is

$$\Sigma(i\omega_n) = g^2 \int \frac{d^2q}{(2\pi)^2} T \sum_m D(\mathbf{q}, i\omega_m) G_0(\mathbf{k} - \mathbf{q}, i\omega_n - i\omega_m), \quad (47)$$

where G_0 is the bare condensate propagator. Summing on Matsubara frequencies and continuing $i\omega_n \rightarrow \omega + i0^+$ yields a threshold singularity:

$$\text{Im } \Sigma^R(\omega) \propto \Theta(\omega - 2\Delta/\hbar) \sqrt{\omega - 2\Delta/\hbar}. \quad (48)$$

Physically, this is the two-particle production threshold: below ω_0 the catalyzon cannot decay into two excitonic quasiparticles, so the imaginary part vanishes; above ω_0 the phase space opens and the imaginary part grows as a square root. The spectral function $A_{\text{cat}}(\mathbf{k}, \omega)$ peaks at $\omega = \omega_0$.

7.3 Brillouin Zone Location of Resonance

For MATBG parameters with working gap $\Delta^* = 0.2324 \text{ meV}$, the resonance frequency is $\omega_0 = 2\Delta^*/\hbar \approx 7.06 \times 10^{11} \text{ rad/s}$, or cyclic $\nu^* = 0.112 \text{ THz}$. Independently, the GPE phonon drive of Section 13 uses the bare gap, $\omega_0 = 2\Delta_0/\hbar = 1.82 \text{ ps}^{-1}$, cyclic $\nu_{\text{GPE}} = 0.290 \text{ THz}$. Graphene longitudinal acoustic phonons at $c_{\text{LA}} = 2.14 \times 10^4 \text{ m/s}$ locate the working-gap resonance at

$$q^* = \frac{\omega_0(\Delta^*)}{c_{\text{LA}}} = \frac{2\Delta^*}{\hbar c_{\text{LA}}} = 0.0330 \text{ nm}^{-1}, \quad (49)$$

with real-space wavelength $\lambda_{\text{res}} = 2\pi/q^* = 190 \text{ nm}$. The ratio $q^*/K_M = 0.102$ (with $K_M = 4\pi/(3\lambda_M) \approx 0.322 \text{ nm}^{-1}$) places the resonance near the Γ point. This is graphene-LA kinematics. The locked package velocity $c_s = 1.06 \times 10^5 \text{ m/s}$ from $\hbar c_s = 0.07 \text{ meV} \cdot \mu\text{m}$ produces a distinct Bogoliubov wavelength $0.95 \mu\text{m}$ at the same Δ^* (Section 10.2). The two lengths are not identified; both remain long compared with λ_M , so the qualitative Γ -point statement stands.

8 The Catalyzon: Vacuum-Dressed Quasiparticle

8.1 Concept and Physical Picture

The catalyzon is the vacuum-dressed collective excitation of the TEVC: a coherent bound state of excitonic quasiparticles hybridized with zero-point fluctuations. Its gap lies below the bare excitonic gap through off-diagonal vacuum-mediated coupling, providing a lower-energy pathway across the Mexican-hat barrier of Section 9. The five-mode Hamiltonian below is a resonant model for that hybridization. Linearization of truncated TQGL about the hopfion (Section 11) yields a single discrete $U(1)$ Goldstone rather than five soft modes, so the meV coupling used in numerics remains the bridge identification $g_0 = \lambda E_{\text{vac}}$.

Locked: $\lambda_{\min} = -1 g_0$ (mult. 4), not $-4g_0$ or $-5g_0$




Figure 23: **Five-mode coupling matrix and eigenvalue spectrum.** (Left) Democratic coupling $G = g_0(J - I)$ with zero diagonal and $\text{Tr } G = 0$. (Right) Exact spectrum: dark/catalyzon subspace $\lambda_{\min} = -g_0$ (multiplicity $N - 1$) and bright mode $\lambda_{\max} = (N - 1)g_0$.

8.2 Five-Mode Resonant Hybridization

The model hybridizes five long-wavelength collective modes at the two-particle threshold $\omega_0 = 2\Delta/\hbar$:

1. **Bogoliubov out-of-phase (amplitude) mode:** Higgs-like oscillation of the condensate magnitude $|\Psi|$. In a BCS-BEC crossover system, this mode corresponds to pair-breaking excitations. Frequency at $k = 0$ is $\omega_H = 2\Delta$ (threshold condition). This mode modulates the local gap $\Delta(\mathbf{r}, t)$ and couples to all other modes through the nonlinear condensate dynamics.
2. **Josephson plasma mode:** Relative-phase oscillation between adjacent TEVC layers, $\theta_j - \theta_{j+1}$ where j labels layers. Frequency $\omega_J = \sqrt{2E_J E_C / \hbar^2}$ where E_J is Josephson coupling and E_C is charging energy. For MATBG at twist $\theta \approx 1.1^\circ$ with $N = 5$ layers, interlayer tunneling produces $E_J \sim 0.1$ meV and capacitive charging gives $E_C \sim 1$ meV, yielding $\omega_J \sim 2\Delta/\hbar$ (near-resonant).
3. **Flat-band plasmon:** Collective density oscillation within the moiré unit cell, $\delta n(\mathbf{r}, t) = n_0 e^{i(\mathbf{q}\cdot\mathbf{r} - \omega t)}$. Flat-band dispersion with bandwidth $W \ll E_F$ produces a plasmon mode $\omega_{\text{pl}}^2 = (4\pi e^2 n_0 / m^*) f_q$ where f_q is a geometric form factor and m^* is the flat-band effective mass. For long-wavelength $q \ll K_M$, $\omega_{\text{pl}} \sim 2\Delta/\hbar$.
4. **Hopfion field mode:** Small deformations of the topological texture, parameterized as $\hat{\mathbf{n}}(\mathbf{r}, t) = \hat{\mathbf{n}}_0(\mathbf{r}) + \delta \hat{\mathbf{n}}(\mathbf{r}, t)$ where $\hat{\mathbf{n}}_0$ is the static hopfion solution. These are analogous to spin-wave excitations in magnets but with nontrivial topology. Restoring forces from $\mathcal{F}_{\text{Skyrme}}$ (higher-derivative stabilization) produce a characteristic frequency $\omega_{\text{hopf}} \sim \sqrt{\kappa_S / (m^* R^2)} \sim 2\Delta/\hbar$ for ring radius $R \sim 50$ nm.
5. **Bogoliubov acoustic (Goldstone) mode:** Phase mode $\theta(\mathbf{r}, t)$ with linear dispersion $\omega = c_s k$ at long wavelength. At $k = 0$ this mode has zero energy (Goldstone mode of broken $U(1)$).

Catalyzon: 5-mode hybridization landscape




Figure 24: **Catalyzon formation.** (*Left*) Schematic collective-mode dispersions meeting at the two-particle threshold. (*Right*) Eigenvalue splitting versus g_0 : four-fold dark multiplet at $\lambda_{\min} = -g_0$ and bright mode at $\lambda_{\max} = 4g_0$ ($N = 5$). Catalyzon hybridization is a parallel channel to the static fold and to truncated GPE dynamics.




Figure 25: **Catalyzon dark-mode basis.** Orthonormal Helmert vectors spanning the sum-zero subspace of the democratic coupling matrix (eigenvalue $\lambda_{\min} = -g_0$).

symmetry). However, at finite $k^* = 2\Delta/(\hbar c_s)$, $\omega(k^*) = 2\Delta/\hbar$, bringing it into resonance with the other four modes.

8.3 Resonance Condition and Mode Degeneracy

In the resonant model the five modes are taken as accidentally degenerate at $\omega_0 = 2\Delta/\hbar$ for the wavevector q^* of Section 7. The degeneracy is not symmetry-protected; it is a working assumption of the five-mode algebra. Linearization of truncated TQGL does not produce five Goldstone modes, as discussed above.

Zero-point fluctuations of the electromagnetic field (Casimir modes with $\mathcal{F}(x) = 0.58068$, Section 4) and phononic field (flat phonon bands, Section 7) introduce vacuum-mediated interactions between these modes. These interactions are off-diagonal: they do not shift individual mode frequencies (no diagonal G_{ii} terms) but instead hybridize the modes through pair-wise couplings G_{ij} for $i \neq j$.

8.4 Coupling Matrix Structure

The five-mode Hamiltonian near ω_0 is

$$H = \sum_{i=1}^5 \hbar\omega_0 a_i^\dagger a_i + \sum_{i<j} G_{ij} (a_i^\dagger a_j + a_j^\dagger a_i), \quad (50)$$

where $a_i^\dagger, a_i$ are creation/annihilation operators for mode i , and $G \in \mathbb{R}^{5 \times 5}$ is the symmetric coupling matrix. By construction:

- $G_{ii} = 0$ (no self-energy corrections at resonance);
- $\text{Tr } G = 0$ (sum rule from charge neutrality and momentum conservation);
- G_{ij} mediated by vacuum field correlators $\langle \phi_{\text{ZPF}}(\mathbf{r}) \phi_{\text{ZPF}}(\mathbf{r}') \rangle$.

Diagonalizing H yields eigenvalues $\hbar\omega_0 + \lambda_i$ where $\{\lambda_i\}_{i=1}^5$ are eigenvalues of G . For the democratic coupling $G = g_0(J_N - I_N)$ the spectrum is exact: $\lambda_{\max} = (N-1)g_0$ (multiplicity 1; bright/in-phase) and $\lambda_{\min} = -g_0$ (multiplicity $N-1$; dark/sum-zero). The catalyzon corresponds to this dark subspace.

Theorem 8.1 (Catalyzon Formation Criterion). *Let $G \in \mathbb{R}^{5 \times 5}$ be symmetric with $G_{ii} = 0$ for all i , $\text{Tr } G = 0$, and $G \neq 0$. Then at least one eigenvalue of G is strictly negative, and the catalyzon gap satisfies*

$$\boxed{\Delta_{\text{cat}} = \Delta_{\text{ex}} + \lambda_{\min} < \Delta_{\text{ex}}.} \quad (51)$$

Proof. Symmetry of G implies eigenvalues $\{\lambda_i\}_{i=1}^5$ are real and $\sum_i \lambda_i = \text{Tr } G = 0$. The Frobenius norm satisfies $\|G\|_F^2 = \sum_{i,j} G_{ij}^2 = \sum_i \lambda_i^2 > 0$ since $G \neq 0$, so not all λ_i vanish. Combined with $\sum_i \lambda_i = 0$ and not all $\lambda_i = 0$, there exist eigenvalues of both signs; in particular $\lambda_{\min} < 0$. $\square$

8.5 Physical Interpretation and Magnitude

The catalyzon eigenvector $|\text{cat}\rangle = \sum_{i=1}^5 c_i |i\rangle$ is a coherent superposition of the five modes with amplitudes c_i determined by diagonalizing G . The negative eigenvalue $\lambda_{\min} < 0$ represents a *binding energy*: vacuum-mediated interactions lower the energy of this particular superposition below the uncoupled resonance.

From the locked vacuum $E_{\text{vac}} = 0.1287 \text{ meV}$ with $g_0 = \lambda^* E_{\text{vac}} = 0.0982 \text{ meV}$, the democratic dark eigenvalue is $|\lambda_{\text{min}}| = g_0$, yielding

$$\Delta_{\text{cat}} = \Delta_{\text{ex}} - g_0^* = 0.502 \text{ meV}. \quad (52)$$

This is a 16.4% reduction from the bare exchange gap $\Delta_{\text{ex}} = 0.600 \text{ meV}$ on the democratic channel, distinct from the static fold reduction at λ^* . At the fold edge the same identification gives $g_0 = 0.109 \text{ meV} = \Delta_c$ (18.2%). The catalyzon thus provides a lower-energy pathway for barrier crossing, parallel to the fold.

8.6 Role in Quantum Vacuum Catalyzation

The catalyzon is the microscopic agent of barrier reduction in QVC. When the system attempts to cross the Mexican-hat potential barrier (Section 9), it can do so by:

1. **Bare thermal activation:** Overcoming barrier Δ_{ex} through thermal fluctuations, with rate $\Gamma_{\text{thermal}} \propto e^{-\Delta_{\text{ex}}/k_B T}$.
2. **Quantum tunneling:** Tunneling through the barrier with WKB suppression $\Gamma_{\text{tunnel}} \propto e^{-S_{\text{inst}}/\hbar}$ where S_{inst} is the instanton action.
3. **Catalyzon-mediated crossing:** Accessing the lower-gap state Δ_{cat} , with enhanced rate $\Gamma_{\text{QVC}} \propto e^{-\Delta_{\text{cat}}/k_B T}$ (thermal) or reduced tunneling action.

At sub-Kelvin temperatures ($T \ll \Delta_{\text{ex}}/k_B \sim 7 \text{ K}$), thermal activation is negligible and quantum tunneling dominates. The catalyzon reduces the effective barrier from Δ_{ex} to Δ_{cat} , exponentially enhancing the tunneling probability.

8.7 Scope of the five-mode model

The five-mode Hamiltonian is a resonant model for $\text{Spec}(G)$, not a Goldstone theorem. Truncated TQGL about the hopfion supplies one discrete Goldstone (Section 11.1). Democratic $\text{Spec}(G)$ is exact algebra at the bridge value g_0^* ; Wigner–Seitz K_0 and Hessian overlaps $G^{(H)}$ are computed kernels, neither equal to $g_0(J - I)$ in meV. A Cartan-torsion origin for G_{ij} (hypothesis H-TOR1) is not evaluated here. Theorem 6.1 still guarantees that any nonzero off-diagonal vacuum coupling of the democratic algebra produces a dark eigenvalue $\lambda_{\text{min}} = -g_0$. That statement concerns $\text{Spec}(G)$, not the number of hopfion soft modes.

9 Topological Ginzburg-Landau Functional

The QVC low-energy field theory is the Topological Ginzburg-Landau (TQGL) functional:

$$\mathcal{F}[\Psi, \mathbf{a}] = \mathcal{F}_{\text{bulk}} + \mathcal{F}_{\text{CS}} + \mathcal{F}_{\text{Berry}} + \mathcal{F}_{\text{ZPF}} + \mathcal{F}_{\text{Skyrme}}, \quad (53)$$

with

$$\mathcal{F}_{\text{bulk}} = \int d^2r \left[\alpha |\Psi|^2 + \frac{\beta}{2} |\Psi|^4 + \kappa |\nabla \Psi|^2 \right], \quad (54)$$

$$\mathcal{F}_{\text{CS}} = \frac{k}{4\pi} \int \mathbf{a} \wedge d\mathbf{a}, \quad k = 2N, \quad (55)$$

$$\mathcal{F}_{\text{Berry}} = \int d^2r \Omega(\mathbf{k}) \cdot \mathbf{j}(\mathbf{r}), \quad (56)$$

$$\mathcal{F}_{\text{ZPF}} = -\lambda \int d^2r A_{\text{ZPF}}(\mathbf{r}) |\Psi(\mathbf{r})|^2, \quad (57)$$

$$\mathcal{F}_{\text{Skyrme}} = \kappa_S \int d^2r |\nabla^2 \Psi|^2. \quad (58)$$




Figure 26: **Mexican-hat potential reshaping (3D visualization).** (*Left*) Bare quadratic-quartic $V(\rho) = \alpha_{\text{GL}}\rho^2 + (\beta/2)\rho^4$ of the Track A family (Section 10). (*Center*) The same potential with Landau mass tracking the catalyzon gap $\Delta_{\text{cat}} = 0.502 \text{ meV}$. (*Right*) Radial cut: well-depth reduction $B_{\text{cat}}/B_0 = (\Delta_{\text{cat}}/\Delta_0)^2$ is 30% from the matched $V(\rho)$, while the parallel catalyzon pathway is the 16.4% gap reduction $\Delta_{\text{ex}} \rightarrow \Delta_{\text{cat}}$.

$\mathcal{F}_{\text{bulk}}$ is the conventional Ginzburg-Landau Mexican-hat potential fixing the condensate amplitude (Fig. 26). $\mathcal{F}_{\text{CS}}$ is a Chern-Simons term for the emergent gauge field $\mathbf{a}$ coupled to the condensate phase; at locked level $k = 2N$ the *dictionary* Hall response of the uncatalyzed state is $\sigma_{xy} = (k/2)e^2/h = Ne^2/h$, with step (75); this is not the unique Wen/Laughlin evaluation $\sigma_{xy} = ke^2/h$ of $\frac{k}{4\pi} \int \mathbf{a} \wedge d\mathbf{a}$ (Remark 10.1)[24, 20]. $\mathcal{F}_{\text{Berry}}$ threads the band Berry curvature through the condensate, ensuring that the topology of the underlying electrons is inherited by the TEVC order parameter[25, 26]. $\mathcal{F}_{\text{ZPF}}$ encodes the vacuum coupling explicitly: where the local zero-point amplitude is large, the condensate is favored. Finally, the higher-derivative Skyrme term $\mathcal{F}_{\text{Skyrme}}$ stabilizes finite-size topological textures against Derrick collapse[27, 28]. The matched numerical values of α_{GL} , β , κ_{GL} , λ^* , g_0 , and the Hall step are collected in Section 10.

10 Matched L1 coefficients, lengths, and catalyzon scale

The preceding sections introduce the frozen-cavity bridge, the democratic catalyzon, and the bulk TQGL functional. Those objects share symbols—lengths, Landau coefficients, a vacuum coupling

g_0 , a Hall step, a retardation factor, and an analogue horizon radius—that must not be identified across distinct geometric settings. This section records one convention for each quantity. Fold, catalyzon, and truncated Gross–Pitaevskii dynamics remain independent channels. Holographic and torsion constructions are not used to derive G_{ij} or κ_{GL} . The matching $T_H = T^*$ (hypothesis H-HAWK1) fixes R_{WS} .

10.1 Length dictionary

The frozen-cavity geometry identifies the fat-torus tube with the Ginzburg–Landau healing length,

$$a_H = r = 8 \text{ nm} \quad (59)$$

A second length is the phonon (Bogoliubov) healing constructed from the same package constants that enter the gap map,

$$\xi_{\text{ph}} = \frac{\hbar c_s}{\Delta_0} = 117 \text{ nm} \quad (60)$$

These two lengths are not interchangeable: a_H is the cavity minor radius and the GPE core scale; ξ_{ph} is the scale on which Δ_0 and $\hbar c_s$ produce a dimensionless ratio of order one.

Retardation is defined from the causal phonon propagator by the algebraic identity

$$\eta_{\text{ret}} = \frac{2\Delta_0\xi}{\hbar c_s} \quad (61)$$

Substituting the two lengths of this dictionary yields the pair

$$\eta_{\text{ret}}(a_H) = 0.137, \quad \eta_{\text{ret}}(\xi_{\text{ph}}) = 2. \quad (62)$$

The second evaluation is an identity: $\xi_{\text{ph}} = \hbar c_s / \Delta_0$ converts Eq. (61) into $\eta_{\text{ret}} = 2$. An alternate coherence length $\xi = 181 \text{ nm}$ produces $\eta_{\text{ret}} \approx 3.1$. Table 2 records the three evaluations.

Table 2: Length dictionary on the frozen cavity. The retardation identity is Eq. (61).

Length	Value	η_{ret}	St.
$a_H = r$ (GL/GPE)	8 nm	0.137	L
$\xi_{\text{ph}} = \hbar c_s / \Delta_0$	117 nm	2	D
ξ (alternate)	181 nm	3.1	Alt

10.2 Locked kinematics: mass, frequencies, two velocities

The gradient weight $\kappa_{\text{GL}} = \Delta_0 a_H^2$ is $\hbar^2 / (2m^*)$, so

$$\frac{m^*}{m_e} = \frac{\hbar^2 / (2m_e)}{\kappa_{\text{GL}}} = 0.99 \quad (63)$$

The healing identification $a_H = \hbar / \sqrt{2m^*\Delta_0}$ with locked $a_H = 8 \text{ nm}$ therefore requires $m^* \approx m_e$, not $0.4 m_e$.

Package $\hbar c_s = 0.07 \text{ meV} \cdot \mu\text{m}$ converts to $c_s = 1.06 \times 10^5 \text{ m/s}$. This velocity enters the Casimir prefactor, the gap map, retardation (61), and the Goldstone resonance $k^* = 2\Delta / (\hbar c_s)$. Independently, graphene’s longitudinal acoustic speed is $c_{\text{LA}} = 2.14 \times 10^4 \text{ m/s}$ and locates the moiré-zone wavevector $q^* = \omega_0 / c_{\text{LA}}$. The two speeds are not identified.

Cyclic frequencies $\nu = \omega/(2\pi)$ at $\omega = 2\Delta/\hbar$ are $\nu_{\text{GPE}} = 0.290$ THz for the GPE drive $\Delta = \Delta_0$ and $\nu^* = 0.112$ THz for the working gap Δ^* . Angular $\omega_0(\Delta_0) = 1.82$ ps $^{-1}$. At Δ^* the graphene-LA kinematics give $q^* = 0.0330$ nm $^{-1}$, $\lambda_{\text{res}} = 2\pi/q^* = 190$ nm, and $q^*/K_M = 0.102$ with $K_M = 4\pi/(3\lambda_M)$. The Bogoliubov wavelength at the same Δ^* is 0.95 μm . The Vakulenko–Kapitansky comparison uses E_{vac} : $N^{-3/4} \approx 0.299$ implies $C > 0.430$ meV.

10.3 Track A Ginzburg–Landau coefficients

The bulk functional of Section 9 and the two-sector split of Section 6 employ a Landau mass α_{GL} , a quartic β , and a gradient weight κ_{GL} , together with a participation λ that feeds the gap-map coupling α of Section 5. These are distinct objects. The gap-map coupling is dimensionless,

$$\alpha = \lambda \frac{E_{\text{vac}}}{\Delta_0}, \quad (64)$$

and at the working point equals $\alpha^* = 0.1636$. The Landau mass is matched to the bare exchange gap of the truncated GPE,

$$\alpha_{\text{GL}} = -\Delta_0 = -0.600 \text{ meV} \quad (65)$$

so that $|\alpha_{\text{GL}}| = \Delta_0$ and α_{GL} is not the number α^* . The gradient weight follows from the healing identification $a_H = r$ in the kinetic coefficient $\hbar^2/(2m^*)$,

$$\kappa_{\text{GL}} = \Delta_0 a_H^2 = 3.84 \times 10^{-5} \text{ meV} \cdot \mu\text{m}^2 \quad (66)$$

Hence $m^* = \hbar^2/(2\kappa_{\text{GL}}) = 0.99 m_e$. This κ_{GL} is not the holographic γ_{QM} of Section 27; that coefficient is an L3 matching and is not imported into Track A.

The quartic is matched so that the Mexican-hat vacuum manifold sits on the hopfion density of the truncated solver. On the ring-weighted initial condition of `hopfion_initial` (box $L = 200$ nm, $n_0 = 1$, core depletion 0.10, noise-free) one finds $n_{\text{ref}} = 0.864$, and therefore

$$\beta = \frac{\Delta_0}{n_{\text{ref}}} = 0.694 \text{ meV} \quad (67)$$

The GPE cubic-quintic potential used in the 1 ps integrator is the related but not identical pair $\alpha_{\text{nl}} = -\Delta_0$, $\beta_{\text{nl}} = \Delta_0/n_0$ with dimensionless $n_0 = 1$; n_{ref} is the live-gap denominator of Section 13. Participation in the dictionary is the working value $\lambda = \lambda^* = 0.7628$, with $\lambda_{\text{fold}} = 0.8475$ the edge of the physical fold branch.

The Skyrme coefficient that stabilizes finite-size textures against Derrick collapse is not derived from a microscopic Hubbard model. A Faddeev–Skyrme balance at the locked tube scale a_H gives the estimate $\kappa_S \sim \kappa_{\text{GL}} a_H^2 = \Delta_0 a_H^4 = 2.46 \times 10^{-9} \text{ meV} \cdot \mu\text{m}^4$. Table 3 collects the Track A card.

10.4 Catalyzon scale $g_0 = \lambda E_{\text{vac}}$

If the same vacuum that feeds the fold also hybridizes the five long-wavelength modes of Section 8, the off-diagonal coupling is the bridge identification

$$g_0 = \lambda E_{\text{vac}}. \quad (68)$$

At the working point λ^* this is $g_0^* = 0.0982$ meV, so the democratic dark eigenvalue $\lambda_{\text{min}} = -g_0$ produces

$$\Delta_{\text{cat}} = \Delta_{\text{ex}} - g_0^* = 0.502 \text{ meV} \quad (69)$$

(16.4% of $\Delta_{\text{ex}} = 0.600$ meV). At the fold edge λ_{fold} one has $g_0 = \lambda_{\text{fold}} E_{\text{vac}} = 0.109$ meV $= \Delta_c$, a 18.2% catalyzon reduction. Both evaluations are parallel to the static fold: $\Delta^* = 0.2324$ meV at λ^* is an amplitude-map number, not a stacked correction to Δ_{cat} . The working vacuum scale is g_0^* ; a coupling larger than E_{vac} is not a participation of the locked vacuum.

Table 3: Track A Ginzburg–Landau coefficients. The gap-map α is dimensionless; α_{GL} and β are energies. Holographic γ_{QM} is not κ_{GL} .

Symbol	Value	Meaning	St.
α^*	0.1636	Gap-map $\lambda^* E_{\text{vac}}/\Delta_0$	D
α_{GL}	-0.600 meV	Landau mass, $ \alpha_{\text{GL}} = \Delta_0$	M
κ_{GL}	$3.84 \times 10^{-5} \text{ meV} \cdot \mu\text{m}^2$	$\Delta_0 a_H^2$ with $a_H = r$	D
m^*/m_e	0.99	$\hbar^2/(2\kappa_{\text{GL}})$	D
λ^*	0.7628	Working participation	L
β	0.694 meV	Δ_0/n_{ref} on hopfion IC	M
n_{ref}	0.864	Ring-weighted hopfion density	C
κ_S	$2.46 \times 10^{-9} \text{ meV} \cdot \mu\text{m}^4$	Derrick estimate at tube scale	A

10.5 Mexican-hat barrier from $V(\rho)$

The bulk potential of Track A is the quadratic-quartic family

$$V(\rho) = \alpha_{\text{GL}} \rho^2 + \frac{\beta}{2} \rho^4, \quad (70)$$

with α_{GL} and β from Table 3. The vacuum manifold sits at $\rho^2 = |\alpha_{\text{GL}}|/\beta = n_{\text{ref}}$, and the well depth measured from $\rho = 0$ is

$$B = \frac{\alpha_{\text{GL}}^2}{2\beta} = \frac{\Delta_0 n_{\text{ref}}}{2}. \quad (71)$$

On the catalyzon channel the Landau mass tracks the dressed gap, $\alpha_{\text{GL}} \rightarrow -\Delta_{\text{cat}}$, at the same β . The well-depth ratio is then independent of n_{ref} ,

$$\frac{B_{\text{cat}}}{B_0} = \left(\frac{\Delta_{\text{cat}}}{\Delta_0} \right)^2, \quad (72)$$

which evaluates to a 30% reduction. Figure 27 shows the radial cuts of Eq. (70) at $\alpha_{\text{GL}} = -\Delta_0$ and $\alpha_{\text{GL}} = -\Delta_{\text{cat}}$. This is the barrier of the matched potential, not a schematic pair of Landau masses.




Figure 27: **Matched Mexican-hat radial cuts.** Equation (70) in units of Δ_0 , with the Track A quartic β scaled out of the dimensionless plot. Well-depth reduction $B_{\text{cat}}/B_0 = (\Delta_{\text{cat}}/\Delta_0)^2$ is 30%. The catalyzon pathway $\Delta_{\text{ex}} \rightarrow \Delta_{\text{cat}}$ remains the parallel 16.4% gap reduction of Section 10.4.

10.6 Hall convention

A single Chern–Simons *dictionary* is used throughout. The topological term in the TQGL functional is

$$\mathcal{F}_{\text{CS}} = \frac{k}{4\pi} \int \mathbf{a} \wedge d\mathbf{a}, \quad k = 2N, \quad (73)$$

and the lens-space identification of Section 26 is $Q_H = 1/N$. These two statements are independent locks; their product $kQ_H = 2$ is algebra, not a pairing theorem that uniquely determines k from Q_H or conversely. The uncatalyzed Hall conductance of the condensate is

$$\sigma_{xy} = N \frac{e^2}{h} = \frac{k}{2} \frac{e^2}{h}. \quad (74)$$

Each hopfion nucleation event changes that conductance by one step

$$\delta\sigma_{xy} = \frac{e^2}{2Nh} = \frac{Q_H e^2}{2h} = \frac{e^2}{kh}. \quad (75)$$

For $N = 5$ this is $\delta\sigma_{xy} = 3.87 \times 10^{-6} \text{ S}$. No second Hall formula is employed as a physics lock.

Remark 10.1 (Dictionary versus textbook Chern–Simons). *The usual continuum evaluation of $\frac{k}{4\pi} \int \mathbf{a} \wedge d\mathbf{a}$ assigns Hall conductance $\sigma_{xy} = k e^2/h$ [24, 20]. At the locked level $k = 2N$ that textbook reading would be $2N e^2/h$, which is not the QVC lock. The paper uses $\sigma_{xy} = (k/2)e^2/h = N e^2/h$ as a dictionary convention so that the uncatalyzed condensate tracks N layers and the step (75) remains $e^2/(2Nh)$. Likewise, textbook flux attachment at level k is $\oint \mathbf{a} = 2\pi/k = \pi/N$; the mean-field background integrated in Section 11.5 is $a_\theta = Q_H/\rho$, whose holonomy is $2\pi Q_H = 2\pi/N$. Both numbers are recorded. Neither is used to reset k , Q_H , σ_{xy} , or $\delta\sigma_{xy}$. The covariant stepper confirms the mean-field holonomy; it does not transport an edge and does not independently derive the Hall staircase.*

10.7 Analogue horizon radius from $T_H = T^*$

The Unruh–Visser acoustic metric of Section 28 has surface gravity $\kappa_{\text{surface}} = 2c_s/R_{\text{WS}}$ and Hawking temperature $T_H = \hbar c_s/(\pi k_B R_{\text{WS}})$. Hypothesis H-HAWK1 is the matching condition $T_H = T^*$, which inverts for the Wigner–Seitz radius

$$R_{\text{WS}} = \frac{\hbar c_s}{\pi k_B T^*} = 2.18 \mu\text{m} \quad (76)$$

using the Schwinger diagnostic $T^* = 118.7 \text{ mK}$ constructed from $E_{\text{pair}} = 0.1963 \text{ meV}$ at g_0^* . The same algebra at $T^* = 285 \text{ mK}$ returns $R_{\text{WS}} = 0.91 \mu\text{m}$; that length is the matching at that temperature, not the present T^* . The Berry Monte Carlo ratio $R_{\text{mean}} = 0.908$ of Section 21 is a dimensionless GUE/GOE comparison of $|\lambda_{\min}|$, not a radius, and is not R_{WS} .

10.8 Wigner–Seitz kernel G_{ij}

Democratic $G = g_0(J_N - I_N)$ has the exact spectrum $\lambda_{\max} = (N-1)g_0$ (multiplicity 1) and $\lambda_{\min} = -g_0$ (multiplicity $N-1$) at the bridge scale g_0^* . That algebra does not evaluate matrix elements and is not inserted into G_{ij} . Two computed kernels share the Macdonald function $K_0(|\mathbf{r} - \mathbf{r}'|/\xi_K)$ with $\xi_K = R = 50 \text{ nm}$ and set $G_{ii} = 0$.

The Gaussian construction places five profiles of width $\sigma \approx 0.18 \lambda_M$ on a regular pentagon inside the moiré Wigner–Seitz hexagon. The output is Hermitian and traceless, with $\lambda_{\min} < 0$,

mean off-diagonal $g_{0,\text{eff}} = 1.365 \times 10^{-4}$ in kernel units, and relative Frobenius distance 0.0667 from the democratic matrix at $\langle G_{\text{off}} \rangle$. The eigenvalues are

$$\text{Spec}(G_{K_0}) \approx (-1.568, -1.568, -1.162, -1.160, +5.458) \times 10^{-4} : \quad (77)$$

λ_{max} tracks $4g_{0,\text{eff}}$, while the dark subspace is split by the pentagon-in-hexagon geometry rather than exactly four-fold. This $g_{0,\text{eff}}$ is not the bridge energy $g_0^* = 0.0982 \text{ meV}$.

A ring-Fourier companion uses angular harmonics of the hopfion envelope, Wannier-transformed to N localized packets on the locked torus. It returns $g_{0,\text{eff}} = 5.472 \times 10^{-4}$ in kernel units and relative Frobenius distance 0.1564, with

$$\text{Spec}(G_{\text{ring}}) \approx (-7.385, -7.247, -4.326, -2.930, +21.888) \times 10^{-4}. \quad (78)$$

Converting either $g_{0,\text{eff}}$ to meV requires a prefactor that is not supplied by the overlap; the meV scale of the catalyzon remains the bridge value g_0^* . Neither kernel is a torsion vertex (hypothesis H-TOR1 is left for a future three-dimensional Weitzenböck quadrature; Section 11) nor a hand-inserted $J - I$.

10.9 Lindblad window

The truncated GPE of Section 13 is kinetic plus cubic-quintic plus V_{ZPF} plus V_{phonon} ; Chern–Simons, Berry, and Skyrme terms are not discretised. A Lindblad dissipator is applied after each split step with phenomenological rates $\gamma_n = \gamma_\phi = 1/\tau_{\text{dec}}$. On the coherent 1 ps coupled-DCE window the integrator returns $\Delta_{\text{live}}(1 \text{ ps}) = 0.6305 \text{ meV}$ and $g^{(1)} = 0.8832$. Combined number-plus-dephasing on that window is Table 4.

Table 4: Truncated GPE plus Lindblad on the 1 ps coupled-DCE window. Combined scan: $\gamma_n = \gamma_\phi = 1/\tau_{\text{dec}}$.

τ_{dec}	Δ_{live}	$g^{(1)}$	St.
coherent	0.6305 meV	0.8832	C
5 ps	0.5720 meV	0.7102	C
20 ps	0.6152 meV	0.8351	C
50 ps	0.6248 meV	0.8639	C

Number-only damping at $\tau_{\text{dec}} = 5 \text{ ps}$ gives $\Delta_{\text{live}} = 0.5697 \text{ meV}$ and $g^{(1)} = 0.8806$; dephasing-only gives $\Delta_{\text{live}} = 0.6309 \text{ meV}$ and $g^{(1)} = 0.7031$. Number damping moves Δ_{live} ; dephasing moves $g^{(1)}$. A 20 ps window with frozen V_{ZPF} (no DCE) isolates the rates: coherent $\Delta_{\text{live}} = 0.6623 \text{ meV}$, $g^{(1)} = 0.1075$; at $\tau_{\text{dec}} = 5, 20, 50 \text{ ps}$ the pairs are (0.0872 meV, 0.0102), (0.3920 meV, 0.0812), and (0.4761 meV, 0.0032). The $\tau_{\text{dec}} = 5 \text{ ps}$ frozen trajectory reaches 0.0872 meV, below $\Delta_c = 0.109 \text{ meV}$. That window is unphysical on the static branch and is not combined with Δ^* or Δ_{cat} .

11 Catalyzon linearization and L1 closures

The dictionary of Section 10 fixes the meV scale of the catalyzon by $g_0^* = \lambda^* E_{\text{vac}}$. This section linearizes truncated TQGL about the hopfion and records additional algebraic closures: a Chern–Simons covariant kinetic equation, a two-band quantum-metric stiffness on the physical fold branch, a Hubbard-scale matching, a Villain dualization with a finite-lattice $g(r)$, and the S^3 Hopf–Whitehead integral. A Cartan-torsion origin of G_{ij} is not evaluated.

11.1 Galerkin linearization about the hopfion

The truncated energy whose variation reproduces the Gross–Pitaevskii equation of Section 13 is

$$F[\psi] = \int d^2r \left[\kappa_{\text{GL}} |\nabla \psi|^2 + \frac{1}{2} \alpha_{\text{nl}} n^2 + \frac{1}{3} \beta_{\text{nl}} n^3 + V_{\text{ZPF}} n \right], \quad (79)$$

with $n = |\psi|^2$, $\alpha_{\text{nl}} = -\Delta_0$, $\beta_{\text{nl}} = \Delta_0$, $\kappa_{\text{GL}} = \Delta_0 a_H^2$, and $V_{\text{ZPF}} = -E_{\text{vac}}$ times the ring Gaussian. Chern–Simons, Berry, and Skyrme terms remain outside F , matching the stepper. A noise-free hopfion ψ_0 of Section 13 is the background. Infinitesimal $U(1)$ rotations $\psi \rightarrow e^{i\varepsilon} \psi$ leave (79) invariant, so $\Phi_{\text{G}} = i\psi_0$ is the Goldstone direction. A Galerkin basis orthonormalizes that Goldstone together with amplitude and phase packets $A(r)e^{in\theta}$ and $i\psi_0 e^{in\theta}$ for $n = 0, \dots, 4$ on a 48^2 periodic box.

The real Hessian $H_{ij} = \partial^2 F / \partial x_i \partial x_j$ along $\psi = \psi_0 + \sum x_k \Phi_k$ is obtained by mixed central differences. Its spectrum on the locked freeze is

$$\text{Spec}(H) \approx (0.052, 0.720, 1.32, 2.05, 2.37, 2.69, 3.01, 3.25, 6.00, 7.32) \text{ meV} \quad (80)$$

The lowest eigenvalue is the discrete Goldstone, lifted by the grid and the finite-difference step to 0.052 meV. The remaining nine directions are gapped by $\mathcal{O}(1)$ meV. The truncated functional about a pinned ring therefore supports a single $U(1)$ Goldstone, not a five-fold Goldstone manifold. Figure 28 shows the stick spectrum.




Figure 28: **Galerkin Hessian of truncated TQGL about the hopfion.** The isolated lowest eigenvalue is the discrete $U(1)$ Goldstone; the truncation does not produce five soft modes.

Off-diagonal entries of the amplitude block of H , with vanishing diagonal as in the catalyzon convention, define a Hessian coupling $G^{(H)}$ in meV. The mean off-diagonal is $g_{0,\text{eff}}^{(H)} = 0.0185 \text{ meV}$, of the same order as the bridge g_0^* but not equal to it (ratio ≈ 0.19). Relative Frobenius distance to $g_0(J - I)$ at that mean is large: the Hessian coupling is not democratic. The meV catalyzon gap therefore remains $\Delta_{\text{cat}} = \Delta_0 - g_0^*$ from Section 10; $G^{(H)}$ is a fluctuation pattern, not a replacement of the bridge.

11.2 Dark versus bright envelopes (H-CAT1)

On the Galerkin frequencies ω_k of (80), a radiation kernel $\gamma_k = \gamma_0(\omega_k/\omega_{\text{ref}})^2$ with $\gamma_0 = 0.05 \text{ ps}^{-1}$ assigns a larger decay rate to the stiffest mode. Evolving $c_k(t) = c_k(0)e^{-\gamma_k t}$ yields envelope times $\tau_{\text{dark}} = 662 \text{ ps}$ and $\tau_{\text{bright}} = 10.1 \text{ ps}$ (ratio $\tau_{\text{dark}}/\tau_{\text{bright}} \approx 65$). Under that kernel the Helmert intuition of hypothesis H-CAT1—that dark sum-zero weight outlives the bright eigenvalue—is realized. The kernel is an ansatz: uniform γ would not distinguish the subspaces, and the existing GPE Lindblad of Section 10.9 acts on ψ , not on a mode basis. Figure 29 shows the envelopes.




Figure 29: **H-CAT1 envelopes.** Dark weight (smallest $|\omega|$) versus the brightest Galerkin mode with $\gamma \propto \omega^2$. The ratio of $1/e$ times is 65 under that rate kernel.

11.3 Torsion proxy (H-TOR1)

Hypothesis H-TOR1—that democratic G_{ij} arises as a Cartan torsion overlap on hopfion modes—is not accepted on the present evidence and is parked as an ansatz for future development. A two-dimensional proxy using the phase Jacobian of ψ_0 as a Hopf-density stand-in q_H and a weight $T = |\nabla q_H|$ produces overlaps $G_{ij} = \int \Phi_i^* T \Phi_j$ that are not near-democratic. A three-dimensional Weitzenböck quadrature on true hopfion torsion is not performed. Democratic $\text{Spec}(G)$ remains locked algebra; the Wigner–Seitz K_0 kernel of Section 10.8 remains a distinct computed overlap; the bridge g_0^* is independent of this proxy. Figure 30 records the parked matrix.

11.4 Catalyzon spectral function

The mode spectral density

$$A_{\text{cat}}(\omega) = \sum_k |c_k|^2 \frac{\Gamma_k/\pi}{(\omega - \omega_k)^2 + \Gamma_k^2} \quad (81)$$

with $\Gamma_k = \hbar\gamma_k$ from the H-CAT1 kernel is shown in Fig. 31. It is the Galerkin-pole mix, not a tunneling lineshape. The two-particle phonon threshold $\omega_0 = 2\Delta/\hbar$ of Section 7 remains a distinct energy from these Hessian poles.




Figure 30: **H-TOR1 two-dimensional proxy G_{ij} , parked.** The pattern is not $g_0(J - I)$. A 3D Weitzenböck quadrature is left for future development and is not used to set G_{ij} or g_0^* .




Figure 31: **Catalyzon mode spectral function.** Lorentzian mix of Galerkin Hessian poles. The bridge gap Δ_{cat} is not read from this plot.

11.5 Chern–Simons constraint equation of motion

Level $k = 2N$ is the dictionary lock. The mean-field constraint integrated here is the static connection $a_\theta = Q_H/\rho$, whose holonomy is $\Phi = 2\pi Q_H = 2\pi/N$. Textbook attachment $2\pi/k = \pi/N$ is a different convention and is not used to reset k or Q_H (Remark 10.1). The truncated kinetic operator $-\kappa\nabla^2$ is replaced by the covariant operator

$$H_{\text{kin}} = \kappa(-i\nabla - a)^2 = \kappa[-\nabla^2 + 2ia \cdot \nabla + i(\nabla \cdot a) + |a|^2]. \quad (82)$$

A Strang split advances half a covariant kinetic RK4 step, the local potential, then the other kinetic half-step, so a sits inside the time stepper. On the locked torus the enclosed flux is $2\pi Q_H$ to relative error 3.8×10^{-4} . In polar continuum, $\nabla\theta = Q_H\hat{e}_\theta/\rho$ for the hopfion phase, so the superfluid velocity $\nabla\theta - a$ vanishes identically away from the origin cell (analytic residual 3.8×10^{-16}). That is flux attachment. A ψ -based current on the fractional winding is polluted by the $2\pi Q_H$ branch cut of $\arg\psi$ and is not used as a Hall diagnostic. Mass in the covariant run drifts by 8.3×10^{-6} . The Hall step remains the dictionary algebra $\delta\sigma_{xy} = e^2/(2Nh)$ of Section 10; the uncatalyzed conductance is $\sigma_{xy} = Ne^2/h$. No edge is transported. The anyon shift $K_c^{\text{anyon}} = (2/\pi)(1 - 1/N)^2 = 0.407$ stays the labeled option of Section 6, not a derivation from (82).

11.6 Quantum-metric stiffness on the physical branch

The two-sector identity

$$\kappa(\alpha) = 2\kappa_{\text{GL}}n_0[\delta_+(\alpha)]^2 + \kappa_{\text{geom}} \quad (83)$$

is derived given a density unit n_0 and a geometric weight κ_{geom} . Option A* fixes $\kappa_{\text{GL}} = \Delta_0 a_H^2$. The healing-disk $n_0 = 1/(\pi a_H^2)$ remains ansatz and implies $\kappa_{\text{conv}}(\alpha = 0) = 2\Delta_0/\pi = 0.382 \text{ meV}$. A uniform phase-twist check of that conventional piece on a periodic box agrees with (83) to machine precision.

A Qi–Wu–Zhang two-band Chern insulator at mass $m = -1$ on a 48×48 Brillouin-zone grid returns Chern number $C = 0.993$ and a Peotta–Törmä geometric weight $\kappa_{\text{geom}}^{\text{QWZ}} = 0.0430 \text{ meV}$ from the Brillouin-zone mean of $\text{tr} g_{ij}$. This is a *labeled C1 estimator*, not a replacement of the vortex-sector primary floor $\kappa_{\text{geom}} = \Delta_0/(2\pi)$ of Eq. (34) (Section 6), whose ultraviolet values are $\kappa_{\text{UV}} \approx 0.477 \text{ meV}$ and $K(\alpha_c, T^*) \approx 10.57$. The previous frozen floor in this section was $\Delta_0/(2\pi) = 0.0955 \text{ meV}$. Adding the toy metric to the conventional ultraviolet piece gives $\kappa_{\text{UV}}^{\text{toy}} = 0.425 \text{ meV}$. The toy is not a magic-angle continuum integral.

Scanning (83) along the locked physical fold branch $\delta_+(\alpha)$ with frozen toy geometry yields $\kappa(\alpha = 0) = 0.425 \text{ meV}$ and $\kappa(\alpha \rightarrow \alpha_c) = 0.0556 \text{ meV}$ (Fig. 32). The Kosterlitz coupling $K = \kappa/T_{\text{eff}}$ at the twin-channel T^* is $K_{\text{UV}} = 41.6$ and $K_{\text{fold}} = 5.44$, both above $K_c = 2/\pi$. There is no KT hit before the fold at T^* on this C1 estimator. A tracks-gap variant $\kappa_{\text{geom}} \propto \delta_+$ and a conventional-only variant likewise remain above K_c at T^* . That miss does not disprove hypothesis H-VRG1 as a vortex-RG implication; it means the present n_0 plus QWZ toy do not by themselves force unbinding before α_c .

11.7 Track B Hubbard-scale matching

Track B does not replace Track A and does not perform a Bistritzer–MacDonald Schrieffer–Wolff projection. It records the unique dimensional identifications from the locked GL/GPE scales. The healing identification $t_{\text{heal}} \equiv \kappa_{\text{GL}}/a_H^2 = \Delta_0$ forces $t_{\text{heal}}/U = 1$ when the on-site scale is matched to the gap, $U \leftrightarrow \Delta_0$. If the tight-binding lattice constant is taken to be the moiré period $\lambda_M = 13 \text{ nm}$ instead,

$$t_\lambda = \kappa_{\text{GL}}/\lambda_M^2 = \Delta_0 (a_H/\lambda_M)^2 = 0.227 \text{ meV}, \quad (84)$$




Figure 32: **C1 physical-branch stiffness.** $\kappa(\alpha) = 2\kappa_{\text{GL}}n_0[\delta_+]^2 + \kappa_{\text{geom}}^{\text{QWZ}}$ with frozen toy geometry. The healing-disk n_0 remains ansatz.

so $t_\lambda/U = 0.379$. A bandwidth proxy $W = \hbar c_s/\lambda_M = 5.38 \text{ meV}$ uses the locked phonon speed as a moiré-velocity stand-in and gives $U/W = 0.111$ at the gap scale. That ratio is not the Coulomb $U/W \rightarrow \infty$ of magic-angle graphene. The holographic γ_{QM} is not imported.

11.8 Villain dualization and $g(r)$

Polar substitution $\Psi = \rho e^{i\theta}$ in (79) with ρ frozen at the saddle produces an XY stiffness κ . The Villain approximation replaces $\exp(K \cos \varphi)$ by a 2π -periodic Gaussian $\sum_{n \in \mathbb{Z}} \exp(-(K_V/2)(\varphi - 2\pi n)^2)$ with $K_V \sim K$ at large K . Poisson summation dualizes integer link variables to a height field whose curl is the integer vortex gas

$$\frac{H}{T} = 2\pi K \sum_{i < j} m_i m_j \ln(|r_i - r_j|/a_c) + \frac{E_{\text{core}}}{T} \sum_i m_i^2, \quad (85)$$

with Kosterlitz $K = \kappa/T_{\text{eff}}$ and $K_c = 2/\pi$ as an effective theory and as a property of TEVC. The defect is a 2D phase vortex, not a hopfion and not a Chern–Simons fluxon. Spin-wave theory on the XY magnet gives $\eta_{\text{sw}} = 1/(2\pi K)$, hence $\eta = 1/4$ at the jump.

Checkerboard Metropolis XY on an $L = 20$ torus at a large- K coupling (capped at $K = 3$, since the physical $K(T^*) = 41.6$ lies far into the spin-wave regime) returns $g(r)$ with log-fit exponent $\eta_{\text{UV}} = 0.050$ against $\eta_{\text{sw}} = 0.053$. At $K = 2/\pi$ the same finite lattice is not a thermodynamic-limit measurement of $\eta \rightarrow 1/4$. Figure 33 records the two correlators; Fig. 34 records $\eta(K)$ against the spin-wave line. Hypothesis H-VRG1 is not upgraded.

11.9 Hopf invariant

The locked charge $Q_H = 1/N$ is Chern–Simons/linking on $L(N, 1)$. Independently, the Hopf fibration $S^3 \rightarrow S^2$ has Whitehead integral equal to one. On $S^3 \subset \mathbb{C}^2$ with $z_1 = \sin \chi e^{i\alpha}$, $z_2 = \cos \chi e^{i\beta}$ and connection $A = -i\langle z, dz \rangle = \sin^2 \chi d\alpha + \cos^2 \chi d\beta$, one has $A \wedge dA = 2 \sin \chi \cos \chi d\chi \wedge d\alpha \wedge d\beta$, so

$$\int_{S^3} A \wedge dA = 4\pi^2, \quad Q = \frac{1}{4\pi^2} \int_{S^3} A \wedge dA = 1 \quad (86)$$




Figure 33: **C7 phase correlator.** Checkerboard Metropolis XY, not equilibrated Villain. The ultraviolet exponent tracks the spin-wave line; $\eta = 1/4$ at $K = 2/\pi$ remains a thermodynamic target.




Figure 34: **C7 exponent $\eta(K)$.** Log fit of $g(r)$ versus spin-wave $\eta = 1/(2\pi K)$. Agreement at large K is the XY Goldstone; the jump $\eta = 1/4$ is not resolved on $L = 20$.

exactly. A Haar quadrature returns $Q = 0.9998$ (absolute error 2×10^{-4}). The domain volume $\int_{S^3} \text{vol} = 2\pi^2$ is recovered to the same order. String-free $\mathbb{R}^3$ Whitehead integrals (Coulomb-gauge padded FFT, and the pullback Hopf connection of the stereographic spinor) remain box-truncated and do not replace (86) or the lock $Q_H = 1/N$. An N -wound toroidal ansatz is a labeled probe, not a derivation of $1/N$.

11.10 Summary of L1 closures

The preceding calculations establish the following results on their stated truncations: the Galerkin Hessian yields a single discrete Goldstone and a non-democratic meV coupling $G^{(H)}$; H-CAT1 dark envelopes outlive bright ones under $\gamma \propto \omega^2$; the covariant Chern–Simons kinetic equation (82) implements a in the stepper with analytic flux-attachment residual 3.8×10^{-16} ; the QWZ Chern number and toy κ_{geom} track the physical branch (83); Track B matching gives $t_{\text{heal}}/U = 1$ and $t_\lambda/U = 0.379$; Villain dualization (85) reproduces ultraviolet η tracking $1/(2\pi K)$; and the S^3 Hopf integral (86) closes to machine precision. Several questions remain open: H-FN1 (five soft modes) is not realized on this truncation, H-TOR1 is parked and not accepted pending a 3D Weitzenböck quadrature, a transported-edge Hall simulation is not performed, MATBG quantum-metric and Bistritzer–MacDonald Hubbard coefficients are not computed, thermodynamic $\eta = 1/4$ is not resolved on $L = 20$, and the $\mathbb{R}^3$ hopfion quadrature does not replace the linking result $Q_H = 1/N$. Crucially, the locked fold, Option A* vacuum, bridge g_0^* , democratic $\text{Spec}(G)$, and $Q_H = 1/N$ are algebraic identities independent of these open questions.

12 Hopfions and Fractional Topological Charge

Hopfions are three-dimensional solitons of a unit-vector field $\hat{\mathbf{n}} : \mathbb{R}^3 \rightarrow S^2$. The topological charge is the Hopf invariant $H \in \mathbb{Z}$, which counts the linking number of the preimages of any two distinct points on S^2 [27, 28, 29, 30]. For a hopfion with n windings around the ring and m windings around the tube, $H = nm$. Independently, Chern–Simons/linking on $L(N, 1)$ with $H_2(L(N, 1); \mathbb{Z}) = 0$ locks

$$Q_H = \frac{1}{N}. \quad (87)$$

This is not a layer-fraction of a 3D Hopf integer, and it is not the Whitehead integral of the Hopf fibration $S^3 \rightarrow S^2$, which equals 1 (Section 11.9 computes $Q = 0.9998$ on S^3). For five-layer MATBG, $Q_H = 1/5$. The lock resolves the 2D/3D topology gap at the level of lens-space holonomy: a strictly 2D curvature integral on a fundamental 2-cycle does not exist.

Each hopfion nucleation event changes the Hall conductance by

$$\delta\sigma_{xy} = \frac{e^2}{2Nh} = \frac{Q_H e^2}{2h}, \quad (88)$$

producing an observable Hall staircase (Section 14).

12.1 Bogoliubov Quasiparticles and Inherited Topology

A fundamental question is how the topology of the underlying Chern bands propagates into the catalyzon spectrum. In a conventional BEC, Bogoliubov quasiparticles (“bogolons”) are obtained from the transformation $\gamma_{\mathbf{k}} = u_{\mathbf{k}}c_{\mathbf{k}} + v_{\mathbf{k}}c_{-\mathbf{k}}^\dagger$ diagonalizing the condensate Hamiltonian. When the




Figure 35: **Hall staircase schematic.** Step structure consistent with $Q_H = 1/N$ from Chern–Simons/linking on $L(N, 1)$, where $H_2(L(N, 1); \mathbb{Z}) = 0$.




Figure 36: **Hopfion topology and BEC vortex structure (3D visualization).** (Left) Hopfion toroidal topology with major radius $R = 50 \text{ nm}$ and minor radius $r = 8 \text{ nm}$, showing Hopf fibration field lines (black) and vortex core centerline (red). Hopf invariant $Q_H = 1/N$ for $N = 5$ layers. (Center) BEC vortex lattice in Abrikosov hexagonal pattern with Thomas-Fermi density profile and vortex core depletion. Red vertical lines mark vortex cores with healing length $\xi \approx 8 \text{ nm}$. (Right) Hopfion nucleation dynamics showing ring expansion from point nucleation at $t = 0$ to final radius $R = 50 \text{ nm}$ over 2 ps , illustrating parametric instability growth.




Figure 37: **Hopf charge numerical convergence.** Two-dimensional Chern–Simons flux of the mean-field connection $a_\theta = Q_H/\rho$ (equivalently the enclosed holonomy of the GPE phase $\arg \psi = Q_H \theta$) versus grid resolution. The integral converges to $Q_H = 1/5 = 0.2$. This is a 2D flux diagnostic, not a 3D Whitehead quadrature on $\mathbb{R}^3$. The lock $Q_H = 1/N$ is Chern–Simons/linking on $L(N, 1)$ (Section 26); the S^3 Hopf integral is Section 11.9.




Figure 38: **Hopf charge scaling $Q_H = 1/N$.** Two-dimensional CS-flux diagnostic versus layer number N for $N = 2$ – 8 , tracking the lock $Q_H = 1/N$ from Chern–Simons/linking on $L(N, 1)$. Error bars indicate grid-convergence uncertainty. This is not a 3D Hopf integer and not an S^3 Whitehead integral.

condensate forms in a band carrying Chern number $C \neq 0$, the Bogoliubov modes inherit Berry curvature through the coherence factors:

$$\Omega_{\text{Bog}}(\mathbf{k}) = (|u_{\mathbf{k}}|^2 - |v_{\mathbf{k}}|^2) \Omega_{\text{band}}(\mathbf{k}) + \Omega_{\text{pair}}(\mathbf{k}), \quad (89)$$

where Ω_{band} is the bare band Berry curvature and Ω_{pair} is an intrinsic contribution from the pairing structure in momentum space. The first term is weighted by the Bogoliubov “quasiparticle weight” ($|u|^2 - |v|^2$), which equals unity deep in the particle regime ($k \gg k_{\text{heal}}$) and vanishes at the condensate coherence scale.

12.1.1 Topological Constraint on Five-Mode Coupling

The inherited Berry curvature constrains the catalyzon coupling matrix G_{ij} (Section 8). Because the Bogoliubov amplitude mode (mode #1) carries curvature Ω_{Bog} , its overlap with other modes is weighted by the Chern number:

$$G_{1j} \sim C \cdot \int \frac{d^2k}{(2\pi)^2} \Omega_{\text{Bog}}(\mathbf{k}) \phi_j^*(\mathbf{k}) \langle \phi_{\text{ZPF}}(\mathbf{k}) \phi_{\text{ZPF}}(-\mathbf{k}) \rangle, \quad (90)$$

where $\phi_j(\mathbf{k})$ is the wavefunction of mode j in momentum space. This yields the scaling prediction:

$$|\lambda_{\text{min}}| \propto |C|^\alpha, \quad \alpha \approx 0.5-1, \quad (91)$$

implying that systems with trivial topology ($C = 0$) exhibit weaker catalyzon binding, while higher Chern numbers enhance the QVC effect. Experimentally, this can be tested by tuning the Chern number via gating or strain in MATBG and measuring the catalyzon gap $\Delta_{\text{cat}}(C)$.

12.1.2 Topological Protection of the Catalyzon

Each of the five hybridizing modes carries distinct topological quantum numbers: the Bogoliubov modes carry Berry phase from pairing, the Josephson mode carries interlayer winding number n_J , the plasmon inherits band Chern number C , the hopfion carries Hopf invariant H , and the acoustic mode is associated with the winding in $\pi_1(U(1)) = \mathbb{Z}$. The catalyzon superposition $|\text{cat}\rangle = \sum_i c_i |i\rangle$ must satisfy a *topological sum rule*:

$$N_{\text{total}} = \sum_{i=1}^5 |c_i|^2 n_i = \text{const.}, \quad (92)$$

where n_i is the topological index of mode i . This conservation law forbids catalyzon decay into channels that violate topological charge, producing selection rules that enhance the catalyzon lifetime beyond the naive estimate from energetics alone:

$$\tau_{\text{cat}} \sim \tau_0 \exp\left(\frac{\Delta_{\text{topo}}}{k_B T}\right), \quad (93)$$

where Δ_{topo} is the topological gap protecting against charge-violating decay channels. The topological protection makes the catalyzon a *robust* quasiparticle—observable in transport and spectroscopy even in the presence of disorder that would destroy conventional collective modes.

12.1.3 Josephson-Chern-Simons Duality

The Josephson plasma mode (mode #2) and the Chern-Simons gauge field in $\mathcal{F}_{\text{CS}}$ are related by a duality:

$$\partial_i \theta_j \longleftrightarrow a_i^{(j)}, \quad \oint \partial_i \theta \, dl = 2\pi \longleftrightarrow \int (\partial_i a_j - \partial_j a_i) \, dS = \frac{2\pi}{N}, \quad (94)$$

where θ_j is the condensate phase on layer j and $a_i^{(j)}$ is the emergent Chern-Simons gauge field. A 2π phase winding (vortex) maps to a fractional Chern-Simons flux quantum $\Phi_{\text{CS}} = 2\pi Q_H = 2\pi/N$ in the mean-field background $a_\theta = Q_H/\rho$ of Section 11.5. Textbook attachment at level k would be $2\pi/k = \pi/N$; that formula is not used to reset k or Q_H (Remark 10.1). The product $Q_H \times k = 2$ is the algebra of the two locks, not an independent pairing theorem.

13 Quantum TQGL Dynamics: Live Gap from ψ

13.1 Time-Dependent Simulation

We solve the quantum time-dependent TQGL equation using a split-step Fourier method. The equation of motion actually integrated is the *truncated* Gross–Pitaevskii equation of `qvc.tqgl.gpe`,

$$i\hbar \frac{\partial \psi}{\partial t} = \left[\kappa_{\text{GL}}(-\nabla^2) + \alpha_{\text{nl}}|\psi|^2 + \beta_{\text{nl}}|\psi|^4 + V_{\text{ZPF}}(\mathbf{r}, t) + V_{\text{phonon}}(\mathbf{r}, t) \right] \psi, \quad (95)$$

with $\kappa_{\text{GL}} = \Delta_0 a_H^2 = \hbar^2/(2m^*)$, $\alpha_{\text{nl}} = -\Delta_0$, $\beta_{\text{nl}} = \Delta_0/n_0$ at dimensionless $n_0 = 1$, matching the cubic-quintic potential $\alpha_{\text{nl}}n + \beta_{\text{nl}}n^2$ that multiplies ψ . There is no separate chemical-potential term in the stepper; $\mu_{\text{nl}}[\psi]$ is a diagnostic. Chern–Simons, Berry, and Skyrme terms in $\mathcal{F}$ (Section 9) remain in the parent functional and are not discretized in this stepper. The phonon drive is $V_{\text{phonon}} = g_{\text{ph}} \sin(\omega_0 t) \exp(-r_{\text{ring}}^2/(4r_{\text{core}}^2))$ at $\omega_0 = 2\Delta_0/\hbar = 1.82 \text{ ps}^{-1}$ (cyclic $\nu_{\text{GPE}} = 0.290 \text{ THz}$), and the Casimir coupling is $V_{\text{ZPF}} = -E_{\text{vac}} (A(t)/A_{\text{ref}}) \exp(-r_{\text{ring}}^2/(2r_{\text{core}}^2))$ with $E_{\text{vac}} = 0.1287 \text{ meV}$. A schematic temperature-dependent quartic $g(T)$ is not present in the integrator; V_{ZPF} already carries the vacuum amplitude.

Initial conditions consist of a planar ring texture: a toroidal density depletion of major radius $R = 50 \text{ nm}$ and minor radius $r_{\text{core}} = 8 \text{ nm}$, with amplitude $|\psi| = \sqrt{n_0}[1 - 0.1 \exp(-r_{\text{ring}}^2/(2r_{\text{core}}^2))]$ and planar phase winding $\arg(\psi) = Q_H \theta$ where $\theta = \arctan(y/x)$ and $Q_H = 1/5$. This is a two-dimensional fractional vortex, not a 3D Hopf map $\mathbb{R}^3 \rightarrow S^2$. A 3% random perturbation seeds dynamical evolution.

Numerical methods.—The computational framework uses: (i) split-step Fourier propagation (128^2 grid, $\Delta t = 2 \text{ fs}$, box $200 \times 200 \text{ nm}^2$) with coherent-window relative $\|\psi\|$ drift 1.30×10^{-14} per ps (covariant CS run: mass drift 8.3×10^{-6}); (ii) a Qi–Wu–Zhang two-band Chern insulator on a 48×48 Brillouin zone as a labeled geometric estimator $C = 0.993$, $\kappa_{\text{geom}}^{\text{QWZ}} = 0.0430 \text{ meV}$ (Section 11.6); Fukui lattice gauge is not in the L1 suite; (iii) checkerboard Metropolis XY on $L = 20$ (hundreds of sweeps; K capped at 3) for the C7 correlator $\eta_{\text{UV}} = 0.050$ versus spin-wave 0.053 (Section 11.8); (iv) RK4 for the DCE ODE and for the covariant kinetic half-step; (v) 14-mode Bessel K_0 sum for the Casimir energy with polynomial approximation (Abramowitz & Stegun 9.8.1–9.8.6, absolute error $< 2 \times 10^{-7}$). Table 8 records internal consistency on these stated solvers.

13.2 Parallel dynamical channels

The time-dependent theory comprises three parallel channels, each with its own observable and baseline.

1. **Static fold.** Participation λ through the bridge (29) selects a physical-branch gap. At $\lambda^* = 0.7628$ the static root is $\Delta^* = 0.2324 \text{ meV}$ (Section 5).
2. **Parametric resonance / catalyzon nucleation.** The phonon-mediated oscillation at $\omega_0 = 2\Delta_0/\hbar$ (GPE channel) or $2\Delta^*/\hbar$ (static working gap) drives the order parameter through the first Mathieu instability tongue. Core depletion and hopfion nucleation counts are the observables of this channel (Section 14). The 1 ps live gap *rises*; it does not drop to Δ^* .
3. **Coherent GPE.** The truncated integrator of this section evolves ψ and *reads* the gap from the field. It does not accept a prescribed $\Delta(t)$.

13.3 Live gap estimator

We simulate on a 128×128 spatial grid covering $200 \times 200 \text{ nm}^2$ for 1 ps (500 time steps of $\Delta t = 2 \text{ fs}$). The gap is extracted from the field,

$$\Delta_{\text{live}}[\psi] = \Delta_0 \sqrt{n_w[\psi]/n_{\text{ref}}}, \quad (96)$$

where n_w is the ring-weighted mean of $|\psi|^2$. On the locked drive the live gap starts at $\Delta_{\text{live}}(0) = 0.6000 \text{ meV}$ and ends at $\Delta_{\text{live}}(1 \text{ ps}) = 0.6305 \text{ meV}$ (a slight rise, from the attractive V_{ZPF} well). The minimum on the trajectory is 0.6000 meV , so the live gap remains above $\Delta_c = 0.109 \text{ meV}$. Phase coherence $g^{(1)} = |\langle \psi \rangle|^2 / \langle |\psi|^2 \rangle$ is 0.8832 at 1 ps. Relative $\|\psi\|$ drift is 1.30×10^{-14} . The estimator is an amplitude proxy and is blind to 2D phase-vortex unbinding (Section 6); Chern–Simons is truncated here as in the default vortex-RG scan.

The coupled DCE amplitude on this window reaches $A(1 \text{ ps})/A_c^{\text{fold}} = 0.9811$. Fold-slaved DCE over 80 ps (Section 20) saturates at $A^*/A_c^{\text{fold}} = 0.9117$ (resonant) and 0.8800 (detuned). Lindblad dissipation is not included in the present integrator.




Figure 39: **Condensate density $|\psi(\mathbf{r})|^2$ evolution.** (*Left*) Initial hopfion texture at $t = 0$ with toroidal vortex ring ($R = 50 \text{ nm}$, $r_{\text{core}} = 8 \text{ nm}$) on a 128×128 grid ($200 \times 200 \text{ nm}^2$). (*Right*) Evolved state at $t = 1 \text{ ps}$ showing ring-localized enhancement from the vacuum coupling V_{ZPF} . The growth is concentrated at the toroidal vortex ring where the Casimir mode sum is maximal, confirming spatial localization of catalyzation to the geometric cavity.




Figure 40: **Live gap $\Delta_{\text{live}}(\psi)$ vs static fold $\Delta(A(t))$.** Solid (teal): gap extracted from the ring-weighted density $n_w[\psi]$, rising from $\Delta_0 = 0.6000$ meV to 0.6305 meV over 1 ps of coherent GPE evolution. Dashed (gold): fold-slaved gap $\Delta(A(t))$ tracking the instantaneous DCE amplitude through the static gap map. Horizontal markers: $\Delta^* = 0.2324$ meV (working point) and $\Delta_c = 0.109$ meV (fold critical gap). The live gap remains above Δ_c throughout, confirming dynamical stability.




Figure 41: **Phase portrait in (A, Δ) space.** Teal curve: GPE trajectory $\Delta_{\text{live}}(A(t))$ over 1 ps. Gold dashed: DCE fold-resonant trajectory $\Delta(A)$ over 80 ps approaching the self-consistent fixed point. Square marker: locked operating point (λ^*, Δ^*) . The coral horizontal marks the fold critical gap $\Delta_c = 0.109$ meV; both trajectories remain above it. The system evolves within the physical branch, confirming the DCE self-limiting mechanism (Section 20).




Figure 42: **TQGL condensate density from the bridge-locked simulation.** Spatial profile $|\psi(\mathbf{r})|^2$ at $t = 1$ ps from the split-step Fourier integrator using bridge-locked parameters ($E_{\text{vac}} = 0.1287$ meV, $\lambda^* = 0.7628$). The ring-localized enhancement at $R = 50$ nm confirms that vacuum catalyzation concentrates at the hopfion toroidal core where the geometric cavity confines the Casimir mode sum.




Figure 43: **Lindblad decoherence scan at $t = 20$ ps.** Gap evolution $\Delta_{\text{live}}(t)$ under combined number-loss and dephasing. With frozen V_{ZPF} and $\tau_{\text{dec}} = 5$ ps the trajectory reaches 0.0872 meV, below $\Delta_c = 0.109$ meV, and is therefore unphysical on the static branch. Coupled dynamical Casimir evolution on the 1 ps window at the same rate remains above the fold ($\Delta_{\text{live}} = 0.5720$ meV, $g^{(1)} = 0.7102$).

14 Experimental Signatures

14.1 Hall Conductance Staircase

By Eq. (75), successive hopfion nucleation events produce quantized conductance steps of height $\delta\sigma_{xy} = e^2/(2Nh)$. For five-layer MATBG, $\delta\sigma_{xy} = 3.87 \times 10^{-6} \text{ S}$, directly measurable via Hall transport. The staircase pattern in σ_{xy} versus A_{ZPF} (tuned via pressure, twist angle, or chemical potential) provides a quantitative measurement of the fractional Hopf charge $Q_H = 1/N$.

14.2 Enhanced Schwinger Pair Production

The Schwinger rate for pair creation by a uniform field E in a medium with effective gap Δ is [31, 32, 33]

$$\Gamma_{\text{sch}}(E) \propto E^2 \exp\left(-\frac{\pi\Delta^2}{\hbar c E/e}\right). \quad (97)$$

In the TEVC the relevant gap in this kinematic regime is the catalyzon gap $\Delta_{\text{cat}} = 0.502 \text{ meV}$. The quantitative many-body rate used in this paper, including the thermal-quantum crossover $T^* = 118.7 \text{ mK}$, is derived in Section 19.

14.3 Laser-Driven QVC Excitation (Triple-Coincidence Test)

A circularly polarized THz drive at $\omega_L = 2\Delta_0/\hbar = 1.82 \text{ ps}^{-1}$ (cyclic $\nu_L = 0.290 \text{ THz}$; the working-gap variant is $\nu^* = 0.112 \text{ THz}$) is applied to a MATBG or TMD-bilayer TEVC at $T \ll T_{\text{BEC}}$. Three simultaneous signatures are predicted:

1. **L1 (Faraday-wave hopfion nucleation):** Stroboscopic imaging reveals sidebands at $\omega_L/2$ (Faraday waves) and discrete hopfion nucleation events scaling with driving amplitude squared (Mathieu instability) [34, 35].
2. **L2 (Hall-staircase response):** Hall conductance σ_{xy} climbs a staircase of quantized steps in lockstep with nucleation events, with step height $\delta\sigma_{xy} = e^2/(2Nh)$.
3. **L3 (Schwinger pair amplification):** THz/FIR emission or absorption shows a superlinear increase in exciton-pair production once the drive amplitude crosses the first Mathieu instability tongue. The pair threshold is set by $2\Delta/\hbar$, of order 0.1 THz .

All three signatures occur *only* when ω_L matches the two-particle threshold $2\Delta/\hbar$ of the channel being driven, providing a robust triple-coincidence test that cannot be explained by conventional (non-vacuum) coupling. The 1 ps GPE live gap of Section 13 rises from Δ_0 to 0.6305 meV and is a distinct parallel channel from the static fold Δ^* drawn in Fig. 44.




Figure 44: **Laser-driven QVC excitation.** (*Left*) Circularly polarized THz drive (σ^+) at $\omega_L = 2\Delta_0/\hbar = 1.82\text{ ps}^{-1}$ (cyclic $\nu_L = 0.290\text{ THz}$) incident on a TEVC sample at $T \ll T_{\text{BEC}}$, with simultaneous optical, transport, and spectroscopic readout. (*Center*) Parametric drive $A(t) = A_0 + \delta A \cos(\omega_L t)$ and subharmonic response $|\Psi|^2$ with Mathieu growth envelope. (*Right*) Schematic three-channel outputs: hopfion nucleation, Hall staircase of height $\delta\sigma_{xy} = e^2/(2Nh)$, and the static-fold gap $\Delta_{\text{ex}} \rightarrow \Delta^*$. The 1 ps GPE live gap rises to 0.6305 meV and belongs to a distinct dynamical sector.

15 Summary of Physical Predictions

The self-consistent framework, anchored by the toroidal Casimir geometry of the hopfion lattice, produces a falsifiable set of parameters for the TEVC platform. Conventions for lengths, Ginzburg–Landau coefficients, g_0 , Hall response, and retardation are collected in Section 10; provenance is classified in Appendix A.

The fat-torus geometry $(r, R) = (8, 50) \text{ nm}$, with aspect ratio $x = 2\pi r/R \approx 1.0053$, determines the vacuum energy through the Bessel sum (22), $F(x) = 0.58068$. Combined with $\hbar c_s = 0.07 \text{ meV} \cdot \mu\text{m}$,

$$E_{\text{vac}} = \frac{1}{4\pi^2} \frac{\hbar c_s}{r} F(x) = 0.1287 \text{ meV}. \quad (98)$$

The geometric amplitude is $A_{\text{geom}} R \approx 0.0919$.

This vacuum energy enters the dimensionless gap map $\delta = 1 + \alpha(\ln \delta + \ell)$ through $\alpha = \lambda E_{\text{vac}}/\Delta_0$, with $\ell = -2.7958$. The saddle-node fold occurs at $\alpha_c = \delta_c = 0.1818$, yielding $\Delta_c = 0.109 \text{ meV}$ and near-fold scaling $\delta - \delta_c \sim \pm \sqrt{\alpha_c - \alpha}$. The bridge (29) sets the maximum physical participation at $\lambda_{\text{fold}} = \alpha_c \Delta_0/E_{\text{vac}} = 0.8475$; the working point $\lambda^* = 0.90 \lambda_{\text{fold}} = 0.7628$ remains on the physical branch with static gap $\Delta^* = 0.2324 \text{ meV}$, a 61.3% reduction from the bare exchange gap.

The democratic coupling $G = g_0(J_N - I_N)$ has eigenvalues $\lambda_{\text{max}} = (N - 1)g_0$ (bright) and $\lambda_{\text{min}} = -g_0$ (catalyzon, multiplicity $N - 1$). With $g_0 = \lambda^* E_{\text{vac}} = 0.0982 \text{ meV}$, the catalyzon gap is $\Delta_{\text{cat}} = 0.502 \text{ meV}$ and the pair threshold is $E_{\text{pair}} = 0.1963 \text{ meV}$. Chern–Simons/linking on $L(N, 1)$ gives $Q_H = 1/N$, selected by $H_2(L(N, 1); \mathbb{Z}) = 0$.

Time-dependent evolution probes a distinct sector. The truncated GPE produces a live gap $\Delta_{\text{live}}(1 \text{ ps}) = 0.6305 \text{ meV}$ with $g^{(1)} = 0.8832$, while fold-slaved dynamical Casimir dynamics over 80 ps saturate at $A^*/A_c^{\text{fold}} = 0.9117$ (resonant) and 0.8800 (detuned). The Schwinger ratio $A_{\text{eff}}/A_c = 0.1636$ sets the thermal–quantum crossover at $T^* = 118.7 \text{ mK}$. Berry-phase Monte Carlo yields $R_{\text{mean}} = 0.908$.

In the vortex sector, the stiffness $\kappa(\alpha)$ along the physical branch remains above the Kosterlitz–Thouless unbinding surface at T^* on the primary closure. If a different material realization reached the KT surface first, the gap closing would be an essential singularity rather than square-root fold scaling—an experimentally distinguishable signature (Theorem 6.1).

16 Consolidated Parameter Table

Table 5 collects all numerical quantities entering the TEVC framework on the fat-torus freeze at $\theta = 1.1^\circ$. The full L1 dictionary with derivation provenance is given in Section 10; epistemic classification (locked inputs, derived results, computed outputs, matched conditions, ansätze, and conjectures) appears in Appendix A.

The static gap $\Delta^* = 0.2324 \text{ meV}$ at working participation λ^* is a 61.3% reduction from the bare exchange scale. Catalyzon hybridization through the dark eigenvalue $\lambda_{\text{min}} = -g_0$ is an independent reduction channel. Coherent GPE dynamics produce a live gap $\Delta_{\text{live}}(1 \text{ ps}) = 0.6305 \text{ meV}$ with $g^{(1)} = 0.8832$; that increase is a dynamical effect, not a static fold crossing. Strong number-loss Lindblad evolution at $\tau_{\text{dec}} = 5 \text{ ps}$ on a frozen potential can drive the live gap below Δ_c and is therefore unphysical on the static branch (Section 10.9). In the vortex sector, the fold is reached before Kosterlitz–Thouless unbinding at T^* on the primary stiffness closure. If a different material reached the KT surface first, an essential-singularity gap closing would replace the square-root fold (hypothesis H-VRG1).

Numerical evaluations of the gap map, Bessel sum, catalyzon well, Schwinger rate, and Hopf charge are collected in Appendix A.1. Source code is available in the accompanying repository.

Table 5: Numerical quantities for MATBG at $\theta = 1.1^\circ$ on the fat-torus freeze. The L1 dictionary (Section 10) provides full derivation context.

Symbol	Value	Meaning
J	0.30 meV	Exchange coupling
$\hbar c_s$	0.07 meV $\cdot\mu\text{m}$	Sound velocity $\times \hbar$
a_H	8 nm	Healing length (frozen cavity)
R	50 nm	Torus major radius
N	5	Number of layers
λ_M	~ 13 nm	Moiré period
$\Delta_{\text{ex}} = 2J$	0.600 meV	Bare exchange gap
$F(x)$	0.58068	Bessel sum $K_0(nx)$
E_{vac}	0.1287 meV	Toroidal vacuum energy
AR	0.0919	Geometric amplitude
$\alpha_c = \delta_c$	0.1818	Fold critical point
Δ_c	0.109 meV	Gap at the fold
λ_{fold}	0.8475	Max participation on physical branch
λ^*	0.7628	Working participation
Δ^*	0.2324 meV	Static gap at λ^* (61.3% reduction)
q^*	0.0330 nm^{-1}	Graphene-LA resonance at Δ^* ($q^*/K_M = 0.102$)
Q_H	$1/N = 1/5$	Fractional Hopf charge
g_0^*	0.0982 meV	$\lambda^* E_{\text{vac}}$ (catalyzon coupling)
Δ_{cat}	0.502 meV	$\Delta_{\text{ex}} - g_0^*$ (16.4% reduction)
κ_{GL}	$3.84 \times 10^{-5} \text{ meV} \cdot \mu\text{m}^2$	$\Delta_0 a_H^2$
m^*/m_e	0.99	From $\kappa_{\text{GL}} = \Delta_0 a_H^2$
ν_{GPE}	0.290 THz	Cyclic $2\Delta_0/h$ (GPE drive)
$\eta_{\text{ret}}(a_H)$	0.137	Retardation at $a_H = r$
$\eta_{\text{ret}}(\xi_{\text{ph}})$	2	Identity at $\xi_{\text{ph}} = \hbar c_s/\Delta_0$
R_{WS}	$2.18 \mu\text{m}$	Acoustic Hawking matching $T_H = T^*$
$\delta\sigma_{xy}$	$3.87 \times 10^{-6} \text{ S}$	$e^2/(2Nh)$ for $N = 5$
$\Delta_{\text{live}}(1 \text{ ps})$	0.6305 meV	Live gap from ψ
$g^{(1)}(1 \text{ ps})$	0.8832	Phase coherence
A^*/A_c^{fold}	0.9117 / 0.8800	DCE resonant / detuned saturation
T^*	118.7 mK	Schwinger crossover ($E_{\text{pair}} = 0.1963 \text{ meV}$)

17 Implications

17.1 Materials Science

If experimentally confirmed, QVC provides a materials-engineering pathway for reducing effective barriers in barrier-crossing phenomena. Flat-band engineering in moiré superlattices may enable designer quantum materials with tunable vacuum coupling, potentially impacting transport phenomena, topological phase transitions, and quantum information processing. The TEVC platform exemplifies a broader class of systems where geometry, topology, and vacuum fluctuations interplay at experimentally accessible scales.

17.2 Speculative Extensions

Highly speculative and requiring substantial future theoretical and experimental work: Understanding vacuum-mediated barrier reduction in condensed matter may inform future studies of barrier-crossing phenomena in other contexts, including nuclear fusion reactions. If analogous vacuum-catalyzation mechanisms exist for Coulomb barriers in fusion plasmas, rates could be enhanced without requiring net energy input from zero-point fields. However, such extensions face formidable challenges: (1) translating the TEVC mechanism to hot, dense plasmas; (2) identifying the relevant zero-point modes in a fusion context; (3) quantifying enhancement factors under realistic conditions. We emphasize that *no energy extraction from the vacuum is claimed*, and any fusion relevance remains conjectural pending rigorous plasma-physics analysis.

18 Connections to Cavity QED and Vacuumronics

Recent experimental and theoretical developments in cavity quantum electrodynamics establish direct connections to the QVC framework. Enkner et al. (2025)[36] demonstrated tunable vacuum-field control of fractional and integer quantum Hall phases using a split-ring resonator, while Jiang (2025)[37] proposed “Vacuumronics” as a paradigm for engineering material properties through cavity-enhanced vacuum fluctuations. These works validate the central premise of QVC: zero-point fluctuations can actively reshape energy landscapes without supplying net energy. In this section, we formalize the connection and derive several extensions.

18.1 Self-Cavitizing TEVC: Moiré Geometry as Intrinsic Cavity

In Enkner et al.’s experiment, an *external* resonator provides tunable vacuum coupling. A key insight of QVC is that the TEVC itself provides an *intrinsic* cavity: the moiré superlattice confines electromagnetic modes laterally at wavelength $\lambda_M \approx 13$ nm, while each graphene layer acts as a partially reflecting mirror due to its universal optical conductivity $\sigma = \pi\alpha e^2/h$ (where $\alpha = 1/137$ is the fine-structure constant).

For an N -layer stack, the cumulative vacuum field enhancement factor is:

$$\mathcal{F}_{\text{moiré}}(N) = \sqrt{1 + (N - 2) r_{\text{eff}}^2 Q_{\text{moiré}}}, \quad (99)$$

where $r_{\text{eff}}^2 = (\pi\alpha)^2 \cdot (U/W)$ is the effective reflectivity per layer enhanced by the flat-band ratio $U/W \approx 10$ (interaction energy over bandwidth), and $Q_{\text{moiré}} = 10 + 2(N - 2)$ is the moiré cavity quality factor. Physically, each additional layer adds one mirror to the cavity, improving mode confinement.

The static gap equation of Section 3 does not include $\mathcal{F}_{\text{moiré}}$: $\alpha = \lambda E_{\text{vac}}/\Delta_0$ is evaluated on $A_{\text{geom}} = E_{\text{vac}}/(\hbar c_s) = 1.839 \mu\text{m}^{-1}$. When the stack is treated as an intrinsic cavity one instead has

$$A_{\text{eff}} = \mathcal{F}_{\text{moiré}}(N) A_{\text{geom}}, \quad \alpha_{\text{cav}} = \lambda \mathcal{F}_{\text{moiré}}(N) \frac{E_{\text{vac}}}{\Delta_0}, \quad (100)$$

and Eq. (12) with A_{geom} replaced by A_{eff} ,

$$\Delta = 2J + \hbar c_s \cdot \mathcal{F}_{\text{moiré}}(N) \cdot A_{\text{geom}} \left[\ln \left(\frac{\Delta a_H}{2\hbar c_s} \right) + \gamma_E \right]. \quad (101)$$

The moiré stack concentrates vacuum fluctuations into the condensate volume. This enhancement is additional to the frozen-cavity evaluation of E_{vac} and λ^* .

Equation (100) grows from 1 (bilayer) through 1.119 at $N = 5$ to 1.52 at $N = 11$. At the operating stack, $A_{\text{eff}} = 2.058 \mu\text{m}^{-1}$. Fold participation on the enhanced map is $\lambda_{\text{fold}}^{(F)} = \lambda_{\text{fold}}/\mathcal{F}_{\text{moiré}} = 0.7573$. The unenhanced working point λ^* times $\mathcal{F}_{\text{moiré}}$ lies beyond the fold, so the cavity working point is retuned to $\lambda_F^* = 0.6816$ with static root $\Delta_F^* = 0.2914 \text{ meV}$.

The cavity-enhanced gap therefore depends on layer number N , with stronger catalyzation for thicker stacks provided participation is held at or below $\lambda_{\text{fold}}^{(F)}$. This is testable by fabricating TEVCs with $N = 3, 5, 7, 9$ layers and measuring the excitonic gap via tunneling spectroscopy at fixed twist angle.

18.2 Catalyzon-Polariton: Coupling to External Cavities

When a TEVC is placed inside an external cavity, the catalyzon hybridizes with the cavity photon. Extending the resonant coupling matrix of Section 8 by a resonator mode ω_c gives

$$\hat{M}_6 = \begin{pmatrix} \hat{M}_5 & \mathbf{g}_c \\ \mathbf{g}_c^\dagger & \omega_c \end{pmatrix}, \quad (102)$$

where $\mathbf{g}_c$ couples the cavity primarily to the flat-band plasmon, which carries electric dipole character, with matrix elements of order g_0^* .

Second-order perturbation in the cavity-plasmon coupling g_c then gives

$$\frac{\Delta_{\text{cat-pol}}}{\Delta_{\text{ex}}} \approx \frac{\Delta_{\text{cat}}}{\Delta_{\text{ex}}} \times \left(1 - \frac{2g_c^2}{\omega_c \Delta_{\text{cat}}} \right), \quad (103)$$

so that an additional gap reduction appears in the ultrastrong regime $g_c/\omega_c \gtrsim 0.1$.

Placing a MATBG device in a tunable split-ring resonator (Enkner geometry) and sweeping sample-resonator distance would vary g_c . Tunneling spectroscopy of the excitonic gap versus coupling is then predicted to show a nonlinear reduction distinct from ordinary linear Rabi splitting.

18.3 Devil's Staircase: Fractal Structure of the Hall Response

Following Rotondo et al.[38], we map the hopfion vortex lattice to a one-dimensional lattice gas in the thin-torus limit. Define occupation $n_i \in \{0, 1\}$ at moiré site i (hopfion present or absent). The effective Hamiltonian:

$$H_{\text{hop}} = \sum_{i < j} V(|i - j|) n_i n_j - \mu_{\text{eff}} \sum_i n_i, \quad (104)$$

where $V(r) \sim V_0/r^\alpha$ is the inter-hopfion repulsion (dipolar: $\alpha = 3$) and the effective chemical potential $\mu_{\text{eff}} = E_{\text{Cas}} + A_{\text{ZPF}} \cdot \partial\Delta/\partial n_{\text{hop}}$ is vacuum-tunable.

This mapping reveals that the Hall response has *devil's staircase* structure: every rational hopfion filling $\nu_H = p/q$ produces a quantized plateau in σ_{xy} with width:

$$w_{p/q} = 2|V_q| \sin\left(\frac{\pi p}{q}\right) \cdot \left(1 + \frac{A_{\text{ZPF}} q^2}{A_c}\right)^{-1}. \quad (105)$$

The Hall conductance at filling $\nu_H = p/q$ is:

$$\sigma_{xy} = \frac{e^2}{h} \cdot \frac{p}{2qN}. \quad (106)$$

The Hausdorff dimension of the staircase complement (the set of transitions between plateaux) is:

$$d_H = 1 - \frac{1}{\alpha + 1} = \frac{3}{4} \quad (\text{for dipolar hopfions, } \alpha = 3). \quad (107)$$

Physical interpretation.—The QVC factor in Eq. (106) narrows higher-order plateaux faster than lower-order ones (because of the q^2 weighting). This means vacuum catalyzation makes the staircase *sharper*—more Cantor-like—with the finest plateaux disappearing first. In the limit $A_{\text{ZPF}} \rightarrow A_c$, only the $q = 1$ plateau survives, corresponding to a completely ordered hopfion lattice.

Prediction.—With fine magnetic field resolution ($\delta B/B \sim 10^{-4}$), the Hall staircase should exhibit sub-steps between the main N -integer steps predicted in Section 14. The fractal dimension of the measured staircase discriminates between hopfion interaction models ($\alpha = 3$ dipolar vs. exponential compact).

18.4 Quantum Atmosphere and Vacuum Information Transfer

Jiang (2025)[37] showed that a Chern insulator radiates a “quantum atmosphere” into surrounding vacuum—the broken time-reversal symmetry modifies the vacuum spectral function with power-law decay:

$$\Delta E_{\text{QF}}(r) = C^2 \left(\frac{a_0}{r}\right)^2 E_0, \quad a_0 \approx 10 \text{ nm}, \quad E_0 \approx 0.06 \text{ } \mu\text{eV}, \quad (108)$$

where C is the Chern number. For a TEVC with $C = 1$ per valley and moiré period $\lambda_M \approx 13$ nm, the quantum atmosphere extends over precisely the length scale of interlayer vacuum coupling.

This provides a new physical interpretation of our self-consistency loop $\Delta \leftrightarrow A_{\text{ZPF}}$ (Section 4): each TEVC layer projects its topological order into the interlayer vacuum via the quantum atmosphere, and these overlapping atmospheres feed back into the gap equation. The vacuum is not merely a passive cavity—it carries topological information.

Information-theoretic consequence.—The mutual information between condensate and vacuum is bounded below by:

$$I(\text{cond} : \text{vac}) \geq \frac{1}{2} \ln |C|, \quad (109)$$

which vanishes for $C = 1$ but grows logarithmically for higher Chern numbers accessible in multilayer stacks. This predicts a T_c enhancement:

$$\frac{\delta T_c}{T_c} = \frac{I(\text{cond} : \text{vac})}{S_{\text{config}}} \approx \frac{\ln |C|}{2 \ln 4} \approx 25\% \quad (C = 2), \quad (110)$$

where $S_{\text{config}} = \ln 4$ accounts for the four-fold spin $\times$ valley degeneracy. The physical meaning: destroying topological order also destroys the vacuum’s information content, which provides an entropic stabilization not captured by standard Ginzburg-Landau theory.

18.5 Vacuum-Catalyzed Topological Phase Transitions

QVC can catalyze transitions between topological phases ($C \rightarrow C'$) by reducing the gap-closing barrier. The transition rate exhibits a striking double-exponential form:

$$\Gamma_{C \rightarrow C'} = \omega_0 \exp\left(-\frac{E_{\text{barrier}}^{(0)}}{k_B T} e^{-A_{\text{ZPF}}/A_c}\right), \quad (111)$$

where $E_{\text{barrier}}^{(0)}$ is the bare gap-closing energy. At the MATBG operating point ($A_{\text{ZPF}}/A_c \approx 0.4$), the barrier is reduced by 33%, and the critical magnetic field for the transition decreases:

$$B_c^{\text{QVC}} = B_c^{(0)} \sqrt{1 - \frac{\Delta_{\text{cat}}}{\Delta_{\text{ex}}}} \approx 0.72 B_c^{(0)}. \quad (112)$$

This 28% reduction in critical field is experimentally detectable and provides a clean test: measure the $C = 1 \rightarrow C = 2$ transition field in a TEVC and compare to the bare value from theory. The double-exponential sensitivity near A_c means even small variations in vacuum coupling produce dramatic changes in transition rates.

Cavity-enhanced QVC predictions (corrected spectrum / fold)




Figure 45: Cavity-enhanced QVC predictions (regenerated). (a) Moiré self-cavity enhancement and A_{ZPF}/A_c vs. N . (b) Hall staircase schematic with $Q_H = 1/N$. (c) Catalyzon-cavity spectrum: dark multiplet at $\lambda_{\min} = -g_0$ (multiplicity 4) plus bright-cavity polaritons. (d) Barrier reduction and rate vs. A_{ZPF}/A_c from the saddle-node fold map, with MATBG operating point marked.




Figure 46: (a) Three-dimensional visualization of the devil's staircase Hall conductance as a function of both effective chemical potential and vacuum coupling strength A_{ZPF} . Higher vacuum coupling narrows the fractional plateaux, sharpening the fractal structure. (b) Quantum atmosphere energy shift ΔE_{QF} as a function of distance r from the TEVC surface for different Chern numbers $C = 1, 2, 3, 5$. Exponential screening beyond $\lambda_M \approx 13$ nm reflects moiré-scale vacuum confinement.

19 Schwinger Twin Channel and Thermal-Quantum Crossover

The many-body analogue of Schwinger pair production provides a second pathway for vacuum-catalyzed barrier crossing, operating in a distinct kinematic regime from the catalyzon mechanism (Section 8). While the catalyzon reduces the effective barrier height through resonant hybridization of collective modes, the Schwinger channel exploits the exponential sensitivity of quantum tunneling to the ratio of zero-point field amplitude to critical amplitude.

19.1 Mapping from Relativistic QED

In relativistic QED, the Schwinger rate for electron-positron pair creation by a uniform electric field E is [32, 39]

$$\Gamma_{\text{Schwinger}}^{\text{QED}} = \frac{(eE)^2}{4\pi^3\hbar^2c} \exp\left(-\frac{\pi m^2c^3}{\hbar eE}\right) = \frac{(eE)^2}{4\pi^3\hbar^2c} \exp\left(-\frac{\pi E_c}{E}\right), \quad (113)$$

where $E_c = m^2c^3/(\hbar e)$ is the critical Schwinger field. The key structural feature is the exponential dependence on the ratio E_c/E : the rate is exponentially suppressed when $E \ll E_c$ and becomes appreciable only as $E \rightarrow E_c$.

In the TEVC condensate, the analogous mapping replaces the QED electric field with the *zero-point phonon displacement amplitude* [40]. The physical reasoning is as follows: excitonic quasiparticles in the flat band experience a local zero-point displacement field A_{ZPF} from the toroidal Casimir modes propagating at the zone-folded optical phonon group velocity c_s . The critical amplitude A_c is defined by the condition that the vacuum-phonon work done over one coherence time $\tau_c = \hbar/\Delta_{\text{ex}}$

equals the excitonic gap:

$$\hbar c_s \cdot A_c = \Delta_{\text{ex}} \implies A_c = \frac{\Delta_{\text{ex}}}{\hbar c_s}. \quad (114)$$

This is the phononic Schwinger threshold: the zero-point amplitude at which vacuum fluctuations can supply enough energy to produce an exciton pair within one coherence cycle. Numerically, $A_c = 0.600 \text{ meV} / 0.07 \text{ meV} \cdot \mu\text{m} = 8.571 \mu\text{m}^{-1}$. In 2+1 dimensions the tunneling rate takes the linear-prefactor form used as the lock,

$$\Gamma_{\text{Sch}} = \omega_0 \left(\frac{A_{\text{eff}}}{A_c} \right) \exp \left(-\pi \frac{A_c}{A_{\text{eff}}} \right), \quad (115)$$

where $\omega_0 = 2\Delta^*/\hbar$ is the pair frequency. The ratio is identified with the frozen-cavity drive, not with a cavity-enhanced amplitude:

$$\frac{A_{\text{eff}}}{A_c} = \lambda^* \cdot \frac{E_{\text{vac}}}{\Delta_{\text{ex}}} \equiv \alpha^* = 0.1636. \quad (116)$$

Moiré $\mathcal{F}_{\text{moiré}}$ of Section 18.1 is a labeled cavity enhancement of A_{geom} and is *not* inserted into this exponent.

19.2 Many-Body Schwinger Rate

The locked many-body Schwinger rate is therefore

$$\Gamma_{\text{Schwinger}} = \omega_0 \left(\frac{A_{\text{eff}}}{A_c} \right) \exp \left(-\frac{\pi A_c}{A_{\text{eff}}} \right) = \omega_0 \alpha^* \exp(-\pi/\alpha^*), \quad (117)$$

with $\omega_0 = 2\Delta^*/\hbar \approx 7.06 \times 10^{11} \text{ rad/s}$. Substituting $\alpha^* = 0.1636$ yields

$$\text{Schwinger exponent: } \frac{\pi}{\alpha^*} = 19.20, \quad (118)$$

$$\text{Prefactor: } \omega_0 \alpha^* \approx 1.15 \times 10^{11} \text{ s}^{-1}, \quad (119)$$

$$\Gamma_{\text{Schwinger}} = 5.30 \times 10^2 \text{ Hz}. \quad (120)$$

Per moiré unit cell this is a slow process; in a typical sample of $\sim 10^6$ unit cells the ensemble rate is of order $5 \times 10^8 \text{ Hz}$, a photoluminescence signal in reach of a dilution-refrigerator measurement.

19.3 Thermal-Schwinger Crossover Temperature T^*

At finite temperature, thermal activation competes with quantum Schwinger tunneling. The thermal pair creation rate for excitonic pairs is:

$$\Gamma_{\text{thermal}} = \omega_0 \exp \left(-\frac{E_{\text{pair}}}{k_B T} \right), \quad (121)$$

where $E_{\text{pair}} = 2|\lambda_{\text{min}}| = 0.1963 \text{ meV}$ is the pair-creation threshold—the energy cost to excite a virtual excitonic pair at the catalyzon resonance. This is distinct from the full catalyzon gap Δ_{cat} : the Schwinger process creates pairs from the dressed vacuum, and the thermal analogue involves activating over the same vacuum-renormalized barrier.

The crossover temperature T^* is defined by equating the two rates:

$$\frac{E_{\text{pair}}}{k_B T^*} = \frac{\pi A_c}{A_{\text{eff}}} = \frac{\pi}{\alpha^*} = 19.20. \quad (122)$$




Figure 47: **Many-body Schwinger rate in the TEVC.** Γ_{Sch} (ps^{-1} , log scale) versus $A_{\text{eff}}/A_{c,\text{Sch}}$, computed with the corrected $E_{\text{vac}} = 0.1287 \text{ meV}$ and democratic $|\lambda_{\text{min}}| = g_0$. Vertical dashed: operating point $A^*/A_{c,\text{Sch}}$ at the self-consistent vacuum amplitude. The exponential suppression $\exp(-\pi A_c/A_{\text{eff}})$ sets the rate hierarchy; at the MATBG parameters, $\Gamma_{\text{Sch}} \sim 5 \times 10^2 \text{ Hz}$ per unit cell.

Solving for T^* :

$$T^* = \frac{E_{\text{pair}}}{k_B} \cdot \frac{\alpha^*}{\pi} = 118.7 \text{ mK}. \quad (123)$$

The physical meaning of T^* is that below 118.7 mK, the dominant mechanism for excitonic pair creation in the TEVC switches from thermal activation over the barrier to quantum Schwinger tunneling *through* the barrier, catalyzed by the zero-point field. This crossover temperature is experimentally accessible in dilution refrigerators and provides a sharp prediction:

Experimental signature.—Plot the pair-creation luminescence rate versus inverse temperature $1/T$. Above T^* , the Arrhenius slope gives Δ_{cat}/k_B . Below T^* , the rate *saturates* to the temperature-independent Schwinger value. The kink at $T^* = 118.7 \text{ mK}$ is a signature of the twin-channel mechanism.

19.4 Twin-Channel Enhancement

The total pair-creation rate combining both channels is:

$$\Gamma_{\text{total}}(T) = \Gamma_{\text{Schwinger}} + \Gamma_{\text{thermal}}(T) = \omega_0 \left[e^{-\pi A_c/A_{\text{eff}}} + e^{-E_{\text{pair}}/k_B T} \right]. \quad (124)$$

At the crossover $T = T^*$, both terms contribute equally and the total rate is $2\Gamma_{\text{Schwinger}}$ —a factor-of-2 enhancement over either channel alone. The smooth crossover (rather than a sharp transition) reflects the additive nature of independent tunneling paths through the barrier landscape.

20 Dynamical Casimir Self-Limiting Theorem

The dynamical Casimir effect (DCE) in the TEVC context describes the parametric amplification of zero-point fluctuations when the effective cavity boundary (the condensate coherence length ξ) oscillates at twice the resonant frequency. While linear DCE theory predicts unbounded exponential growth $A(t) \propto \cosh(\Lambda t)$, the self-consistent coupling to the gap equation introduces a feedback mechanism that naturally saturates the growth—producing a stable operating point for the “vacuumronic transistor” concept.

20.1 Linear DCE Growth

Consider a time-dependent modulation of the excitonic gap:

$$\Delta(t) = \Delta_0 + \delta\Delta \cos(\omega_{\text{pump}} t), \quad (125)$$

where $\omega_{\text{pump}} = 2\omega_0 = 4\Delta_0/\hbar$ is twice the catalyzon frequency (parametric resonance condition). The coherence length $\xi \propto 1/\Delta$ oscillates accordingly, modulating the effective cavity size. By Uhlmann et al.[41], photon creation in a cavity with time-dependent dielectric $\epsilon(t) = \epsilon_0 + \chi \cos(\omega_{\text{pump}} t)$ produces exponential growth:

$$\langle N(t) \rangle = \sinh^2 \left(\frac{k_{\parallel}^2}{\omega} \cdot \frac{\chi}{\epsilon^{(2)}} \cdot \frac{a}{L} \cdot t \right), \quad (126)$$

where $k_{\parallel}$ is the transverse wavevector, χ the modulation depth, $\epsilon^{(2)}$ the second-order susceptibility, and a/L the geometric factor.

In the TEVC mapping, the effective zero-point amplitude grows as:

$$A_{\text{ZPF}}(t) = A_0 \cosh(\Lambda t), \quad \Lambda = \frac{\delta\Delta}{2\hbar} \cdot \frac{Q_{\text{moiré}}}{2\pi}, \quad (127)$$

where $Q_{\text{moiré}} \sim 10\text{--}50$ is the quality factor of the moiré cavity and Λ is the parametric growth rate. For typical modulation $\delta\Delta/\Delta_0 \sim 0.1$ and $Q_{\text{moiré}} = 20$:

$$\Lambda \approx \frac{0.1 \times 0.2324 \text{ meV}}{2\hbar} \cdot \frac{20}{2\pi} \approx 5.6 \times 10^{10} \text{ s}^{-1} \quad (128)$$

using Δ^* . This predicts e -folding on a timescale $\tau_{\text{DCE}} = 1/\Lambda \approx 18 \text{ ps}$.

20.2 Self-Consistent Feedback: The Detuning Mechanism

The linear growth (Eq. 128) cannot persist indefinitely because increasing A_{ZPF} feeds back into the gap equation (Eq. 12), reducing Δ . As Δ drops, the catalyzon frequency $\omega_{\text{cat}} = 2\Delta/\hbar$ shifts away from the parametric resonance condition $\omega_{\text{pump}}/2$, suppressing further growth. The self-consistent dynamics are governed by the coupled ordinary differential equations:

$$\frac{dA}{dt} = \Lambda(\Delta(A)) \cdot A \cdot \mathcal{L}\left(\frac{\delta\omega}{\Gamma_{\text{cav}}}\right), \quad (129)$$

$$\Delta(A) = 2J + \hbar c_s A \left[\ln\left(\frac{\Delta(A) \cdot a_H}{2\hbar c_s}\right) + \gamma_E \right], \quad (130)$$

$$\delta\omega = \frac{\omega_{\text{pump}}}{2} - \frac{2\Delta(A)}{\hbar}, \quad (131)$$

where $\mathcal{L}(x) = 1/(1+x^2)$ is the Lorentzian detuning function and $\Gamma_{\text{cav}} = \omega_0/Q_{\text{moiré}}$ is the cavity linewidth. The detuning $\delta\omega$ (Eq. 132) measures how far the instantaneous catalyzon frequency has drifted from the pump's half-frequency.

The physical content of this self-limiting mechanism deserves emphasis. The Lorentzian response $\mathcal{L}(\delta\omega/\Gamma_{\text{cav}}) = 1/(1+(\delta\omega/\Gamma_{\text{cav}})^2)$ depends implicitly on A through the detuning $\delta\omega(A)$: as A grows, it modifies Δ via the gap map (Eq. 131), which shifts the parametric resonance frequency $\omega_0 = 2\Delta(A)/\hbar$ progressively away from the drive frequency $\omega_{\text{pump}}/2$. This continuous detuning reduces the effective parametric gain at every step, constituting an intrinsic negative feedback loop that requires no external regulation. Crucially, the fixed point A^* occurs where the detuned gain exactly balances the threshold condition for net growth—it is the system's own back-reaction on its resonance condition, not an externally imposed dissipation channel, that enforces saturation. For comparison with experimental DCE implementations[42, 43], where modulation supplies energy and material dissipation determines the steady state, one should augment the idealized dynamics with phenomenological cavity and phonon losses κ_{loss} and a nonlinear saturation term ηA^3 :

$$\frac{dA}{dt} = \left[\Lambda(\Delta(A)) \mathcal{L}\left(\frac{\delta\omega(A)}{\Gamma_{\text{cav}}}\right) - \kappa_{\text{loss}} \right] A - \eta A^3. \quad (132)$$

The ηA^3 term provides additional saturation above threshold via nonlinear mode-mode scattering, while κ_{loss} sets the minimum gain required to sustain amplification; together they ensure that the operating point A^* is robust against parameter variations in realistic material environments.

20.3 Fixed-Point Analysis and Saturation

Theorem 20.1 (DCE Self-Limiting Theorem). *The augmented dynamical system (Eq. 133) with detuning feedback (Eqs. 131–132) possesses a unique stable fixed point A^* satisfying $dA/dt = 0$, given implicitly by:*

$$\delta\omega(A^*) = \pm\Gamma_{\text{cav}} \cdot \sqrt{\frac{\Lambda(\Delta(A^*))}{\Lambda_0} - 1}, \quad (133)$$

where $\Lambda_0 \equiv \kappa_{\text{loss}}$ is the total dissipation rate (the threshold growth rate below which parametric amplification ceases). The system exponentially approaches A^* with relaxation rate $\gamma_{\text{relax}} = \partial(dA/dt)/\partial A|_{A^*} < 0$.

Proof. At the fixed point, $dA/dt = 0$ requires either $A^* = 0$ (trivial) or $\mathcal{L}(\delta\omega/\Gamma_{\text{cav}}) = 0$, which has no solution. Instead, the non-trivial fixed point requires the detuned gain to balance dissipation, $\Lambda(\Delta(A^*)) \cdot \mathcal{L}(\delta\omega/\Gamma_{\text{cav}}) = \Lambda_0$. Equating growth to loss in the augmented system (Eq. 133) at leading order in A (neglecting the ηA^3 correction):

$$\frac{\Lambda(\Delta(A^*))}{1 + (\delta\omega/\Gamma_{\text{cav}})^2} = \Lambda_0 \equiv \kappa_{\text{loss}}. \quad (134)$$

This has a unique solution because $\Lambda(\Delta)$ decreases monotonically with A (since Δ decreases, reducing the parametric coupling), while the Lorentzian suppression increases with detuning. The intersection defines A^* .

Stability follows from the negative slope: for $A > A^*$, the detuning increases, the Lorentzian suppresses growth below the loss rate, and A decreases back to A^* . Conversely for $A < A^*$. $\square$

Physical consequence.—The DCE in a TEVC is *self-limiting*: the zero-point amplitude cannot grow beyond A^* because detuning acts as a negative feedback. This makes the vacuumtronic transistor a stable device with a well-defined operating point, rather than a runaway instability.

20.4 Numerical Solution

Solving the augmented DCE dynamics (Eq. 133 with gap feedback Eqs. 131–132) numerically with MATBG parameters ($Q_{\text{moiré}} = 20$, $\delta\Delta/\Delta_0 = 0.1$, $\kappa_{\text{loss}} = 10^9 \text{ s}^{-1}$) on the frozen cavity, the fold-slaved evolution reaches

$$A^*/A_c^{\text{fold}} = 0.9117 \quad (\text{resonant}), \quad \Delta_{\text{ss}} = 0.2240 \text{ meV}, \quad (135)$$

$$A^*/A_c^{\text{fold}} = 0.8800 \quad (\text{detuned}), \quad \Delta_{\text{ss}} = 0.2461 \text{ meV}. \quad (136)$$

The Lorentzian is not parked at zero in the detuned run ($L(0) = 0.20$). The approach to A^* follows exponential relaxation with characteristic time $\tau_{\text{relax}} \approx 25 \text{ ps}$ after the initial cosh-growth phase ($\sim 10 \text{ ps}$). The total transient from pump-on to steady state is $\sim 35 \text{ ps}$.

Operating regimes.—Three distinct regimes emerge:

1. **Sub-threshold** ($\delta\Delta/\Delta_0 < \delta\Delta_{\text{thr}}$): Parametric gain insufficient to overcome loss; A remains at thermal equilibrium value.
2. **Transient growth** ($t < \tau_{\text{relax}}$): Exponential cosh-growth as in linear DCE (Eq. 128).
3. **Saturated steady state** ($t \gg \tau_{\text{relax}}$): Self-consistent fixed point A^* maintained indefinitely as long as pump is active.




Figure 48: **DCE amplitude dynamics** $A(t)$. Teal: fold-slaved resonant drive. Coral: detuning-feedback drive (Lorentzian not parked at zero). Horizontal markers: A_c^{fold} (gold dashed) and $A(\lambda^*)$ (blue dotted). Both drives saturate: $A^*/A_c^{\text{fold}} = 0.9117$ (resonant) and 0.8800 (detuned), confirming the DCE self-limiting theorem. The Lorentzian detuning provides the negative-feedback mechanism that prevents runaway parametric amplification.

21 Berry Phase Enhancement of the Catalyzon

The catalyzon formation theorem (Section 8, Theorem 6.1) was originally proven for real symmetric coupling matrices G . In materials with non-trivial band topology—precisely the class of systems where QVC operates—the Berry curvature $\Omega(\mathbf{k})$ introduces complex phases into the mode coupling, promoting G from real symmetric to complex Hermitian. The Berry phase modifies the distribution of $|\lambda_{\min}|$; a GUE-like random-matrix ensemble yields a mean ratio ≈ 1.43 relative to GOE, while a magnitude-preserving Hermitian draw on the locked g_0 yields $R_{\text{mean}} = 0.908$.

21.1 Complex Hermitian Extension

When the band carries Chern number $C \neq 0$, the vacuum-mediated coupling between collective modes acquires a geometric phase from the parallel transport of Bloch states around the Fermi surface. The coupling matrix becomes:

$$G_{ij} \rightarrow G_{ij}^{(\text{Berry})} = |G_{ij}| \exp \left(i \oint_{\text{FS}} \mathcal{A}_i^*(\mathbf{k}) \cdot \nabla_{\mathbf{k}} \mathcal{A}_j(\mathbf{k}) d\mathbf{k} \right), \quad (137)$$

where $\mathcal{A}_i(\mathbf{k}) = \langle u_i(\mathbf{k}) | \nabla_{\mathbf{k}} | u_i(\mathbf{k}) \rangle$ is the Berry connection for mode i and the integral runs over the Fermi surface. The resulting $G^{(\text{Berry})}$ is Hermitian (not symmetric): $G_{ij}^* = G_{ji}$ but generically $G_{ij} \neq G_{ji}$.

Theorem 21.1 (Complex Hermitian Catalyzon Formation). *Let $G \in \mathbb{C}^{5 \times 5}$ be Hermitian with $G_{ii} = 0$ for all i and $G \neq 0$. Then $\lambda_{\min}(G) < 0$, i.e., the catalyzon exists.*

Proof. The proof proceeds identically to Theorem 6.1:

1. Hermiticity guarantees all eigenvalues $\{\lambda_i\}$ are real (spectral theorem).
2. $G_{ii} = 0$ implies $\text{Tr}(G) = \sum_i \lambda_i = 0$.
3. $G \neq 0$ implies $\|G\|_F^2 = \sum_{ij} |G_{ij}|^2 = \sum_i \lambda_i^2 > 0$, so not all λ_i vanish.
4. From (2) and (3): eigenvalues of both signs exist, hence $\lambda_{\min} < 0$.

□

The extension is mathematically trivial because the key properties—zero trace and nonzero Frobenius norm—are shared by real symmetric and complex Hermitian matrices alike. However, the *magnitude* of $\lambda_{\min}$ differs substantially between the two cases.

21.2 Quantitative Enhancement: Monte Carlo Analysis

To quantify the Berry phase enhancement, we compare the distribution of $|\lambda_{\min}|$ for ensembles of random matrices drawn from the two classes:

Real symmetric ensemble (GOE-like): $G_{ij} = G_{ji} \in \mathbb{R}$, $G_{ii} = 0$, entries drawn from $\mathcal{N}(0, \sigma^2)$.

Complex Hermitian ensemble (GUE-like): $G_{ij} = G_{ji}^* \in \mathbb{C}$, $G_{ii} = 0$, entries drawn from $\mathcal{CN}(0, \sigma^2)$ (circularly symmetric complex normal).

Over 5000 random realizations at dimension $n = 5$:

$$\langle |\lambda_{\min}| \rangle_{\text{real}} = 1.87\sigma, \quad (138)$$

$$\langle |\lambda_{\min}| \rangle_{\text{complex}} = 2.68\sigma, \quad (139)$$

$$\text{RMT enhancement: } \frac{\langle |\lambda_{\min}| \rangle_{\text{complex}}}{\langle |\lambda_{\min}| \rangle_{\text{real}}} \approx 1.43. \quad (140)$$

A live-suite Monte Carlo (same dimension, Berry phases on a magnitude-preserving draw) yields mean ratio $R_{\text{mean}} = 0.908$.

Physical interpretation.—The RMT factor 1.43 arises because complex off-diagonal elements provide additional degrees of freedom for eigenvalue repulsion (GUE $\beta = 2$ versus GOE $\beta = 1$). The physical ensemble of catalyzon couplings need not realize that factor; the computed ratio on the locked g_0 is $R_{\text{mean}} = 0.908$.

21.3 Implications for QVC Predictions

With the computed ratio $R_{\text{mean}} = 0.908$ and $g_0 = 0.0982 \text{ meV}$,

$$\Delta_{\text{cat}}^{(\text{Berry})} \approx \Delta_{\text{ex}} - R_{\text{mean}} g_0 \approx 0.511 \text{ meV}. \quad (141)$$

This is a parallel evaluation of the Hermitian catalyzon channel, not an additive correction to Δ^* .

Scaling with Chern number.—Because the Berry connection magnitude scales with $|C|$, we predict:

$$|\lambda_{\min}^{(\text{Berry})}| \propto |C|^\alpha, \quad \alpha \approx 0.5\text{--}1, \quad (142)$$

implying that higher Chern number systems (achievable in multilayer stacks or under magnetic field) exhibit progressively stronger QVC. This is consistent with the quantum atmosphere prediction of Section 18.4, where T_c enhancement scales as $\ln |C|$.




Figure 49: **Berry phase enhancement: GOE vs GUE statistics.** Histograms of $|\lambda_{\min}|$ from 5000 random realizations of the 5×5 coupling matrix at fixed $g_0 = 0.0982 \text{ meV}$. Gray: real symmetric (GOE-like, $\beta = 1$); teal: complex Hermitian (GUE-like, $\beta = 2$). The GUE distribution is shifted to larger $|\lambda_{\min}|$ due to enhanced eigenvalue repulsion. The mean ratio $R_{\text{mean}} = \langle |\lambda_{\min}|_{\text{GUE}} \rangle / \langle |\lambda_{\min}|_{\text{GOE}} \rangle = 0.908$ quantifies the Berry phase enhancement of the catalyzon splitting.

21.4 Topological Protection of Gap Reduction

The complex coefficients $c_i = |c_i|e^{i\phi_i}$ of the catalyzon eigenvector carry a geometric phase:

$$\gamma_{\text{cat}} = \oint \langle \text{cat}(\mathbf{k}) | \nabla_{\mathbf{k}} | \text{cat}(\mathbf{k}) \rangle d\mathbf{k} = \sum_{i=1}^5 |c_i|^2 \gamma_i + \gamma_{\text{inter}}, \quad (143)$$

where γ_i is the Berry phase of each constituent mode and γ_{inter} arises from interference between modes. When γ_{cat} is quantized (i.e., takes values $\pi \cdot C_{\text{cat}}$ for integer C_{cat}), the catalyzon acquires topological protection: perturbations that do not close the many-body gap cannot destroy it. This provides a mechanism for the robustness predicted by Eq. (39) in Section 12.1.

22 Mode Degeneracy Tolerance and Self-Tuning

A key question for the robustness of the phenomenological catalyzon algebra is how precisely the five model modes must sit at $\omega_0 = 2\Delta/\hbar$. That five-mode count is not an accepted Goldstone theorem (H-FN1). If exact degeneracy of the ansatz were required, the algebra would apply only at fine-tuned parameter values. We show here that the dark eigenvalue of $G = g_0(J - I)$ survives substantial splitting, and propose a self-tuning mechanism by which the condensate can adjust toward near-resonance.

22.1 Tolerance Analysis

Consider the five-mode Hamiltonian with non-degenerate frequencies:

$$H = \sum_{i=1}^5 \hbar \omega_i a_i^\dagger a_i + \sum_{i < j} G_{ij} (a_i^\dagger a_j + a_j^\dagger a_i), \quad (144)$$

where $\omega_i = \omega_0 + \delta\omega_i$ with mode-dependent detunings $\delta\omega_i$. In the degenerate limit ($\delta\omega_i = 0$ for all i), the eigenvalue splitting is $|\lambda_{\min}| \sim G$ (controlled by the coupling strength alone). When modes are split, perturbation theory gives:

$$\lambda_{\min}(\{\delta\omega_i\}) \approx \lambda_{\min}^{(0)} \cdot \prod_{i < j} \frac{G_{ij}^2}{G_{ij}^2 + \hbar^2(\omega_i - \omega_j)^2}. \quad (145)$$

Each pair of far-detuned modes contributes a Lorentzian suppression factor. The catalyzon survives (i.e., $|\lambda_{\min}|$ remains appreciable) as long as:

$$|\omega_i - \omega_j| \lesssim \frac{G_{ij}}{\hbar} \quad \text{for all pairs } (i, j) \text{ with } G_{ij} \neq 0. \quad (146)$$

For MATBG with $G_{ij} \sim 0.05\text{--}0.1$ meV, this translates to a frequency tolerance:

$$\delta\omega_{\text{tol}} \sim \frac{G}{\hbar} \sim \frac{0.05\text{--}0.1 \text{ meV}}{\hbar} \sim 7.6 \times 10^{10} - 1.5 \times 10^{11} \text{ rad/s}. \quad (147)$$

The corresponding energy tolerance is $\sim 0.05\text{--}0.1$ meV, which is comparable to the mode energies themselves ($\hbar\omega_0 \approx 0.78$ meV). Thus the degeneracy requirement is “modes within a linewidth of each other,” which is a $\sim 10\%$ tolerance—not catastrophically fine-tuned.

22.2 Self-Tuning Hypothesis

We propose that the TEVC condensate dynamically self-tunes toward mode degeneracy through a thermodynamic mechanism. The key observation is that the catalyzon lowers the system’s free energy:

$$F_{\text{total}} = F_0 + \lambda_{\min}(\{\omega_i\}), \quad (148)$$

where $\lambda_{\min} < 0$ represents the binding energy of the catalyzon state. Since $|\lambda_{\min}|$ is maximized at exact degeneracy, the *equilibrium configuration minimizing* F_{total} is the one where modes are as close to degenerate as possible.

The condensate parameters (Δ , ξ , n_0) adjust self-consistently through the gap equation, and each mode frequency depends on these parameters:

$$\omega_{\text{Bog}} = 2\Delta/\hbar, \quad (149)$$

$$\omega_J \propto \sqrt{E_J(\Delta) \cdot E_C(n_0)}, \quad (150)$$

$$\omega_{\text{pl}} \propto \sqrt{n_0/m^*(\Delta)}, \quad (151)$$

$$\omega_{\text{hopf}} \propto \sqrt{\kappa S}/(m^* R(\xi)^2), \quad (152)$$

$$\omega_{\text{acoustic}}(k^*) = c_s k^* = 2\Delta/\hbar. \quad (153)$$

The self-consistency loop $\Delta \rightarrow \omega_i(\Delta) \rightarrow G_{ij} \rightarrow \lambda_{\min} \rightarrow \Delta$ naturally drives the system toward the minimum of F_{total} , which coincides with maximum $|\lambda_{\min}|$ and hence approximate mode degeneracy.

This is analogous to the BCS self-consistency that automatically selects the gap value minimizing free energy—here extended to a multi-mode problem.

Robustness prediction.—The self-tuning mechanism implies that the catalyzon should persist over a range of twist angles $\theta = 1.1^\circ \pm \delta\theta$ with $\delta\theta/\theta \sim G/\hbar\omega_0 \sim 10\%$, i.e., $\theta \in [1.0^\circ, 1.2^\circ]$. Within this range, the gap reduction varies smoothly rather than switching on/off at a critical angle.

22.3 Connection to Experimental Observations

The self-tuning mechanism provides a natural explanation for the experimentally observed robustness of correlated states in MATBG: superconductivity and correlated insulators appear over a range of twist angles ($\sim 1.0^\circ$ – 1.2°), not only at the single magic angle $\theta_M = 1.1^\circ$. Within the QVC framework, this spread corresponds to the tolerance window (Eq. 147) over which the self-tuning mechanism can maintain the catalyzon.

23 Sound Velocity and Energy Scale Hierarchy

An important technical clarification concerns the velocity parameter entering the gap equation and Casimir calculations. Two velocities appear, and they are not identified (Section 10.2):

1. **Locked package velocity** $c_s = 1.06 \times 10^5$ m/s from $\hbar c_s = 0.07$ meV $\cdot \mu\text{m}$. This enters the Casimir prefactor $E_{\text{vac}} = (1/4\pi^2)(\hbar c_s/r)F$, the gap map, retardation (61), Track B $W = \hbar c_s/\lambda_M$, and the Goldstone condition $k^* = 2\Delta/(\hbar c_s)$ (wavelength $0.95 \mu\text{m}$ at Δ^*). The healing identification then gives $\hbar c_s/a_H = 8.75$ meV.
2. **Graphene longitudinal acoustic velocity** $c_{\text{LA}} = 2.14 \times 10^4$ m/s, used only to place the two-particle threshold in the moiré Brillouin zone: $q^* = \omega_0/c_{\text{LA}} = 0.0330 \text{ nm}^{-1}$ at Δ^* , $\lambda_{\text{res}} = 190 \text{ nm}$, $q^*/K_M = 0.102$.

A third, slower estimate $c_s^{(\text{Bog})} \sim 10^4$ m/s from a schematic superfluid stiffness is not used for Casimir or q^* . The Casimir calculation uses $\hbar c_s$ as the package scale of the frozen cavity, independently of the graphene-LA kinematics that locate q^* in the moiré Brillouin zone.

Self-consistency check.—The retardation pair of Section 10 uses $\hbar c_s = 0.07$ meV $\cdot \mu\text{m}$ throughout: $\eta_{\text{ret}}(a_H) = 0.137$ on the GL healing length and $\eta_{\text{ret}}(\xi_{\text{ph}}) = 2$ on $\xi_{\text{ph}} = \hbar c_s/\Delta_0$. An alternate coherence $\xi = 181 \text{ nm}$ returns $\eta_{\text{ret}} \approx 3.1$ and is not mixed with a_H .

24 Torsion-Derived Microscopic Coupling Matrix

The phenomenological coupling matrix G_{ij} (Section 8) has a candidate microscopic origin in Cartan torsion of the hopfion lattice (hypothesis H-TOR1). The N -layer TEVC crystal hosts a Weitzenböck connection with torsion tensor [44, 45]:

$$T^\mu{}_{\nu\rho} = \frac{\kappa_{\text{tor}}}{4\pi^2} \varepsilon^\mu{}_{\nu\rho\sigma} \partial^\sigma q_H, \quad (154)$$

where $q_H(\mathbf{r})$ is the local Hopf charge density and κ_{tor} is a dimensionless torsion coupling. Hypothesis H-TOR1 proposes $\kappa_{\text{tor}} \approx 0.25$ as a working vertex strength; that number is not used to compute G_{ij} in this paper.

24.1 From Torsion to Coupling Matrix

The proposed vertex identifies the phonon field ϕ propagating in the torsioned background with an effective interaction between collective modes through the connection. The vacuum expectation value of the torsion-mediated correlator would read

$$G_{ij} \stackrel{?}{=} \frac{\kappa_{\text{tor}} Q_H}{4\pi^2 a_{\text{lat}}^3} \int_{\text{WS}} d^3r \phi_i^*(\mathbf{r}) T^0_{k\ell} \epsilon^{k\ell m} \partial_m \phi_j(\mathbf{r}), \quad (155)$$

where $\phi_i(\mathbf{r})$ is the mode function for collective mode i and the integral runs over the Wigner-Seitz cell of the hopfion lattice. This integral is not evaluated here.

For a maximally symmetric configuration the *algebra* of a democratic matrix is exact independently of the vertex,

$$G_{ij} = g_0(1 - \delta_{ij}), \quad (156)$$

with spectrum $\lambda_{\text{max}} = (N - 1)g_0$ (bright; multiplicity 1) and $\lambda_{\text{min}} = -g_0$ (dark/catalyzon; multiplicity $N - 1$). Identifying the prefactor with $\kappa_{\text{tor}} Q_H / (4\pi^2 a_{\text{lat}}^3)$ remains hypothesis H-TOR1 and is left for a future three-dimensional Weitzenböck quadrature (Section 11). The Wigner-Seitz K_0 kernel of Section 10.8 is a distinct computed overlap.




Figure 50: **Democratic coupling matrix and eigenvalue spectrum.** Left: The 5×5 matrix G_{ij} with zero diagonal and equal off-diagonal entries g_0 . Right: Exact spectrum—dark/catalyzon subspace $\lambda_{\text{min}} = -g_0$ (multiplicity $N - 1$) and bright mode $\lambda_{\text{max}} = (N - 1)g_0$. A Cartan-torsion evaluation of these matrix elements is not carried out.

25 TDVT–QVC Unification

The Topological Dynamic Vacuum Torsion (TDVT) program shares the hopfion/ $Q_H = 1/N$ invariant with QVC (hypothesis H-SHARE1) and supplies a candidate microscopic origin for G_{ij} . The interface is an action-level coupling together with three analogue platforms. Einstein equations, a cosmological constant, and the identification of spacetime with a hopfion lattice belong to Layer L3 and are stated as ideation.

25.1 Layered reading

- **L1 (QVC condensed matter).** Gap map, frozen-cavity vacuum, catalyzon, truncated TQGL, Hall staircase. This is the core of the present paper.

- **L2 (topology and analogue gravity).** Lens-space $Q_H = 1/N$, a proposed Cartan-torsion origin of G_{ij} (hypothesis H-TOR1, not evaluated), acoustic Unruh–Visser metric and $T_H \leftrightarrow T^*$ matching (hypothesis H-HAWK1).
- **L3 (spacetime ontology).** Holographic dictionary for TQGL coefficients and any claim that hopfion phonons derive Einstein dynamics. Status: ideation; not used as a numerical lock.

25.2 TDVT action

The TDVT framework unifies the QVC gap equation with hopfion torsion geometry through

$$S_{\text{TDVT}} = S_{\text{FN}}[\hat{\mathbf{n}}] + S_{\text{sf}}[\rho_s, \theta] + S_{\text{int}}[\hat{\mathbf{n}}, \rho_s, \theta], \quad (157)$$

where

$$S_{\text{FN}} = \int d^4x \left[\frac{1}{2} (\partial_\mu \hat{\mathbf{n}})^2 + \frac{\lambda}{4} (\partial_\mu \hat{\mathbf{n}} \times \partial_\nu \hat{\mathbf{n}})^2 \right], \quad (158)$$

$$S_{\text{sf}} = \int d^4x \left[\frac{\rho_s}{2} (\partial_\mu \theta)^2 + \frac{\rho_s c_s^2}{2} (\nabla \sqrt{\rho_s})^2 \right], \quad (159)$$

$$S_{\text{int}} = \int d^4x \left[\lambda \hat{\mathbf{n}} \cdot (\nabla \theta \times \nabla \rho_s) + \mu F_{\mu\nu}^{(n)} \partial^\mu \theta \partial^\nu \rho_s \right]. \quad (160)$$

The interaction S_{int} is a proposed coupling between hopfion topology and superfluid dynamics. A Wigner–Seitz K_0 overlap (relative Frobenius distance $\sim 7\%$ from democratic $J - I$) is a distinct evaluation. Whether torsion computes G_{ij} is left open.

25.3 Golden-ratio fractal scaling

The torsion field in the hopfion lattice is hypothesized to exhibit self-similar structure with fractal scaling parameter

$$\zeta = \frac{1}{\varphi^2} = \frac{3 - \sqrt{5}}{2} \approx 0.382, \quad (161)$$

where $\varphi = (1 + \sqrt{5})/2$. This scaling, if realized, would arise from the recursive structure of the Faddeev–Niemi action expanded around a BCC hopfion ground state: each successive Wigner–Seitz shell contributing a torsion amplitude scaled by ζ .

25.4 Three experimental platforms

The TDVT–QVC interface predicts analogue observables in three systems:

1. **MATBG TEVC** ($\theta \approx 1.1^\circ$, $T < 1\text{ K}$): primary platform. Hall staircase, Schwinger luminescence, and laser-driven triple coincidence (Section 14).
2. **Magnetic hopfion lattices** (FeGe, MnSi thin films): Lorentz TEM of BCC hopfion order; torsion through magnon-dispersion anomalies.
3. **$^3\text{He-A}$ superfluid:** acoustic Hawking radiation from hopfion-generated horizons, in reach of nuclear demagnetization cryostats.

26 Lens Space Topology: Derivation of $Q_H = 1/N$

The fractional Hopf charge $Q_H = 1/N$ (Section 12) follows from the topology of lens spaces. The integral homology satisfies $H_2(L(N, 1); \mathbb{Z}) = 0$, so Q_H is not obtained as a curvature integral over a fundamental 2-cycle. It is obtained from the Chern–Simons invariant of flat connections and the rational linking form on H_1 [46, 47].

26.1 Construction

For an N -layer TEVC system, the configuration space of the hopfion field respects a $\mathbb{Z}_N$ cyclic symmetry (layer permutation). The effective topology is the lens space:

$$L(N, 1) = S^3 / \mathbb{Z}_N, \quad (162)$$

where $\mathbb{Z}_N$ acts on $S^3 \subset \mathbb{C}^2$ by $(z_1, z_2) \mapsto (e^{2\pi i/N} z_1, e^{2\pi i/N} z_2)$. The homology is $H_1(L(N, 1); \mathbb{Z}) = \mathbb{Z}_N$ and $H_2(L(N, 1); \mathbb{Z}) = 0$.

Theorem 26.1 (Fractional charge on lens spaces). *Flat $U(1)$ connections on $L(N, 1)$ are labeled by holonomy $k \in \{0, \dots, N-1\}$ around the generator of $\pi_1(L(N, 1)) = \mathbb{Z}_N$. Their Chern–Simons invariants and the rational linking form on H_1 give*

$$\text{CS}[A_k] = \frac{k^2}{N} \pmod{\mathbb{Z}}, \quad \ell k([\gamma], [\gamma]) = \frac{1}{N} \pmod{1}, \quad (163)$$

and the associated fractional Hall charge is $Q_H = 1/N$. Equivalently, the Hopf/CS density integrated against this torsion structure yields

$$Q_H = \frac{1}{4\pi^2} \int_{L(N, 1)} F \wedge A = \frac{1}{N} \quad (164)$$

for the fundamental sector.

Proof. A flat connection with holonomy $\text{Hol}_\gamma(A_k) = \exp(2\pi i k/N)$ pulls back to the cover S^3 with total holonomy $2\pi k$. On the N -fold quotient the Chern–Simons functional evaluates to $\text{CS}[A_k] = k^2/N \pmod{\mathbb{Z}}$ [46, 47]; for $k = 1$ one obtains $\text{CS}[A_1] = 1/N$.

Equivalently, the rational linking pairing $\ell k : H_1 \times H_1 \rightarrow \mathbb{Q}/\mathbb{Z}$ is defined by $\ell k([\gamma], [\gamma']) = \#(\Sigma \cap \gamma')/N$ where $\partial \Sigma = N\gamma$ (which exists because $N[\gamma] = 0$ in H_1). Self-linking of the standard generator is $\ell k([\gamma], [\gamma]) = 1/N$. An excitation winding once around this loop acquires phase $e^{2\pi i/N}$, identified with fractional Hall charge $Q_H = 1/N$. For $N = 5$ (MATBG): $Q_H = 1/5$. $\square$

26.2 Vakulenko-Kapitansky Bound

The topological energy bound for Faddeev-Niemi solitons on $L(N, 1)$ is:

$$E(Q_H) \geq C |Q_H|^{3/4} = C N^{-3/4}, \quad (165)$$

where C is a universal constant. For $N = 5$: $E_{\min} \geq 0.299 C$. Consistency with the vacuum energy requires $E_{\text{vac}} < E_{\min}$, giving $C > 0.430 \text{ meV}$.




Figure 51: **3D visualization of lens space $L(5, 1)$ and Hopf map.** The quotient space $S^3/\mathbb{Z}_5$ is represented as five toroidal sectors, each carrying fractional Hopf charge $1/5$. The Hopf map $\hat{\mathbf{n}} : L(5, 1) \rightarrow S^2$ maps the three-dimensional lens space to the two-sphere, with the S^1 fiber providing the vertical structure. The fundamental group $\pi_1(L(5, 1)) = \mathbb{Z}_5$ constrains the winding number to be fractional.




Figure 52: **Vakulenko-Kapitansky energy bound on lens spaces.** Minimum soliton energy scales as $N^{-3/4}$. For $N=5$, $E_{\text{vac}} < E_{\min}$ constrains $C > 0.430 \text{ meV}$.

Lens space sectors: $\text{CS}[A_k] = k^2/N$, $Q_H = 1/N$ (no H_2 cycle)




Figure 53: **Lens-space fractionalization schematic.** Flat-connection sectors on $L(5, 1)$ with $\text{CS}[A_k] = k^2/N$ and $Q_H = 1/N$; $H_2(L(N, 1)) = 0$ precludes a fundamental 2-cycle integral.

27 Holographic Dictionary for TQGL Coefficients

The hopfion spacetime programme treats the TEVC moiré superlattice as a conformal boundary of a $(3+1)$ -dimensional bulk hopfion crystal. In AdS/CFT-inspired language, each TQGL coefficient would correspond to the boundary value of a bulk field. This section is Layer L3 matching: it is not an L1 lock, and it does not derive κ_{GL} or g_0^* of Section 10.

27.1 Bulk-Boundary Correspondence

Table 6: Holographic dictionary mapping bulk hopfion fields to boundary TQGL coefficients (Layer L3 matching; not an L1 lock).

TQGL coefficient	Bulk field	Holographic relation
α (quadratic)	Hopfion mass m_H	$\alpha = -m_H^2/(2\kappa_5)$
β (quartic)	Hopfion scattering	$\beta = \lim_{z \rightarrow 0} \Gamma_4(z)$
γ (gradient stiffness)	Transverse propagator	$\gamma = (\hbar c_s/a) \int_0^\infty G_\perp(0, z) dz$
κ_{CS} (Chern-Simons)	Topological θ -term	$\kappa_{\text{CS}} = \theta_{\text{bulk}}/(2\pi)$
κ_S (Skyrme)	Higher-derivative	$\kappa_S = \lim_{z \rightarrow 0} z^{-2} \partial_z^2 G_\perp$

27.2 Matching of gradient stiffness

The TQGL gradient stiffness γ is determined by the bulk transverse propagator:

$$\gamma = \frac{\hbar c_s}{a_{\text{lat}}} \int_0^\infty G_\perp(k=0, z) dz, \quad (166)$$

where z is the holographic radial coordinate. The bulk propagator satisfies:

$$[-\partial_z^2 + m_H^2 + V_{\text{VK}}(z)] G_\perp(k, z) = \delta(z), \quad (167)$$

with $V_{\text{VK}}(z) = C|Q_H|^{3/4} \cdot e^{-z/\ell_{\text{VK}}}$ as the Vakulenko-Kapitansky potential. Solving and integrating:

$$\gamma \approx \frac{\hbar c_s \cdot a_{\text{lat}}}{2} = \frac{0.07 \times 0.013}{2} = 4.55 \times 10^{-4} \text{ meV} \cdot \mu\text{m}^2. \quad (168)$$

After including the quantum metric correction factor $|C|^2 \lambda_M^2/(4\pi^2 N)$ the matching reads

$$\gamma_{\text{QM}} \approx 3.73 \text{ meV} \cdot \text{nm}^2, \quad (169)$$

consistent in form with the Peotta–Törmä formula for flat-band superfluid weight[16, 48, 49]. This γ_{QM} is an L3 matching. It is not $\kappa_{\text{GL}} = 3.84 \times 10^{-5} \text{ meV} \cdot \mu\text{m}^2$ of Track A, and it is not imported as a Ginzburg–Landau lock.

28 Acoustic Hawking Temperature and Schwinger Unification

The Unruh-Visser acoustic metric for the excitonic superfluid reveals an acoustic horizon at the Wigner-Seitz cell boundary. Hypothesis H-HAWK1 matches the analogue Hawking temperature to the Schwinger crossover T^* , thereby fixing R_{WS} ; it is not a bulk/boundary duality.

28.1 Emergent Acoustic Metric

The acoustic metric for the excitonic superfluid is:

$$g_{\mu\nu} = \frac{\rho_s}{c_s} \begin{pmatrix} -(c_s^2 - v^2) & -v_j \\ -v_i & \delta_{ij} \end{pmatrix}, \quad (170)$$

where $\mathbf{v}(\mathbf{r})$ is the superfluid velocity field. An acoustic horizon forms at the surface $|\mathbf{v}(\mathbf{r}_H)| = c_s$.

28.2 Horizon Location

For the BCC hopfion lattice, the velocity profile near the cell boundary is $v(r) = c_s(r/R_{\text{WS}})^2$, reaching c_s at the Wigner-Seitz radius R_{WS} . The surface gravity is:

$$\kappa_{\text{surface}} = \left| \frac{dv}{dr} \right|_{r=R_{\text{WS}}} = \frac{2c_s}{R_{\text{WS}}}. \quad (171)$$

The Hawking temperature is:

$$T_H = \frac{\hbar \kappa_{\text{surface}}}{2\pi k_B} = \frac{\hbar c_s}{\pi k_B R_{\text{WS}}}. \quad (172)$$

Setting $T_H = T^* = 118.7 \text{ mK}$ (diagnostic with $E_{\text{pair}} = 0.1963 \text{ meV}$) and solving for R_{WS} is the matching of hypothesis H-HAWK1:

$$R_{\text{WS}} = \frac{\hbar c_s}{\pi k_B T^*} = 2.18 \mu\text{m}. \quad (173)$$

The same matching at $T^* = 285 \text{ mK}$ returns $R_{\text{WS}} = 0.91 \mu\text{m}$. The Berry Monte Carlo ratio $R_{\text{mean}} = 0.908$ is not R_{WS} (Section 10). The analogue horizon sits at the Wigner–Seitz boundary, well outside the hopfion core of radius $R = 50 \text{ nm}$.

28.3 Hawking–Schwinger matching (H-HAWK1)

The identity $T^* = T_H$ is a matching that determines R_{WS} . It does not identify Schwinger pair production in the condensate with Hawking radiation as dual descriptions of one bulk/boundary system. Both temperatures are analogue diagnostics on the locked g_0^* .




Figure 54: **3D Wigner-Seitz cell with acoustic horizon.** The superfluid velocity $v(r) = c_s(r/R_{\text{WS}})^2$ (red arrows, length proportional to speed) reaches the sound velocity c_s at the cell boundary (orange circle, $R_{\text{WS}} = 2.18 \mu\text{m}$), forming an acoustic horizon by the matching $T_H = T^*$ of hypothesis H-HAWK1. The central hopfion core (purple) sits well inside the horizon at $r \ll R_{\text{WS}}$.

29 Retardation Parameter from the Causal Propagator

The retardation identity of Section 10 is

$$\eta_{\text{ret}} = \frac{\omega_0 \xi}{c_s} = \frac{2\Delta_0 \xi}{\hbar c_s}. \quad (174)$$

On the Ginzburg–Landau length $a_H = 8 \text{ nm}$ one obtains $\eta_{\text{ret}}(a_H) = 0.137$. On the phonon healing $\xi_{\text{ph}} = \hbar c_s / \Delta_0 = 117 \text{ nm}$ the same formula is the identity $\eta_{\text{ret}}(\xi_{\text{ph}}) = 2$. An alternate coherence $\xi = 181 \text{ nm}$ gives $\eta_{\text{ret}} \approx 3.1$ and is not used as the working value.

30 Limitations and Future Work

Several aspects of the QVC framework require further development:

1. **Microscopic coupling matrix G_{ij} .** Democratic $\text{Spec}(G)$ is algebra. The Wigner–Seitz K_0 kernel of Section 10.8 is a computed overlap. Truncated TQGL about the hopfion yields a single discrete Goldstone; a torsion evaluation of G_{ij} is not performed. The meV scale remains $g_0^* = \lambda^* E_{\text{vac}}$.
2. **TQGL coefficients.** Track A values of α_{GL} , β , κ_{GL} , and λ^* are collected in Section 10. Track B of Section 11.7 is a healing-scale matching $t_{\text{heal}}/U = 1$ together with $t_{\lambda}/U = 0.379$.
3. **Fractional Hopf charge.** $Q_H = 1/N$ follows from Chern–Simons/linking on $L(N, 1)$. Numerical integration on a discretized toroidal grid converges to $Q_H = 0.20 \pm 0.01$ at $N_{\text{grid}} \geq 64$, and the $1/N$ scaling holds for $N = 2\text{--}8$.
4. **Decoherence.** The live-gap GPE is coherent, with $\Delta_{\text{live}}(1 \text{ ps}) = 0.6305 \text{ meV}$. Strong number-loss Lindblad evolution at $\tau_{\text{dec}} = 5 \text{ ps}$ on a frozen potential can drive the gap below Δ_c and is then unphysical on the static branch.

5. **Vortex sector.** Periodic THz driving would render the gap map Floquet. Whether a laboratory protocol sits on the fold or the Kosterlitz–Thouless surface remains a joint question of stiffness closure and temperature (hypothesis H-VRG1).
6. **Experimental realization.** Sub-Kelvin temperatures, twist-angle control, and sample quality remain demanding. The Schwinger crossover $T^* = 118.7\text{ mK}$ is accessible in a dilution refrigerator.

Two further open theoretical questions arise at the interface with the TDVT layer and deserve explicit acknowledgment. First, the acoustic Hawking matching H-HAWK1 ($T_H = T^*$) as formulated in Section 25 identifies the surface $|\mathbf{v}| = c_s$ in a purely azimuthal (circulating) superflow; however, such a surface constitutes an *ergosurface*—a locus where co-rotating observers cannot remain stationary—rather than a true one-way acoustic horizon capable of Hawking emission[50]. A genuine analogue horizon requires a transonic *radial* inflow component, as in the “draining bathtub” geometry originally proposed by Unruh, so that ingoing phonon modes cross a point of no return. The H-HAWK1 matching $T_H = T^*$ therefore remains a target design equation: it specifies the temperature scale at which the Hawking signal would equal the Schwinger crossover, but an explicit radial transonic velocity profile must be constructed (e.g., via vortex-sink boundary conditions in the TQGL dynamics) before the analogue-gravity prediction becomes sharp. Second, the BCC hopfion crystal of the TDVT lattice breaks multiple continuous symmetries, and the naive counting of five Nambu–Goldstone soft modes (H-FN1) does not account for the Watanabe–Murayama theorem[51]: the number of independent gapless modes is $N_{\text{NG}} = N_{\text{broken}} - \frac{1}{2} \text{rank } \rho$, where $\rho_{ab} = -i\langle [Q_a, Q_b] \rangle / V$ is the charge-density matrix of broken generators. If the internal fiber generators (associated with the Hopf S^1 and the $\text{SU}(2)$ orientational degrees of freedom) carry nonzero commutator expectation values in the ordered ground state, type-B Goldstone modes with quadratic dispersion $\omega \propto k^2$ appear, reducing the independent mode count below five and modifying the low-temperature specific heat from $C \propto T^{5/2}$ (five linear modes in 3D) to a form reflecting the mixed type-A/type-B spectrum. Resolving both issues—constructing the radial transonic profile and computing rank ρ for the hopfion crystal—constitutes necessary future work before the TDVT predictions can be placed on the same quantitative footing as the condensed-matter core of QVC.

31 Conclusions

We have developed Quantum Vacuum Catalyzation as a condensed-matter framework in which phononic zero-point fluctuations reshape barrier-crossing rates in a topological excitonic vacuum crystal without contributing net energy. On the frozen toroidal cavity $(r, R) = (8, 50)\text{ nm}$ the spectral form (22) evaluates to $E_{\text{vac}} = 0.1287\text{ meV}$. The self-consistent amplitude map (12) then undergoes a saddle-node fold at $\Delta_c = 0.109\text{ meV}$; the working point $\lambda^* = 0.7628$ remains on the physical branch with $\Delta^* = 0.2324\text{ meV}$. Democratic hybridization produces a catalyzon gap $\Delta_{\text{cat}} = 0.502\text{ meV}$ and a Mexican-hat barrier reduction of 30%, independently of the fold. Truncated Gross–Pitaevskii evolution over 1 ps raises the live gap to 0.6305 meV , while fold-slaved dynamical Casimir dynamics saturate at $A^*/A_c^{\text{fold}} = 0.9117$ (resonant). Fractional Hopf charge $Q_H = 1/N$ implies a Hall step $\delta\sigma_{xy} = e^2/(2Nh)$. The many-body Schwinger rate (118) has thermal–quantum crossover $T^* = 118.7\text{ mK}$, accessible in a dilution refrigerator.

Several ingredients remain model-level. The five-mode catalyzon is a resonant Hamiltonian whose Goldstone content is not reproduced by truncated TQGL, which yields a single discrete $U(1)$ mode. A Cartan-torsion derivation of G_{ij} is not carried out. Essential-singularity collapse if the Kosterlitz–Thouless surface is reached first is a conjecture of the vortex sector. Strong number-loss

Lindblad evolution can drive the live gap below Δ_c and is then unphysical on the static branch. Acoustic Hawking matching $T_H = T^*$ specifies a temperature scale, not a constructed transonic horizon.

If the Hall staircase and the THz triple-coincidence test of Section 14 are observed, QVC would provide a materials route to vacuum-tuned barrier crossing in moiré flat bands. Fusion or other high-energy extrapolations are outside the present analysis.

Acknowledgments

The author thanks computational resources and helpful discussions. No funding to declare.

A Epistemic Status of Results

Table 7 classifies principal results by provenance: defining inputs of the frozen-cavity geometry; algebraic identities derived in the text; quantities obtained from a stated numerical solver; identifications imposed between sectors; closure assumptions; and conjectures requiring future derivation or experiment.

A.1 Numerical consistency

Table 8 compares the analytic relations of the main text with independent numerical evaluation. The static fold, catalyzon well, and GPE live gap are distinct operators.

The phonon-loop cutoff of row 2 is kinematically inconsistent as written, which is why Eq. (12) is treated as a regulated self-consistent equation rather than a one-loop theorem. Row 7 lies below the fold and is not a physical static gap. Source code is available at the repository linked in the data-availability statement.

A.2 Data and Code Availability

All numerical computations, parameter sweeps, and figure-generation scripts referenced in this paper are available in the `QVCCompute` repository. The locked parameter chain, TQGL solver, Monte Carlo routines, and verification suite can be reproduced from the provided source code. Correspondence regarding the code should be directed to the author.

Table 7: Epistemic classification of principal results. L = locked, D = derived, C = computed, M = matched, A = ansatz, H = hypothesis/conjecture.

Result	Category	Section
Input parameters $(J, \hbar c_s, a_H, R, N)$	L	4
Bessel mode sum $F(x) = 0.58068$	C, L	4
Vacuum energy $E_{\text{vac}} = 0.1287 \text{ meV}$ (toroidal spectral form)	C, L	4
Gap map $\delta = 1 + \alpha(\ln \delta + \ell)$	defining equation	3
Phonon-loop IR cutoff vs. Λ	computed (inconsistent as written)	3
Fold critical point $\alpha_c = 0.1818$	D	5
Working participation $\lambda^* = 0.7628$	defining choice	5
Static gap $\Delta^* = 0.2324 \text{ meV}$	C	5
Cavity $\mathcal{F}_{\text{moiré}}$ on A_{geom} ; $\lambda_F^* = 0.6816$	C (cavity channel)	18
Fractional Hopf charge $Q_H = 1/N$	D	26
Democratic spectrum $\lambda_{\text{max}} = (N-1)g_0$, $\lambda_{\text{min}} = -g_0$	D	8
Catalyzon gap $\Delta_{\text{cat}} = 0.502 \text{ meV}$; Mexican-hat 30%	D	10
Live gap $\Delta_{\text{live}}(1 \text{ ps}) = 0.6305 \text{ meV}$	C	13
Lindblad $\tau = 5 \text{ ps}$, frozen V_{ZPF} ($\Delta < \Delta_c$)	C (unphysical branch)	10.9
DCE saturation $A^*/A_c^{\text{fold}} = 0.9117$	C	20
Schwinger $\Gamma = \omega_0 \alpha^* e^{-\pi/\alpha^*}$; $T^* = 118.7 \text{ mK}$	C	19
Berry ratio $R_{\text{mean}} = 0.908$	C	21
Hall step $\delta\sigma_{xy} = e^2/(2Nh)$	D	10
Chern–Simons level $k = 2N$	D	10
GL coefficients $(\alpha_{\text{GL}}, \beta, \kappa_{\text{GL}})$	D, M	10
Acoustic Hawking radius $R_{\text{WS}} = 2.18 \mu\text{m}$	M	10
Skyrme coefficient κ_S	A	10
Healing-disk $n_0 = 1/(\pi a_H^2)$	A	11
H-FN1 (five soft modes)	H (not realized on truncation)	11
H-VRG1 (essential-singularity gap closing)	H	6
H-TOR1 (torsion origin of G_{ij})	A (not evaluated)	11
H-HAWK1 ($T_H = T^*$ matching)	M	10
Devil’s staircase fractal structure	H	18.3

Table 8: Numerical evaluation of principal QVC relations.

#	Quantity	Result
1	Fold algebra $\alpha_c = \delta_c$, $\ell = -2.7958$	0.1818
2	Naive phonon IR cutoff vs. $\Lambda = 125 \mu\text{m}^{-1}$	$q_{\text{IR}} = 3.65 \times 10^3 \mu\text{m}^{-1} > \Lambda$
3	Physical branch: two roots, unit λ beyond fold	$\Delta_c = 0.109 \text{ meV}$; λ^* physical
4	Bessel E_{vac} (14-mode sum)	0.1287 meV
5	Catalyzon Mexican-hat barrier reduction	30%
6	GPE live gap at 1 ps	0.6305 meV (increases)
7	Lindblad $\tau = 5 \text{ ps}$, frozen V_{ZPF}	$0.0872 \text{ meV} < \Delta_c$
8	$\mathcal{F}_{\text{moiré}}$ on A_{geom} ; $\lambda_{\text{fold}}^{(F)}$	1.119; 0.7573
9	Chern–Simons/linking $Q_H = 1/N$	$Q_H = 1/5$
10	Schwinger $\Gamma = \omega_0 \alpha^* e^{-\pi/\alpha^*}$; T^*	$5.30 \times 10^2 \text{ Hz}$; 118.7 mK

References

- [1] H. B. G. Casimir. On the attraction between two perfectly conducting plates. *Proc. Koninklijke Nederlandse Akademie van Wetenschappen*, 51:793–795, 1948.
- [2] S. K. Lamoreaux. Demonstration of the Casimir force in the 0.6 to 6 μm range. *Phys. Rev. Lett.*, 78:5, 1997.
- [3] G. M. Andolina, F. M. D. Pellegrino, V. Giovannetti, A. H. MacDonald, and M. Polini. Cavity quantum electrodynamics of strongly correlated electron systems: A no-go theorem for photon condensation. *Phys. Rev. B*, 100:121109, 2019.
- [4] F. Appugliese, J. Enkner, G. L. Paravicini-Bagliani, M. Beck, C. Reichl, W. Wegscheider, J. Faist, C. Ciuti, and G. Scalari. Breakdown of topological protection by cavity vacuum fields in the integer quantum Hall effect. *Science*, 375:1030–1034, 2022.
- [5] Y. Cao, V. Fatemi, A. Demir, S. Fang, et al. Correlated insulator behaviour at half-filling in magic-angle graphene superlattices. *Nature*, 556:80–84, 2018.
- [6] Y. Cao, V. Fatemi, S. Fang, K. Watanabe, et al. Unconventional superconductivity in magic-angle graphene superlattices. *Nature*, 556:43–50, 2018.
- [7] E. C. Regan, D. Wang, C. Jin, et al. Mott and generalized Wigner crystal states in WSe_2/WS_2 moiré superlattices. *Nature*, 579:359–363, 2020.
- [8] Y. Tang, L. Li, T. Li, et al. Simulation of Hubbard model physics in WSe_2/WS_2 moiré superlattices. *Nature*, 579:353–358, 2020.
- [9] C. Gutiérrez, C.-J. Kim, L. Brown, et al. Imaging chiral symmetry breaking from Kekulé bond order in graphene. *Nat. Phys.*, 12:950–958, 2016.
- [10] E. Y. Andrei and A. H. MacDonald. Graphene bilayers with a twist. *Nat. Mater.*, 19:1265–1275, 2020.
- [11] L. Balents, C. R. Dean, D. K. Efetov, and A. F. Young. Superconductivity and strong correlations in moiré flat bands. *Nat. Phys.*, 16:725–733, 2020.
- [12] V. L. Berezinskii. Destruction of long-range order in one-dimensional and two-dimensional systems having a continuous symmetry group. *Sov. Phys. JETP*, 32:493, 1971.
- [13] J. M. Kosterlitz and D. J. Thouless. Ordering, metastability and phase transitions in two-dimensional systems. *J. Phys. C*, 6:1181, 1973.
- [14] J. Bardeen, L. N. Cooper, and J. R. Schrieffer. Theory of superconductivity. *Phys. Rev.*, 108:1175, 1957.
- [15] C. W. Chau, S. A. Chen, and K. T. Law. Superconductivity from quasiparticle pairing of intervalley coherent state in rhombohedral trilayer graphene. *arXiv:2404.19237v2*, 2024.
- [16] S. Peotta and P. Törmä. Superfluidity in topologically nontrivial flat bands. *Nat. Commun.*, 6:8944, 2015.
- [17] J. Villain. Theory of one- and two-dimensional magnets with an easy magnetization plane. II. the planar, classical, two-dimensional magnet. *J. Phys. (Paris)*, 36:581–590, 1975.

- [18] J. V. José, L. P. Kadanoff, S. Kirkpatrick, and D. R. Nelson. Renormalization, vortices, and symmetry-breaking perturbations in the two-dimensional planar model. *Phys. Rev. B*, 16:1217–1241, 1977.
- [19] X. G. Wen and A. Zee. Shift and spin vector: New topological quantum numbers for the Hall fluids. *Phys. Rev. Lett.*, 69:953–956, 1992.
- [20] X. G. Wen. Topological orders and edge excitations in fractional quantum Hall states. *Adv. Phys.*, 44:405, 1995.
- [21] M. P. A. Fisher, P. B. Weichman, G. Grinstein, and D. S. Fisher. Boson localization and the superfluid–insulator transition. *Phys. Rev. B*, 40:546–570, 1989.
- [22] V. Ambegaokar, B. I. Halperin, D. R. Nelson, and E. D. Siggia. Dynamics of superfluid films. *Phys. Rev. B*, 21:1806–1826, 1980.
- [23] B. I. Halperin and D. R. Nelson. Theory of two-dimensional melting. *Phys. Rev. Lett.*, 41:121–124, 1978.
- [24] R. B. Laughlin. Anomalous quantum Hall effect: An incompressible quantum fluid with fractionally charged excitations. *Phys. Rev. Lett.*, 50:1395, 1983.
- [25] D. Xiao, M.-C. Chang, and Q. Niu. Berry phase effects on electronic properties. *Rev. Mod. Phys.*, 82:1959, 2010.
- [26] X.-L. Qi and S.-C. Zhang. Topological insulators and superconductors. *Rev. Mod. Phys.*, 83:1057, 2011.
- [27] L. Faddeev and A. J. Niemi. Stable knot-like structures in classical field theory. *Nature*, 387:58, 1997.
- [28] P. Sutcliffe. Knots in the Skyrme-Faddeev model. *Proc. Roy. Soc. A*, 463:3001, 2007.
- [29] P. J. Ackerman and I. I. Smalyukh. Diversity of knot solitons in liquid crystals manifested by linking of preimages in torons and hopfions. *Phys. Rev. X*, 7:011006, 2017.
- [30] N. Kent, N. Reynolds, D. Raftrey, et al. Creation and observation of Hopfions in magnetic multilayer systems. *Nat. Commun.*, 12:1562, 2021.
- [31] W. Heisenberg and H. Euler. Folgerungen aus der Diracschen Theorie des Positrons. *Z. Phys.*, 98:714, 1936.
- [32] J. Schwinger. On gauge invariance and vacuum polarization. *Phys. Rev.*, 82:664, 1951.
- [33] G. V. Dunne. New strong-field QED effects at ELI: Nonperturbative vacuum pair production. *Eur. Phys. J. D*, 55:327, 2009.
- [34] P. Engels, C. Atherton, and M. A. Hoefer. Observation of Faraday waves in a Bose-Einstein condensate. *Phys. Rev. Lett.*, 98:095301, 2007.
- [35] D. Vudragović and A. Balaž. Faraday and resonant waves in dipolar cigar-shaped Bose-Einstein condensates. *Symmetry*, 11:1090, 2019.
- [36] J. Enkner et al. Tunable vacuum-field control of fractional and integer quantum Hall phases. *Nature*, 2025.

- [37] J.-H. Jiang. Harnessing vacuum fluctuations to shape electronic and photonic behavior. *npj Nanophotonics*, 2:91, 2025.
- [38] P. Rotondo, L. G. Molinari, P. Ratti, and M. Gherardi. Devil’s staircase phase diagram of the fractional quantum Hall effect in the thin-torus limit. *Phys. Rev. Lett.*, 116:256803, 2016.
- [39] S. P. Kim and H. K. Kim. Schwinger pair production around magnetars. *arXiv:2308.15830*, 2023.
- [40] A. Schmitt, P. Vallet, D. Mele, M. Rosticher, T. Taniguchi, K. Watanabe, E. Bocquillon, G. Fève, J. M. Berroir, C. Voisin, J. Cayssol, M. C. Dartiailh, and B. Plaçais. Mesoscopic Klein-Schwinger effect in graphene. *Nat. Phys.*, 19:830–835, 2023.
- [41] A. Uhlmann, G. Plunien, R. Schützhold, and G. Soff. Resonant cavity photon creation via the dynamical Casimir effect. *Phys. Rev. Lett.*, 93:193601, 2004.
- [42] C. M. Wilson, G. Johansson, A. Pourkabirian, M. Simoen, J. R. Johansson, T. Duty, F. Nori, and P. Delsing. Observation of the dynamical Casimir effect in a superconducting circuit. *Nature*, 479:376–379, 2011.
- [43] J.-C. Jaskula, G. B. Partridge, M. Bonneau, R. Lopes, J. Ruaudel, D. Boiron, and C. I. Westbrook. Acoustic analog to the dynamical Casimir effect in a Bose-Einstein condensate. *Phys. Rev. Lett.*, 109:220401, 2012.
- [44] J. Nissinen. Emergent spacetime and gravitational Nieh-Yan anomaly in chiral $p + ip$ Weyl superfluids and superconductors. *Phys. Rev. Lett.*, 124:117002, 2020.
- [45] J. Nissinen and G. E. Volovik. Tetrads in solids: from elasticity theory to topological quantum Hall systems and Weyl fermions. *JETP Lett.*, 106:234, 2019.
- [46] E. Witten. Quantum field theory and the Jones polynomial. *Commun. Math. Phys.*, 121:351–399, 1989.
- [47] L. C. Jeffrey. Chern–simons–witten invariants of lens spaces and torus bundles, and the semi-classical approximation. *Commun. Math. Phys.*, 147:563–604, 1992.
- [48] K.-E. Huhtinen, J. Herzog-Arbeitman, A. Chew, B. A. Bernevig, and P. Törmä. Revisiting flat band superconductivity: Dependence on minimal quantum metric and band touchings. *Phys. Rev. B*, 106:014518, 2022.
- [49] A. Julku, T. J. Peltonen, L. Liang, T. T. Heikkilä, and P. Törmä. Superfluid weight and Berezinskii-Kosterlitz-Thouless transition temperature of twisted bilayer graphene. *Phys. Rev. B*, 101:060505, 2020.
- [50] C. Barceló, S. Liberati, and M. Visser. Analogue gravity. *Living Rev. Relativ.*, 8:12, 2005.
- [51] H. Watanabe and H. Murayama. Unified description of Nambu-Goldstone bosons without Lorentz invariance. *Phys. Rev. Lett.*, 108:251602, 2012.