

Validation of the **MadAnalysis 5** implementation of CMS-EXO-16-012

Jiwon Park, Seohyun Ahn (Hanyang Unibversity)
Wenxing Zhang (ITP of Chinese Academy of Science)

email: jiwon.park@cern.ch

August 26, 2017

1 Setup

In this document, the **MadAnalysis 5** implementation of search for associated production of dark matter with a Higgs boson decaying to $b\bar{b}$ or $\gamma\gamma$ at $\sqrt{s} = 13$ TeV (2.3 fb^{-1}), (see also arXiv:1703.05236) is validated.

For this purpose, model UFO, MG5 cards, and a pythia8 card for Monte Carlo production were provided by CMS to generate events with **MadGraph MG5_aMC**, showered with **Pythia 8**


This paper is written in the context of Z' -two-Higgs-doublet model, where a high-mass resonance Z' decays into a pseudoscalar boson A and a CP-even scalar Higgs boson, and the A decays to a pair of dark matter particles.

To generate signal sample, model UFO files is provided by CMS. From the CMS genproduction github repository one can retrieve the cards used for **MadGraph MG5_aMC** event generation for each mass point of Z' . The run card used in **MadGraph MG5_aMC** and proc card were retrieved from there. Also we applied some custom settings according to the mass of Z' . Some examples for a mass point $M_{Z_p} = 1000$ MeV is presented in appendix A. To see full information, please look at linked github pages.

Since **MadGraph MG5_aMC** cannot handle the decay of standard model higgs properly, higgs decay and parton shower was handled by **Pythia 8**. For example, specific **Pythia 8** card is used in this process. The pythia settings are then retrieved from the CMS software github repository:

- **Pythia8CUEP8M1Settings** and
- **Pythia8CommonSettings**. Also:
- The genfragment file is used.

Models studied are shown in 1. For further theoretical aspects of this model, see the paper arXiv:1402.7074

Figure 1: The Z' 2HDM model with pseudoscalar A

For detector simulation, we used **Delphes 3** with latest version of delphes card used for CMS EXO-16-037 recasting. The difference between custom card and default card is presented in appendix B. We added some lines to make neutralino not to deposit energy on calorimeter.

2 Cut flow

This analysis is a diphoton, missing transverse momentum and zero lepton analysis. Unfortunately we couldn't find out detailed cutflow. Here we present the product of acceptance and efficiency for signal in the SR for each mass point only. The error is defined as $(1 - (A \cdot \epsilon)^{MA5} / (A \cdot \epsilon)^{CMS})$ (%)

m_{Z_p} (MeV)	Acceptance \times efficiency ($A \cdot \epsilon$)		
	CMS EXO-16-012	MA5	Error
600	0.317 ± 0.004	0.317 ± 0.001	0 %
800	0.399 ± 0.004	0.410 ± 0.001	-2.8 %
1000	0.444 ± 0.004	0.441 ± 0.001	0.6 %
1200	0.474 ± 0.004	0.343 ± 0.001	27.6 %
1400	0.492 ± 0.004	0.221 ± 0.001	55.1 %
1700	0.493 ± 0.004	0.129 ± 0.0004	73.8 %
2000	0.351 ± 0.004	0.082 ± 0.0002	76.4 %
2500	0.213 ± 0.004	0.046 ± 0.0001	78.4 %

3 Distributions of observables

Since missing transverse energy is important, we overlayed

Appendices

A MadGraph MG5_aMC and Pythia 8 card settings

```

import model Zp2HDM
generate p p > Zp > h A0, A0 > n1 n1~
output Zprime_A0h_A0chichi_MZp1000_MA0300 -nojpeg

set param_card mass 32 1000
set param_card mass 26 300
set param_card mass 27 300
set param_card mass 28 300
set param_card ZpINPUTS 1 1
set param_card ZpINPUTS 2 0.8
set param_card DECAY 32 20.22504
set param_card DECAY 28 8.95228

####From the run card

'lhpdf' = pdlabel      ! PDF set
263400  = lhaid        ! if pdlabel=lhpdf, this is the lhpdf number

####In the shower card
!CUEP8M1 Tune
SLHA:minMassSM=1000.
SLHA:keepSM=on
SLHA:useDecayTable = off
Main:timesAllowErrors=10000
MultipartonInteractions:expPow=1.6
MultipartonInteractions:ecmPow=2.5208
MultipartonInteractions:pT0Ref=2.4024
ParticleDecays:limitTau0=on
ParticleDecays:allowPhotonRadiation=on
ParticleDecays:tau0Max=10
Check:epTolErr=1.0000000000e-02
Tune:ee=7
Tune:pp=14

!This is for decay mode
25:m0 = 125.0
25:onMode=off
25:OnIfMatch=22 22

```

B Delphes card settings

```
#####
# MC truth jet finder
#####

module FastJetFinder GenJetFinder {
  set InputArray NeutrinoFilter/filteredParticles

  set OutputArray jets

  # algorithm: 1 CDFJetClu, 2 MidPoint, 3 SIScone, 4 kt, 5 Cambridge/Aachen, 6
  antikt
  set JetAlgorithm 6
  set ParameterR 0.4

  set JetPTMin 20.0
}

#####
# Jet finder
#####

module FastJetFinder FastJetFinder {
# set InputArray Calorimeter/towers
  set InputArray EFlowMerger/eflow

  set OutputArray jets

  # algorithm: 1 CDFJetClu, 2 MidPoint, 3 SIScone, 4 kt, 5 Cambridge/Aachen, 6
  antikt
  set JetAlgorithm 6
  set ParameterR 0.4

  set JetPTMin 20.0
}

#####
# b-tagging
#####

module BTagging BTagging {
  set JetInputArray JetEnergyScale/jets

  set BitNumber 0

  # add EfficiencyFormula {abs(PDG code)} {efficiency formula as a function of
  eta and pt}
  # PDG code = the highest PDG code of a quark or gluon inside DeltaR cone
  around jet axis
  # gluon's PDG code has the lowest priority
}
```

```

add EfficiencyFormula {0} { (pt >= 30.0 && pt < 130.0) * (0.124 - 1.0*10^-3*pt
+ 1.06*10^-5*pt^2 - 3.18*10^-8*pt^3 + 3.13*10^-11*pt^4) +
                                (pt >= 130.0) * (0.055 + 4.53*10^-4*pt -
                                1.60*10^-7*pt^2) }

add EfficiencyFormula {4} { (pt >= 30.0 && pt < 205.0) * (0.40 + 1.23*10^-3*pt
- 4.60*10^-6*pt^2 + 5.71*10^-9*pt^3) +
                                (pt >= 205.0) * (0.478 + 1.573*10^-4*pt)}

add EfficiencyFormula {5} { (pt >= 30.0 && pt < 150.0) * (0.707 + 5.6*10^-3*pt
- 6.27*10^-5*pt^2 + 3.10*10^-7*pt^3 - 5.63*10^-10*pt^4) +
                                (pt >= 150.0) * (0.906 - 6.39*10^-5*pt +
                                4.11*10^-8*pt^2) }
}

#####
#   ECAL
#####

add EnergyFraction {18} {0.0}

#####
#   HCAL
#####

add EnergyFraction {18} {0.0}

```