SYNTHETIC COMMUNICATIONS® Vol. 34, No. 4, pp. 599–605, 2004

Magnesium-Catalyzed Proficient Reduction of Oximes to Amines Using Ammonium Formate

K. Abiraj and D. Channe Gowda*

Department of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore, Karnataka, India

ABSTRACT

Various aldoximes and ketoximes were selectively reduced to the corresponding amines by catalytic transfer hydrogenation employing low cost magnesium powder and ammonium formate at room temperature. Many other functionalities such as halogens, –OH, –OCH₃, –COOH and –CH₃ remained unaffected. The hydrogenation is fast, mild, clean, cost effective and high yielding.

Key Words: Oximes; Amines; Ammonium formate; Hydrogenation; Magnesium powder.

599

0039-7911 (Print); 1532-2432 (Online)

www.dekker.com

DOI: 10.1081/SCC-120027707Copyright © 2004 by Marcel Dekker, Inc.

^{*}Correspondence: D. Channe Gowda, Department of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore, Karnataka, India-570 006; Fax: 091-0821-2421263/2518835; E-mail: dcgowda@yahoo.com.

There has been growing interest in the use of metals in synthetic chemistry. Zinc, [11] indium, [12] tin, [13] magnesium [14–18] have been used in the synthesis of many organic compounds. The utility of magnesium for the synthesis of pinacols, [14] silaspiro compounds [15] has been demonstrated. Further, the magnesium-mediated reduction of aromatic nitro compounds, [16] Barbier reaction [17] and in situ Grignard reactions [18] has also been demonstrated. The application of ammonium formate [19,20] in the field of catalytic transfer hydrogenation for the reduction of variety of organic compounds and in the synthesis of peptides is peer reviewed.

Nowadays, the heterogeneous catalytic transfer hydrogenation method has proved to be a potent choice for reduction of organic compounds over traditional hydrogenation or other methods of reduction as it involves mild reaction condition, easy work-up and high degree of selectivity. [19–24] Earlier reports reveal that catalytic transfer hydrogenation of oximes to amines had been achieved with systems like ammonium formate/10% Pd/C, [23] cyclohexene/10% Pd/C. [24] But these systems require long reaction times as long as 5–10 hours at reflux temperature, expensive catalyst and also offer very low yield. Moreover, stringent precautions must be taken while employing palladium on carbon because of its flammable nature in presence of air.

Recently, we have explored the utility of magnesium powder and hydrazinium monoformate; a new hydrogen source for deblocking some commonly used protecting groups in peptide synthesis. [25] Herein we report a rapid, selective and simple reduction of oximes to the corresponding primary amines by using low cost magnesium powder and ammonium formate at room temperature as depicted in the Scheme 1. Various other functionalities like halogens, -OH, -OCH₃, -COOH and -CH₃ are tolerated.

Reduction of Oximes to Amines Using Ammonium Formate

$$R^1$$
 NOH $Mg/HCOONH_4$ R^1 NH₂ R^2 NH₂

Scheme 1. R^1 , $R^2 = H$ or alkyl or phenyl or substituted phenyl group.

The results given in the Table 1 reveal the viability of using Mg/HCOONH₄ system for the reduction of oximes. The course of reaction was monitored by TLC and IR spectra. The work-up and isolation of the products were easy. Thus, the oximes reduced (few examples are listed in the Table 1) by this system were obtained in good yield. The products were characterized by comparison of their boiling points or melting points, TLC and IR spectra with authentic samples. The disappearance of strong absorption bands between 1690–1640 cm⁻¹ due to C=N stretching and between 3650–3500 cm⁻¹ due to O-H stretching and appearance of two strong absorption bands between 3500–3300 cm⁻¹ of -NH₂ group clearly show that the oximes were reduced to corresponding primary amines.

A control experiment was carried out using oximes with ammonium formate, but without magnesium powder does not yield any reduced product and the starting material is recovered in 100%. This confirms the role of magnesium as catalyst. The reaction was also carried out for several hours with ammonium chloride instead of ammonium formate expecting an efficient reduction, but the yield is very low. Further, we observed that in the case of nitro oximes, the nitro group at aryl residue and also oxime group underwent reduction to yield respective diamine product at room temperature.

In summary, the reduction of oximes can be accomplished in a short time with magnesium powder instead of expensive catalyst like palladium^[23,24] at room temperature and many other functionalities are tolerated. The yields are almost quantitative and the compounds analytically pure. This magnesium-catalyzed procedure provides a very efficient, inexpensive, mild, and general methodology for reduction of oximes to amines.

EXPERIMENTAL

Materials

The oximes were either commercially available or prepared from the corresponding carbonyl compound by standard methods. In cases where the oxime was obtained as an E/Z-mixture, no attempts were made to separate such mixtures and they are used as such for the reduction. Magnesium powder



 $\it Table~1.$ Magnesium-catalyzed reduction of oximes to amines using ammonium formate.

$$\begin{array}{c|c}
R^1 & & & \\
\hline
 & NOH & & & \\
\hline
 & Mg/HCOONH_4 & & \\
\hline
 & MeOH, r. t. & & \\
\hline
 & R^2 & 2
\end{array}$$

					Boiling point		
1/2	R^1	R^2	Time (min)	Yield ^a (%)	Found	Literature	Ref.
a	Ph	Н	38	91	182-184	185	[26]
b	Me	Н	40	66 ^b	166 ^c	165	[26]
c	Me	Me	42	64 ^b	152 ^c	150	[26]
d	Ph	Me	40	89	184	184-186	[27]
e	Ph	Ph	45	93	295-296	295	[26]
f	p-(OH)C ₆ H ₄ CH ₂	Me	52	92	123°	125-126	[27]
g	p-(OCH ₃) ₃ C ₆ H ₄	Н	55	85	236	236-237	[26]
h	$3,4,5-(OCH_3)C_6H_2$	Н	52	84	122	121	[26]
i	p-(Cl)C ₆ H ₄	Н	32	94	212 - 214	215	[26]
j	d	d	50	81	157-159	158-160	[26]
k	e	e	55	79	135	134	[26]
l	f	f	52	88	100	101	[26]

^aIsolated yields are based on single experiment and the yields were not optimized.

^cMelting point.

was purchased from SISCO Research Laboratories Pvt. Ltd., Bombay (India) and was treated with 0.01 N hydrochloric acid for about 2 min. It was filtered through a sintered glass funnel and washed with water, dry methanol and dry ether. Thus obtained magnesium was vacuum dried and stored. Ammonium formate and 60–120-mesh silica gel (for column chromatography) were purchased from E. Merck (India) Ltd. All of the solvents used were analytical grade or were purified according to standard procedures. Thin layer chromatography was carried out on silica gel plates obtained from Whatman

^bLow yield is due to low boiling point of the product and are isolated as their hydrochloride salts.

Reduction of Oximes to Amines Using Ammonium Formate

Inc. The melting points were determined by a Thomas—Hoover melting point apparatus and are uncorrected. IR spectra were recorded on a Shimadzu FTIR-8300 spectrometer.

Typical Procedure

To a solution of the substrate (10 mmol) in methanol or in any other suitable solvent (20 mmol) was added ammonium formate (30 mmol) and magnesium powder (1 g, 0.041 mol). The mixture was stirred under nitrogen atmosphere at room temperature. The reaction was exothermic and effervescent. After the completion of reaction (monitored by TLC), the reaction mixture was filtered through celite. The organic layer is evaporated and the residue was dissolved in chloroform or dichloromethane or ether and washed with saturated sodium chloride solution to remove excess ammonium formate. The organic layer was dried over anhydrous sodium sulphate and evaporation of the organic layer followed by purification either by preparative TLC or by column chromatography to yield the desired product.

In order to obtain the volatile aliphatic amines in good yield, the reaction was carried out using a condenser cooled with ice water and the reaction flask immersed in a cold-water bath. After filtration, the reaction mixture was neutralized with HCl. The solvent was evaporated under reduced pressure. The residue was lyophilized or subjected to column chromatography by using 60–120 mesh silica gel and a suitable eluting system [50:50 chloroform:benzene (for entry 2e), 60:40 chloroform:benzene (for entries 2a & 2d), 80:20 chloroform:benzene (for entry 2j), 90:10 chloroform:benzene (for entry 2k), 80:20 chloroform:methanol (for entries 2b & 2c), 85:15 chloroform:methanol (for entries 2f & 2g), 90:10 chloroform:methanol (for entry 2i), 95:5 chloroform:methanol (for entries 2h & 2l)]. Aliphatic amines were obtained as their hydrochloride salts in up to 65% yield.

ACKNOWLEDGMENT

The authors wish to thank University Grants Commission, New Delhi, India for providing financial assistance.

REFERENCES

1. Carr, D.; Iddon, B.; Suschitzky, H. Synthesis and some reactions of 3,4,5,6-tetrahydro-3,6-diemthyl-2,6-methano-3-benzazocin-1(2H-one2,5-

- dimethyl-8-oxo-67-benzomorphan). J. Chem. Soc. Perkin Trans. 1. 1980,
- 2. Herscovici, J.; Egron, M.J.; Antonakis, K. Preparation of 4'-amino hexopyranose nucleosides an approach to amino nucleoside antibiotics. J. Chem. Soc. Perkin Trans. 1. 1988, 1219-1226.
- 3. Hutchins, R.O.; Hutchins, M.K. Reduction of C=N to CHNH by metal hydrides. In Comprehensive Organic Synthesis; Trost, B.M., Fleming, I., Eds.; Pergamon Press: Oxford, 1991; Vol. 8; 60-70.
- 4. Demir, A.S.; Tanyeli, C.; Sesenoglu, O.; Demic, S.A. Simple NaBH₄ in presence of transition metal compounds. Tetrahedron Lett. 1996, 37, 407 - 410.
- 5. Diab, Y.; Laurent, A.; Mison, P. Synthesis d'aziridines secondaires cis disubstituees. Tetrahedron Lett. 1974, 17, 1605-1607.
- 6. Rahman, M.A.; Fraser-Reid, B. Actinobolin via the anomeric effect. J. Am. Chem. Soc. 1985, 107, 5576-5578.
- 7. Pitts, M.R.; Harrison, J.R.; Moody, C.J. Indium metal as a reducing agent in organic synthesis. J. Chem. Soc. Perkin Trans. 1. 2001, 9, 955–977.
- 8. Cipolla, L.; Lay, L.; Nicotra, F.; Panza, L.; Russo, G. First synthesis of the phosphono analogue of N-acetyl- α -D-mannosamine 1-phosphate. J. Chem. Soc. Chem. Commun. **1995**, 1993–1994.
- 9. Booker-Milburn, K.I.; Dunkin, I.R.; Kelly, F.C.; Khalaf, A.I.; Learmonth, D.A.; Proctor, G.R.; Scopes, D.I.C. Azabenzocycloheptanones. Part 20. Synthesis and utilization of 4-amino-1,2,3,4-tetrahydro-1(1H)-benzazepines. J. Chem. Soc. Perkin Trans. 1. **1997**, 3261–3273.
- 10. Popp, F.D.; Schultz, H.P. Electrolytic reduction of organic compounds. Chem. Rev. **1962**, 62, 19–40.
- 11. Ranu, B.C.; Majee, A.; Das, A.R. A convenient synthesis of β, γ unsaturated ketones through zinc-mediated allylation of acid chlorides. Tetrahedron Lett. **1996**, *37* (7), 1109–1112.
- 12. Ranu, B.C.; Dutta, P.; Sarkar, A. Indium promoted reductive homocoupling of alkyl and aryl halides. Tetrahedron Lett. **1998**, 39 (57), 9557-9558.
- 13. Zhou, J.Y.; Chen, Z.G.; Wu, S.H. Tin promoted stereocontrolled intramolecular allylation of carbonyl compounds: a facile and stereoselective method for ring construction. J. Chem. Soc. Chem. Commun. **1994**, 24, 2783-2784.
- 14. House, H.O. Dissolving metal reductions and related reactions. In *Modern* Synthetic Reactions; Bartlett, P.D., Curtin, D., Johnson, W.S., Roberts, J.D., Woodward, R.B., Eds.; Benzamin, Inc.: USA, 1972; 145 - 227.
- 15. Salomon, R.G. A fecile one-step synthesis of 5-silaspiro [4,4]nona-2,7diene. J. Org. Chem. 1974, 39 (24), 3602.



Reduction of Oximes to Amines Using Ammonium Formate

- 16. Prajapathi, D.; Borah, H.N.; Sandhu, J.S.; Ghosh, A.C. Ammonium sulphate-magnesium promoted selective reduction of aromatic nitro compounds. Synth. Commun. **1995**, *25* (24), 4025–4028.
- 17. Blomberg, C.; Hartog, F.A. The barbier reaction—a one-step alternative for syntheses via organomagnesium compounds. Synthesis **1977**, *1*, 18–30.
- 18. Merker, R.L.; Scott, M.J. Grignard reactions of polyhalocarbons. In situ grignard reactions of carbon tetrahalides with organochlorosilanes. J. Org. Chem. **1964**, *29* (4), 953–955.
- 19. Johnstone, R.A.W.; Wilby, A.H.; Entwistle, I.D. Heterogeneous catalytic transfer hydrogenation and its relation to other methods for reduction of organic compounds. Chem. Rev. **1985**, 85, 129–170.
- 20. Ram, S.; Ehrenkaufer, R.E. Ammonium formate in organic synthesis: a versatile agent in catalytic transfer hydrogenation. Synthesis **1988**, *1*, 91–97.
- 21. Gowda, S.; Gowda, D.C. Application of hydrazinium monoformate as new hydrogen donor with raney nickel; a facile reduction of nitro and nitrile moieties. Tetrahedron **2002**, *58* (11), 2211–2213.
- 22. Gowda, D.C.; Gowda, A.S.P.; Baba, A.R.; Gowda, S. Nickel-catalyzed formic acid reductions, a selective method for the reduction of nitro compounds. Synth. Commun. **2000**, *30* (16), 2885–2889.
- 23. Jnaneshwara, G.K.; Sudalai, A.; Deshpande, V.H. Palladium catalyzed transfer hydrogenation of azo compounds and oximes using ammonium formate. J. Chem. Res. (S). **1998**, *3*, 160–161.
- 24. Brieger, G.; Nestrick, T.J. Catalytic transfer hydrogenation. Chem. Rev. **1974**, *74*, 567–580.
- 25. Gowda, D.C. Magnesium/hydrazinium monoformate: a new hydrogenation method for the removal of some commonly used protecting groups in peptide synthesis. Tetrahedron Lett. **2002**, *43*, 311–313.
- Vogel, A.I. Text Book of Practical Organic Chemistry, 5th Ed.; Revised by Furniss, B.S., Hannaford, A.J., Smith, P.W.G., Tatchell, A.R., Eds.; Addison Wesley Longman Limited: UK, 1997; 1298.
- 27. Budavari, S, Ed.; *The Merck Index*, 11th Ed.; Merck & Co. Inc.: USA, 1989.

Received in the USA August 14, 2003

Copyright of Synthetic Communications is the property of Marcel Dekker Inc. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.

Copyright of Synthetic Communications is the property of Taylor & Francis Ltd and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.