

---

Section 3.1:

1. Refer to the recurrence relation for the Fibonacci sequence in Definition 3.1

- (a) Answer Fibonacci's question by calculating  $F(12)$ .

$$F(12) = 144$$

- (b) Write  $F(1000)$  in terms of  $F(999)$  and  $F(998)$ .

$$F(1000) = F(999) + F(998)$$

- (c) Write  $F(1000)$  in terms of  $F(998)$  and  $F(997)$ .

$$F(1000) = 2F(998) + F(997)$$

2. In Fibonacci's model, rabbits live forever. The following modification of Definition 3.1 accounts for dying rabbits:

$$G(n) = \begin{cases} 0 & \text{if } n \leq 0, \\ 1 & \text{if } n = 1 \text{ or } n = 2, \\ G(n-1) + G(n-2) - G(n-8) & \text{if } n > 2. \end{cases}$$

- (a) Compute  $G(n)$  for  $n = 1, 2, \dots, 12$ :

$$G(1) = 1$$

$$G(2) = 1$$

$$G(3) = 2$$

$$G(4) = 3$$

$$G(5) = 5$$

$$G(6) = 8$$

$$G(7) = 13$$

$$G(8) = 21$$

$$G(9) = 33$$

$$G(10) = 53$$

$$G(11) = 86$$

$$G(12) = 136$$

- (b) In this modified model, how long do rabbits live?  
The rabbits live for 8 months.

3. Consider the following recurrence relation:

$$H(n) = \begin{cases} 0 & \text{if } n \leq 0, \\ 1 & \text{if } n = 1 \text{ or } n = 2, \\ H(n-1) + H(n-2) - H(n-3) & \text{if } n > 2. \end{cases}$$

(a) Compute  $H(n)$  for  $n = 1, 2, \dots, 10$

$$H(1) = 1$$

$$H(2) = 1$$

$$H(3) = 2$$

$$H(4) = 2$$

$$H(5) = 3$$

$$H(6) = 3$$

$$H(7) = 4$$

$$H(8) = 4$$

$$H(9) = 5$$

$$H(10) = 5$$

(b) Using the pattern from part(a), guess what  $H(100)$  is

$$H(100) = 50$$

4. The Lucas numbers  $L(n)$  have almost the same definition as the Fibonacci numbers:

$$L(n) = \begin{cases} 1 & \text{if } n = 1, \\ 3 & \text{if } n = 2 \\ L(n-1) + L(n-2) & \text{if } n > 2. \end{cases}$$

(a) How is the definition of  $L(n)$  different from the definition of  $F(n)$  in Definition 3.1?

The definition of  $L(n)$  is different from  $F(n)$  when  $n = 2$ .

$L(2) = 3$  and  $F(2) = 1$ .

(b) Compute the first 12 Lucas numbers.

$$L(1) = 1$$

$$L(2) = 3$$

$$L(3) = 4$$

$$L(4) = 7$$

$$L(5) = 11$$

$$L(6) = 18$$

$$L(7) = 29$$

$$L(8) = 47$$

$$L(9) = 76$$

$$L(10) = 123$$

$$L(11) = 199$$

$$L(12) = 322$$

5. Compute the first seven terms of the following recurrence relations:

(a)  $C(n)$  of Example 3.5

$$C(n) = \begin{cases} 1 & \text{if } n = 0, \\ 2.C(n-1) & \text{if } n > 0 \end{cases}$$

$$C(0) = 1$$

$$C(1) = 2$$

$$C(2) = 4$$

$$C(3) = 8$$

$$C(4) = 16$$

$$C(5) = 32$$

$$C(6) = 64$$

(b)  $V(n)$  of Example 3.6

$$V(n) = \begin{cases} 1 & \text{if } n = 0, \\ 2.V(n-1) + 1 & \text{if } n > 0 \end{cases}$$

$$V(0) = 1$$

$$V(1) = 3$$

$$V(2) = 7$$

$$V(3) = 15$$

$$V(4) = 31$$

$$V(5) = 63$$

$$V(6) = 127$$

(c)  $E(n)$  of Example 3.7

$$E(n) = \begin{cases} 0 & \text{if } n = 1, \\ E(n-1) + n - 1 & \text{if } n > 1 \end{cases}$$

$$E(1) = 0$$

$$E(2) = 1$$

$$E(3) = 3$$

$$E(4) = 6$$

$$E(5) = 10$$

$$E(6) = 15$$

$$E(7) = 21$$

6. Consider the following recurrence relation.

$$P(n) = \begin{cases} 0 & \text{if } n = 0 \\ [P(n-1)]^2 - n & \text{if } n > 0 \end{cases}$$

Use this recurrence relation to compute  $P(1)$ ,  $P(2)$ ,  $P(3)$ , and  $P(4)$

$$P(1) = -1$$

$$P(2) = -1$$

$$P(3) = -2$$

$$P(4) = 0$$

7. Given the following definition, compute  $P(4)$ .

$$Q(n) = \begin{cases} 1 & \text{if } n = 0 \\ n \times [P(n-1)] & \text{if } n > 0 \end{cases}$$

$$P(4) = 4 \times P(3) = 4 \times 3 \times P(2) = 4 \times 3 \times 2 \times P(1) = 4 \times 3 \times 2 \times 1 \times P(0)$$

$$4 \times 3 \times 2 \times 1 \times 1 = 24$$

8. Given the following definition, compute  $Q(5)$ .

$$Q(n) = \begin{cases} 0 & \text{if } n = 0 \\ 1 & \text{if } n = 1 \\ 2 & \text{if } n = 2 \\ Q(n-1) + Q(n-2) + Q(n-3) & \text{if } n > 2 \end{cases}$$

$$Q(5) = Q(4) + Q(3) + Q(2) = Q(3) + Q(2) + Q(1) + Q(2) + Q(1) + Q(0) + Q(2)$$

$$= Q(2) + Q(1) + Q(0) + Q(2) + Q(1) + Q(2) + Q(1) + Q(0) + Q(2)$$

$$= 2 + 1 + 0 + 2 + 1 + 2 + 1 + 0 + 2 = 11$$

9. Consider the recurrence relation defined in Example 3.3. Suppose that, as in the example, you borrow \$500, but you pay her back \$100 each week. Each week, Ursula charges you 10% interest on the amount you still owe, after your \$100 payment is taken into account.

- (a) Write down a recurrence relation for  $M(n)$ , the amount owed after  $n$  weeks.

$$M(n) = \begin{cases} 500 & \text{if } n = 0 \\ 1.1 \times (M(n-1) - 100) & \text{if } n > 0 \end{cases}$$

- (b) How much will you owe after four weeks?

$$M(4) = 1.1 \times (M(3) - 100) = 1.1 \times (1.1 \times (M(2) - 100) - 100)$$

$$= 1.1 \times (1.1 \times (1.1 \times (M(1) - 100) - 100) - 100)$$

$$= 1.1 \times (1.1 \times (1.1 \times (1.1 \times (M(0) - 100) - 100) - 100) - 100)$$

$$= 1.1 \times (1.1 \times (1.1 \times (1.1 \times (500 - 100) - 100) - 100) - 100) = 221.54$$

10. Every year, Alice gets a raise of \$3,000 plus 5% of her previous year's salary. Her starting salary is \$50,000. Give a recurrence relation for  $S(n)$ , Alice's salary after  $n$  years, for  $n \geq 0$ .

$$S(n) = \begin{cases} 50000 & \text{if } n = 0 \\ 1.05 \times S(n-1) + 3000 & \text{if } n > 0 \end{cases}$$

11. Suppose that today (year 0) your car is worth \$10,000. Each year your car loses 10% of its value, but at the end of each year you add customizations to your car that increase its value by \$50. Write a recurrence relation to model this situation. Let  $P(n)$  is the value of the car after  $n$  year(s).

$$P(n) = \begin{cases} 10000 & \text{if } n = 0 \\ 0.9 \times P(n-1) + 50 & \text{if } n > 0 \end{cases}$$

Section 3.4:

13. In Exercise 3 of Section 3.1, we defined the following recurrence relation:

$$H(n) = \begin{cases} 0 & \text{if } n \leq 0 \\ 1 & \text{if } n = 1 \text{ or } n = 2 \\ H(n-1) + H(n-2) - H(n-3) & \text{if } n > 2 \end{cases}$$

Prove that  $H(2n) = H(2n-1) = n$  for all  $n \geq 1$ .

**Step 1:**

For  $n = 1$ :

$$H(2 \times 1) = H(2) = 1$$

$$H(2 \times 1 - 1) = H(1) = 1$$

Hence,  $H(2 \times 1) = H(2 \times 1 - 1) = 1$ .

Then  $H(2n) = H(2n-1) = n$  is true for  $n = 1$ .

For  $n = 2$ :

$$H(2 \times 2) = H(4) = H(3) + H(2) - H(1) = H(2) + 1 - 1 = 1 + 1 - 0 = 2$$

$$H(2 \times 2 - 1) = H(3) = 1 + 1 - 0 = 2$$

Hence,  $H(2 \times 2) = H(2 \times 2 - 1) = 2$ .

**Step 2:**

For  $n = k$ , assume  $H(2k) = H(2k-1) = k$  is true.

**Step 3:**

For  $n = k + 1$ :

$$H(2(k+1)) = H(2(k+1) - 1)$$

$$H(2k+2) = H(2k+2-1)$$

$$H(2k+2) = H(2k+1)$$

$$H(2k+1) + H(2k) - H(2k-1) = H(2k) + H(2k-1) - H(2k-2)$$

$$H(2k+1) + k - k = k + k - H(2k-2)$$

$$H(2k) + H(2k-1) - H(2k-2) = 2k - H(2k-2)$$

$$k + k - H(2k-2) = 2k - H(2k-2)$$

$$0 = 0(true)$$

Hence,  $H(2(k+1)) = H(2(k+1)-1)$  is true.

$$H(2(k+1)) = k+1$$

$$H(2k+2) = k+1$$

$$H(2k+1) + H(2k) - H(2k-1) = k+1$$

$$H(2k+1) + k - k = k+1$$

$$H(2k+1) = k+1$$