

Name:.....

Class:.....



Mathematics for Engineering

Exercise Book

Trần Trọng Huỳnh - 2018

CALCULUS

Chapter 1: Function and Limit

1. Find the domain of each function:

a. $f(x) = \frac{2}{(x+2)\sqrt{x+1}}$

b. $f(x) = \frac{2x-1}{\sqrt{x|x-4|}}$

c. $f(x) = \ln(x+1) - \frac{x}{\sqrt{x-1}}$

d. $f(x) = \frac{\sqrt{x^2-4x+3}}{\lg(x-2)}$

2. Find the range of each function:

a. $f(x) = \frac{3x+5}{2x-1}$

b. $f(x) = x^2 - 2x$

c. $f(x) = \frac{x^2-x+1}{x^2+x+1}$

3. Determine whether is even, odd, or neither

a) $f(x) = \frac{|x-1|+|x+1|}{x^3}$

c) $f(x) = \ln(x + \sqrt{1+x^2})$

b) $f(x) = \frac{1}{2}(a^x + a^{-x})$

d) $f(x) = \lg \frac{1+x}{1-x}$

4. Explain how the following graphs are obtained from the graph of $f(x)$

a. $f(x-4)$

b. $f(x)+3$

c. $f(x-2)-3$

d. $f(x+5)-4$

5. Suppose that the graph of $f(x) = \sqrt{x}$ is given. Describe how the graph of the

function $y = \sqrt{x-1} + 2 = f(x-1) + 2$ can be obtained from the graph of f .

6. Let $f(x) = \sqrt{x}$ and $g(x) = \sqrt{2-x}$. Find each function

a. $f \circ g$

b. $g \circ f$

c. $g \circ g$

d. $f \circ f$

7. Let $f(x) = \frac{x^2+x+1}{x} = x+1+\frac{1}{x}$. Find

a. $f\left(x+\frac{1}{x}\right)$

b. $f(2x-1)$

8. Use the table to evaluate each expression

- a. $f(g(1))$ b. $g(f(1))$ c. $f(f(1))$ d. $g(g(1))$
e. $(g \circ f)(3)$ f. $(g \circ f)(6)$


x	1	2	3	4	5	6
$f(x)$	3	1	4	2	2	5
$g(x)$	6	3	2	1	2	3

9. Evaluate the following limits

- a. $\lim_{x \rightarrow 3} \frac{x^2 + x - 12}{x - 3}$ b. $\lim_{x \rightarrow 1} \frac{x^6 - 1}{x^{10} - 1}$ c. $\lim_{x \rightarrow 0} \frac{\tan 3x + 2x}{\tan 5x - \sin x}$ d. $\lim_{h \rightarrow 0} \frac{(3+h)^2 - 9}{h}$
e. $\lim_{t \rightarrow 0} \frac{\sqrt{t^2 + 9} - 3}{t^2}$ f. $\lim_{x \rightarrow \infty} \frac{x^2 + x - 12}{x^3 - 3}$ g. $\lim_{x \rightarrow 0^-} \left(\frac{1}{x} - \frac{1}{|x|} \right)$ h. $\lim_{x \rightarrow 1} \frac{x^2 - 1}{|x - 1|}$

a-e ; b-g ; c-f ; d-h

10. Determine whether each curve is the graph of a function of x . If it is, state the domain and range of the function.




11. The graph of f is given.

a. Find each limit, or explain why it does not exist.

- i. $\lim_{x \rightarrow 0^+} f(x)$, $\lim_{x \rightarrow 0^-} f(x)$ and $\lim_{x \rightarrow 0} f(x)$
ii. $\lim_{x \rightarrow 1} f(x)$ and $\lim_{x \rightarrow 4} f(x)$

b. At what numbers is discontinuous?



12. Determine where the function $f(x)$ is continuous

a. $f(x) = \frac{2x^2 + x - 1}{x - 2}$ b. $f(x) = \frac{x - 9}{\sqrt{4x^2 + 4x + 1}}$ c. $f(x) = \ln(2x + 5)$

13. Find the constant m that makes f continuous on its domain

a. $f(x) = \begin{cases} x^2 - m^2, & x < 4 \\ mx + 20, & x \geq 4 \end{cases}$ b. $f(x) = \begin{cases} mx^2 + 2x, & x < 2 \\ x^3 - mx, & x \geq 2 \end{cases}$

c. $f(x) = \begin{cases} \frac{e^{2x} - 1}{x}, & x \neq 0 \\ m, & x = 0 \end{cases}$ d. $f(x) = \begin{cases} \frac{x^2 - 1}{x - 1}, & x \neq 1 \\ m + 1, & x = 1 \end{cases}$

14. Find the numbers at which the function $f(x) = \begin{cases} x + 2, & x < 0 \\ 2x^2, & 1 \geq x \geq 0 \\ 2 - x, & x > 1 \end{cases}$ is discontinuous.

Chapter 2: Derivatives

1. Find an equation of the **tangent line** to the curve at the given point:

a. $y = \frac{x-1}{x-2}, \quad (3, 2)$

b. $y = \frac{2x}{x^2+1}, \quad (0, 0)$

c. $y = 3 - 2x + x^2, \quad x = 1$

d. $y = \frac{3-2x}{x-1}, \quad y = -1$

2. Find y'

a. $y = x^2 - x\sqrt{x} + \frac{1}{x} + 2$

b. $y = \sqrt{x + \sqrt{x}}$

c. $y = \frac{x^2}{x+1}$

d. $y = x\sqrt{x+2}$

e. $y = \ln(x^2 + 1) - \frac{1}{x}$

f. $y = e^x \sin(2x + 1)$

3. Find y''

a. $y = xe^{3x-1}$

b. $y = \sqrt[3]{2x+1}$

c. $y = e^{-x} \cos x$

4. Find dy/dt for:

a. $y = x^3 + x + 2, dx/dt = 2$ and $x = 1$

b. $y = \ln x, dx/dt = 1$ and $x = e^2$

c. $y = \tan \sqrt{t}$ and $t = \frac{\pi^2}{16}$

d. $\begin{cases} y = \sin \varphi \\ t = \cos \varphi \end{cases}$ and $\varphi = \frac{\pi}{3}$

5. Find dy for: $dy = y'(x)dx$


a. $y = \frac{1}{x^2 + 1}$

b. $y = \sqrt{x+1}, x = 3$


c. $y = \ln(x^2 + 1), x = 1$ and $dx = 0.1$

6. The graph of is given. State the numbers at which is not differentiable

a.



b.



7. A table of values for f, f', g and g' is given

x	$f(x)$	$g(x)$	$f'(x)$	$g'(x)$
1	3	2	4	6
2	1	8	5	7
3	7	2	7	9

- a. If $h(x) = f(g(x))$, find $h'(1)$ b. If $H(x) = (g \circ f)(x) = g(f(x))$, find $H'(1)$
c. If $F(x) = (f \circ f)(x)$, find $F'(2)$ d. If $G(x) = g \circ g(x)$, find $G'(3)$
8. If $h(x) = \sqrt{4 + 3f(x)}$, where $f(1) = 7, f'(1) = 4$, find $h'(1)$.

9. For the circle $x^2 + y^2 = 25$.

a. Find dy/dx

b. Find an equation of the tangent to the circle at the point (3, 4).

10. Let $(L): x^3 + y^3 = 6xy$

a. Find dy/dx

b. Find an equation of tangent to the curve (L) at the point (3, 3)

11. Find y' by implicit differentiation

a. $x^4 + y^4 = 16x + y$

b. $\sqrt{x} + \sqrt{y} = 4$

c. $x^3 + xy = y^2$

12. Find f' in terms of g'

a. $f(x) = g(\sin 2x)$

b. $f(x) = g(e^{1-3x})$

13. Each side of a square is increasing at a rate of 6 cm/s. At what rate is the area of the square increasing when the area of the square is 16 cm^2 ?

14. If $x^2 + y^2 = 25$ and $dy/dt = 6$, find dx/dt when $y = 4$ and $x > 0$.

15. If $z^2 = x^2 + y^2$ ($z > 0$), $dx/dt = 2, dy/dt = 3$, find dz/dt when $x = 5, y = 12$

16. Find the linearization $L(x)$ of the function at a.

a. $f(x) = \frac{1}{\sqrt{2+x}}$, $a = 2$

b. $f(x) = \sqrt[3]{5-x}$, $a = -3$

17. The equation of motion is $s(t) = 3\sin t - 4\cos t + 1$ for a particle, where s is in meters and t is in seconds. Find the **acceleration** (in m/s^2) after 3 seconds.

Chapter 3: Applications of Differentiation

1. Find the absolute maximum and absolute minimum values of the function on the given interval

a. $f(x) = 3x^2 - 12x + 5, [0; 3]$

b. $f(x) = x^3 - 3x + 5, [0; 3]$

c. $f(x) = x\sqrt{4 - x^2}, [-1; 2]$

d. $f(x) = x - \ln x, \left[\frac{1}{2}; 2\right]$

2. Find the critical numbers of the function

a. $f(x) = 5x^2 + 4x$

b. $f(x) = \frac{x-1}{x^2 - x + 1}$

c. $f(x) = x \ln x$

3. Find all numbers that satisfy the conclusion of the **Rolle's Theorem**

a. $f(x) = x\sqrt{x+2}, [-2; 0]$

b. $f(x) = (x-2)x^2, [0; 2]$

4. Find all numbers that satisfy the conclusion of the **Mean Value Theorem**

a. $f(x) = 3x^2 + 2x + 5, [-1; 1]$

b. $f(x) = e^{-2x}, [0; 3]$

5. If $f(1) = 10$ and $f'(x) \geq 2, \forall x \in [1; 4]$, how small can $f(4)$ possibly be?

6. Find where the function $f(x) = 3x^4 - 4x^3 - 12x^2 + 1$ is increasing and where it is decreasing.

7. Find the **inflection points** for the function

a. $f(x) = x^4 - 4x + 1$

b. $f(x) = x^6$

c. $f(x) = xe^x$

8. Find $f(x)$ for $f'(x) = \sqrt{2x+1}$ and $f(0) = 1$.

9. Find the point on the parabola $y^2 = 2x$ that is closest to the point $(1; 4)$

10. Find two numbers whose difference is 100 and whose product is a minimum.

11. Find two positive numbers whose product is 100 and whose sum is a minimum.

12. Use Newton's method with the specified initial approximation x_1 to find x_3

a. $x^3 + 2x - 4 = 0, x_1 = 1$

b. $x^5 + 2 = 0, x_1 = -1$

c. $\ln(x^2 + 1) - 2x - 1 = 0, x_1 = 1$

d. $\ln(4 - x^2) = x, x_1 = 1$

13. Find the most general anti-derivative of the function.

a. $f(x) = 6x^2 - 2x + 3$ b. $f(x) = \sqrt[6]{x} + \frac{1}{x^2}$

c. $f(x) = \frac{x^2 + x + 2}{x}$ d. $f(x) = 2x(x^2 + 1)$

14. Find the anti-derivative of that satisfies the given condition

a. $f(x) = 5x^4 - 2x^5, F(0) = 4$ b. $f(x) = 4 - \frac{2x}{x^2 + 1}, F(0) = 1$


15. A particle is moving with the given data. Find the position of the particle

a. $v(t) = \sin t - \cos t, s(0) = 0$

b. $v(t) = 10\sin t + 3\cos t, s(\pi) = 0$

c. $v(t) = 10 + 3t - 3t^2, s(2) = 10$

16. The figure shows the graph of the derivative f' of a function f



a. On what intervals is f increasing or decreasing?

b. For what values of x does f have a local maximum or minimum?

Chapter 4 - 6: Integration

1. Estimate the area under the graph of $y = f(x)$ using 6 rectangles and left endpoints

a. $f(x) = \frac{1}{x} + x$, $x \in [1, 4]$

b. $f(x) = x^2 - 2$, $x \in [-1, 2]$

c. A table of values for f is given

x	1	2	3	4	5	6	7
$f(x)$	5	6	3	2	7	1	2

3. Repeat part (1) using right endpoints

4. For the function $f(x) = x^3$, $x \in [-2, 2]$. Estimate the area under the graph of using four approximating rectangles and taking the sample points to be

a. Right endpoints

b. Left endpoints

c. Midpoints

5. Use (a) the Trapezoidal Rule, (b) the Midpoint Rule, and (c) Simpson's Rule to approximate the given integral with the specified value of n .

a. $\int_0^3 \sqrt{x} dx$, $n = 4$

b. $\int_1^3 \frac{\sin x}{x} dx$, $n = 6$

6. Let $I = \int_0^2 \frac{dx}{x^2 + 1}$. Find the approximations L_4 , R_4 , M_4 , T_4 and S_4 for I .

7. Find the derivative of the function $g(x) = \int_0^x \sqrt{t^2 + 1} dt$

8. Find g'

a. $g(x) = \int_1^{x^4} \frac{1}{\cos t} dt$

b. $g(x) = \int_1^{\sqrt{x}} \frac{\sin u}{u} du$

c. $g(x) = \int_{2x}^{x^2+x+2} \frac{e^t}{t} dt$

d. $g(x) = \int_{\sin x}^{\cos x} (1+v^2)^{10} dv$

9. Find the average value of the function on the given interval

a. $f(x) = x^2, \quad [-1, 1]$

b. $f(x) = \frac{1}{x}, \quad [1, 5]$

c. $f(x) = x\sqrt{x}, \quad [1, 4]$

d. $f(x) = x \ln x, \quad [1, e^2]$

10. A particle moves along a line so that its velocity at time t is $v(t) = t^2 - t - 6$ (m/s)

a. Find the displacement of the particle during the time period $1 \leq t \leq 4$

b. Find the distance traveled during this time period

11. Suppose the acceleration function and initial velocity are $a(t) = t + 3$ (m/s²), $v(0) = 5$ (m/s). Find the velocity at time t and the distance traveled when $0 \leq t \leq 5$.

12. A particle moves along a line with velocity function $v(t) = t^2 - t$, where t is measured in meters per second. Find the **displacement** and the **distance traveled** by the particle during the time interval $t \in [0, 2]$.

13. Evaluate the integral

a. $\int_0^2 x^2 \cdot \sqrt{x^3 + 1} \, dx$

b. $\int x e^{x^2} \, dx$

c. $\int \left(\frac{1}{x} + \sqrt{x} - 3x^2 \right) dx$

d. $\int_0^1 y(1 + y^2)^5 \, dy$

e. $\int \frac{\ln x}{x} \, dx$

f. $\int \frac{t}{t^2 + 1} \, dt$

a-b, d-f

14. Evaluate the integral

a. $\int x e^x \, dx$

b. $\int_0^1 x^2 e^{-x} \, dx$

c. $\int x \sin x \, dx$

d. $\int \ln x \, dx$

e. $\int_1^e x \ln x \, dx$

f. $\int e^{\sqrt{x}} \, dx$

14b-13c, 14f-13e

15. Suppose $f(x)$ is differentiable, $f(1) = 4$ and $\int_0^1 f(x) \, dx = 5$. Find $\int_0^1 x f'(x) \, dx$

16. Suppose $f(x)$ is differentiable, $f(1) = 3$, $f(3) = 1$ and $\int_1^3 x f'(x) \, dx = 13$. What is the

average value of f on the interval $[1, 3]$?

17. Let $f(x) = \begin{cases} -x-1, & -3 \leq x \leq 0 \\ -\sqrt{1-x^2}, & 0 \leq x \leq 1 \end{cases}$. Evaluate $\int_{-3}^1 f(x) dx$

18. Find $g'(0)$ for

a. $g(x) = \int_x^{x^2} e^{2t+1} dt$ b. $\int_{2x-1}^{x^3} t\sqrt{t+1} dt$

19. Determine whether each integral is convergent or divergent. Evaluate those that are convergent.

a. $\int_1^{\infty} \frac{dx}{(3x+1)^2}$ b. $\int_{-\infty}^0 \frac{dx}{2x-5}$ c. $\int_{-\infty}^{-1} \frac{dx}{\sqrt{2-x}}$ d. $\int_0^{\infty} \frac{xdx}{(x^2+2)^2}$
e. $\int_4^{\infty} e^{-\frac{y}{2}} dy$ f. $\int_{-\infty}^{-1} e^{-2t} dt$ g. $\int_{-\infty}^{\infty} xe^{-x^2} dx$

i. $\int_0^1 \frac{dx}{4x-1}$ j. $\int_3^4 \frac{dx}{\sqrt{x-3}}$ k. $\int_{-1}^1 \frac{1}{\sqrt[3]{x^2}} dx$ l. $\int_0^1 \frac{dx}{\sqrt{x}}$

20. Use the Comparison Theorem to determine whether the integral is convergent or divergent

a. $\int_1^{\infty} \frac{\cos^2 x dx}{1+x^2}$ b. $\int_1^{\infty} \frac{2+e^{-x}}{x} dx$ c. $\int_1^{\infty} \frac{dx}{x+e^{2x}}$
d. $\int_1^{\infty} \frac{xdx}{\sqrt{1+x^6}}$ e. $I = \int_0^{\frac{\pi}{2}} \frac{\cos x dx}{\sqrt{\sin x}}$ f. $\int_0^1 \frac{2dx}{\sqrt{x^3}}$

LINEAR ALGEBRA

Chapter 1: Systems of Linear Equations

1. Write the **augmented matrix** for each of the following systems of linear equations and then solve them.

$$\text{a. } \begin{cases} -x + y + 2z = 1 \\ 2x + 3y + z = -2 \\ 5x + 4y + 2z = 4 \end{cases}$$

$$\text{b. } \begin{cases} 2x + 3y + z = 10 \\ 2x - 3y - 3z = 22 \\ 4x - 2y + 3z = -2 \end{cases}$$

$$\text{c. } \begin{cases} x + y + z = 0 \\ 2x - y + 2z = 0 \\ x + z = 0 \end{cases}$$

$$\text{d. } \begin{cases} x_1 + 2x_2 - x_3 + x_4 = 0 \\ 2x_1 + 3x_2 - 2x_3 + 3x_4 = 0 \\ x_1 + x_2 - 3x_3 + x_4 = 0 \end{cases}$$

2. Compute the rank of each of the following matrices.

$$\text{a. } A = \begin{pmatrix} 1 & 1 & 2 \\ 3 & -1 & 1 \\ -1 & 3 & 4 \end{pmatrix}$$

$$\text{b. } B = \begin{pmatrix} -2 & 3 & 3 \\ 3 & -4 & 1 \\ -5 & 7 & 2 \end{pmatrix}$$

$$\text{c. } C = \begin{pmatrix} 1 & 1 & -1 & 4 \\ 2 & 1 & 3 & 0 \\ 0 & 1 & 5 & 8 \end{pmatrix}$$

$$\text{d. } D = \begin{pmatrix} 1 & 1 & -1 & 3 \\ -1 & 4 & 5 & -2 \\ 1 & 6 & 3 & 4 \end{pmatrix}$$

11. Determine the values of m such that the rank of the matrix is 2

$$\text{a. } \begin{pmatrix} 1 & -1 & 0 \\ 2 & 3 & 5 \\ 1 & 2 & m \end{pmatrix}$$

$$\text{b. } \begin{pmatrix} 1 & 2 & 1 & 4 \\ 2 & 1 & 1 & 5 \\ -3 & 6 & 1 & m \end{pmatrix}$$

$$\text{c. } \begin{pmatrix} 1 & 2 & 3 \\ 2 & -1 & -1 \\ 3 & 1 & 2 \\ m & 3 & 5 \end{pmatrix}$$

$2b - 11b, 2d - 11c$

3. Find all values of k for which the system has **nontrivial solutions** and determine all solutions in each case.

$$\text{a. } \begin{cases} x - y + 2z = 0 \\ -x + y - z = 0 \\ x + ky + z = 0 \end{cases} \quad (1)$$

$$\text{b. } \begin{cases} x - 2y + z = 0 \\ x + ky - 3z = 0 \\ x - 6y + 5z = 0 \end{cases}$$

$$\text{c. } \begin{cases} x + y + z = 0 \\ x + y - z = 0 \\ x + y + kz = 0 \end{cases}$$

$$\text{d. } \begin{cases} x + y - z = 0 \\ ky - z = 0 \\ x + y + kz = 0 \end{cases}$$

4. Determine the values of m such that the system of linear equations has **exactly one solution**.

$$\text{a. } \begin{cases} x - y + 2z = m \\ -x + y - z = 0 \\ -x + my - z = 1 - m \end{cases} \quad (1)$$

$$\text{b. } \begin{cases} mx + y + z = 1 \\ x + my + z = m \\ x + y + mz = m^2 \end{cases}$$

$$\text{c. } \begin{cases} x + y - z = 1 \\ x + my + 2z = m \\ x + 2y + z = 2 \end{cases}$$

$$\text{d. } \begin{cases} x + my - mz = m \\ 2x + y - z = 2 \\ x + y + z = 0 \end{cases}$$

5. Determine the values of m such that the system of linear equations is **inconsistent**.

$$\text{a. } \begin{cases} x - y + 2z = m \\ -x + y - z = 0 \\ x - y + 3z = 1 - m \end{cases}$$

$$\text{b. } \begin{cases} x - 2y + 2z = m \\ x + my + z = 0 \\ 2x + y + mz = 2 - m \end{cases} \quad (1)$$

6. Find a , b and c so that the system $\begin{cases} x + ay + cz = 0 \\ bx + cy - 3z = 1 \\ ax + 2y + bz = 5 \end{cases}$ has the solution $(3, -1, 2)$

7. Consider the matrix $A = \left(\begin{array}{cc|c} 2 & -1 & 3 \\ -4 & 2 & k \\ 4 & -2 & 6 \end{array} \right)$

a. If A is the **augmented matrix** of a system of linear equations, determine the number of equations and the number of variables.

b. If A is the augmented matrix of a **system of linear equations**, find the value(s) of k such that the system is **consistent**.

8. Find all values of k so that the system of equations has **no solution**.

$$\text{a. } \begin{cases} x + y - z = 2 \\ -2y + z = 3 \\ 4y - 2z = k \end{cases}$$

$$\text{b. } \begin{cases} x + y - z = 1 \\ 2x + (k + 5)y - 2z = 4 \\ x + (k + 3)y + (k - 1)z = k + 3 \end{cases}$$

9. Find all values of a and b for which the system of equations
$$\begin{cases} x + y + 3z = 2 \\ x + 2y + 5z = 1 \\ 2x + 2y + az = b \end{cases}$$

is **inconsistent**.

10. Solve the system of linear equation corresponding to the given **augmented matrix**

a. $A = \begin{pmatrix} 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 \end{pmatrix}$

b. $B = \begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 2 & 4 \\ 0 & 0 & 0 & 0 \end{pmatrix}$

Chapter 2: Matrix Algebra

1. Let $A = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$, $B = \begin{pmatrix} -1 & 1 \\ 3 & 2 \end{pmatrix}$ and $C = \begin{pmatrix} 1 & 3 & -4 \\ -1 & 2 & 1 \end{pmatrix}$. Compute the matrix

- a. $2A - B^T$ b. AB c. BA d. AC
 e. CC^T f. C^TC g. A^3 h. B^2A^T

2. Suppose that A and B are $n \times n$ matrices. Simplify the expression

- a. $(A+B)^2 - (A-B)^2$
 b. $A(BC-CD) + A(C-B)D - AB(C-D)$

3. Let $A = \begin{pmatrix} 3 & 1 & 2 \\ 4 & 8 & 0 \\ 0 & 1 & 2 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 5 & 2 & 1 \\ 1 & 8 & 0 & -6 \\ 1 & 4 & 3 & 7 \end{pmatrix}$.

- a. Compute AB
 b. Compute $f(A)$ if $f(x) = x^2 - 3x + 2 = x^2 - 3x + 2x^0$

4. Find the **inverse** of each of the following matrices.

- a. $\begin{pmatrix} 1 & 5 \\ 2 & -1 \end{pmatrix}$ b. $\begin{pmatrix} 2 & 1 \\ 2 & -4 \end{pmatrix}$ c. $\begin{pmatrix} 1 & -1 & 2 \\ -5 & 7 & -11 \\ -2 & 3 & -5 \end{pmatrix}$ d. $\begin{pmatrix} 1 & -1 & 3 \\ 2 & 0 & 5 \\ -1 & 1 & 0 \end{pmatrix}$

5. Given $A^{-1} = \begin{pmatrix} 1 & -1 & 3 \\ 2 & 0 & 5 \\ -1 & 1 & 0 \end{pmatrix}$. Find a matrix X such that

- a. $AX = \begin{pmatrix} 1 \\ -1 \\ 3 \end{pmatrix}$ b. $AX = \begin{pmatrix} 1 & -1 & 2 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{pmatrix}$ c. $XA = \begin{pmatrix} 1 & 2 & -1 \\ 3 & 1 & 1 \end{pmatrix}$

6. Find A when

- a. $(3A)^{-1} = \begin{pmatrix} 1 & 2 \\ 0 & -2 \end{pmatrix}$ b. $(I + 2A)^{-1} = \begin{pmatrix} 2 & 1 \\ 3 & 2 \end{pmatrix}$ c. $(A^{-1} - 2I)^T = -2 \begin{pmatrix} 1 & 4 \\ 3 & 11 \end{pmatrix}$

7. Write the system of linear equations in matrix form and then solve them.

$$\text{a. } \begin{cases} 2x - y = 4 \\ 3x + 2y = -4 \end{cases}$$

$$\text{b. } \begin{cases} 2x + 3y + z = 10 \\ 2x - 3y - 3z = 22 \\ 4x - 2y + 3z = -2 \end{cases}$$

$$\text{c. } \begin{cases} x + y = a \\ 2x + 3y = 1 - 2a \end{cases} (a \in R)$$

8. Find A^{-1} if

$$\text{a. } A^2 - 6A + 5I = 0$$

$$\text{b. } A^2 + 3A - I = 0$$

$$\text{c. } A^4 = I$$

9. Solve for X

$$\text{a. } \begin{pmatrix} 1 & 2 \\ 2 & 3 \end{pmatrix} X = \begin{pmatrix} 1 & -1 \\ 3 & 3 \end{pmatrix}$$

$$\text{b. } ABXC = B^T$$

$$\text{c. } AX^T BC = B$$

(where A , B and C are $n \times n$ invertible matrices)

$$10. \text{ Compute } \begin{pmatrix} -1 & 3 \\ 0 & 1 \end{pmatrix}^{101}$$

11. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation, and assume that $T(1,2) = (-1,1)$ and $T(0,3) = (-3,3)$

$$\text{a. Compute } T(11,-5)$$

$$\text{b. Compute } T(1,11)$$

$$\text{c. Find the matrix of } T$$

$$\text{d. Compute } T^{-1}(2,3)$$

12. Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation such that the matrix of T is $\begin{pmatrix} 1 & 2 \\ -1 & 3 \end{pmatrix}$.

Find $T^{-1}(3,-2)$

13. The (2;1)-entry of the product $\begin{pmatrix} 1 & 2 & 0 & 1 \\ 0 & 2 & 5 & 1 \\ 4 & -1 & 2 & 3 \end{pmatrix} \begin{pmatrix} 4 & 2 & 1 \\ 2 & 3 & 2 \\ 5 & 1 & 0 \\ 0 & 4 & 3 \end{pmatrix}$

Chapter 3: Determinants and **Diagonalization** → sự chéo hóa

1. Evaluate the determinant

a. $\begin{vmatrix} x-2 & -1 \\ -3 & x \end{vmatrix}$ b. $\begin{vmatrix} -2 & 0 & 0 \\ 4 & 6 & 0 \\ -3 & 7 & 2 \end{vmatrix}$ c. $\begin{vmatrix} -3 & 2 & 1 \\ 4 & 5 & 6 \\ 2 & -3 & 1 \end{vmatrix}$ d. $\begin{vmatrix} 2 & -1 & 1 \\ 0 & 2 & 1 \\ 0 & 0 & 4 \end{vmatrix}$

e. $\begin{vmatrix} x & y & 1 \\ -1 & -2 & 1 \\ 1 & 5 & 1 \end{vmatrix}$ f. $\begin{vmatrix} m & -1 & 0 \\ 1 & 2 & 1 \\ 2 & m & -3 \end{vmatrix}$

2. Find the minors and the cofactors of the matrix

a. $A = \begin{pmatrix} 1 & 3 \\ -2 & 4 \end{pmatrix}$ b. $B = \begin{pmatrix} -3 & 4 & 2 \\ 6 & 3 & 1 \\ 4 & -7 & -8 \end{pmatrix}$ c. $C = \begin{pmatrix} 1 & 3 & 1 \\ 2 & -1 & 1 \\ 1 & 2 & m \end{pmatrix}$

3. Find the adjugate and the inverse of the matrix $A = \begin{pmatrix} 1 & 0 & 3 \\ 0 & -1 & 2 \\ 2 & 1 & 0 \end{pmatrix}$

4. Let $A = \begin{pmatrix} 1 & * & * & * \\ 0 & -1 & * & * \\ 0 & 0 & 2 & * \\ 0 & 0 & 0 & 2 \end{pmatrix}$. Find $|A| = 1 \cdot (-1) \cdot 2 \cdot 2 = -4$

a. $|2A^{-1}| =$

b. $|AA^T| =$

c. $|\text{adj } A| =$

d. $|-A^3| =$

e. $|(2A)^{-1}| =$

f. $|A^{-1} - 2\text{adj } A| =$

5. Let A and B be square matrices of order 4 such that $|A| = -5$ and $|B| = 3$. Find

a. $|2AB| =$

b. $|\text{adj}(AB)| =$

c. $|5A^{-1}B^T| =$

d. $|A^T B^{-1} A^2| =$

6. Find all values of m, k for which the matrix is **not invertible**

a. $A = \begin{pmatrix} 1 & 3 \\ k & 2 \end{pmatrix}$ b. $B = \begin{pmatrix} m & 1 & 3 \\ 1 & 3 & 2 \\ -1 & 4 & 5 \end{pmatrix}$ c. $C = \begin{pmatrix} m & 2 & 0 \\ 1 & m & 1 \\ 2 & 3 & 1 \end{pmatrix}$

7. Find the **characteristic polynomial** of the matrix

a. $A = \begin{pmatrix} 3 & 5 \\ 1 & 2 \end{pmatrix}$ b. $B = \begin{pmatrix} 1 & 3 \\ 1 & 3 \end{pmatrix}$

c. $C = \begin{pmatrix} 1 & 0 & 0 \\ -2 & 2 & 1 \\ 1 & 2 & 1 \end{pmatrix}$ d. $D = \begin{pmatrix} 1 & 2 & -1 \\ 0 & 1 & 2 \\ -1 & 1 & 1 \end{pmatrix}$

8. Find the **eigenvalues** and corresponding **eigenvectors** of the matrix

a. $A = \begin{pmatrix} -3 & 5 \\ 10 & 2 \end{pmatrix}$ b. $B = \begin{pmatrix} 5 & 4 \\ 2 & 1 \end{pmatrix}$

c. $C = \begin{pmatrix} 1 & 0 & 0 \\ -2 & 3 & 0 \\ 0 & 0 & 4 \end{pmatrix}$ d. $D = \begin{pmatrix} -3 & 2 & -1 \\ 0 & 1 & 0 \\ 4 & 1 & 1 \end{pmatrix}$

9. Find the determinant of the matrix $A = \begin{pmatrix} 5 & 1 & 2 & 4 \\ 1 & 0 & -1 & -3 \\ 1 & 1 & 6 & 1 \\ 1 & 0 & 0 & -4 \end{pmatrix}$

10. Find the **(1, 2)-cofactor** and **(3,1) - cofactor** of the matrix $A = \begin{bmatrix} -1 & 3 & -2 \\ 4 & 5 & -7 \\ 7 & 8 & 1 \end{bmatrix}$

11. Let $A = \begin{pmatrix} 1 & 3 & 1 \\ 0 & 1 & 0 \\ 2 & -1 & x \end{pmatrix}$. For which values of x is A **invertible**?

Chapter 5: The Vector Space R^n

1. Let $x = (-1, -2, -2), u = (0, 1, 4), v = (-1, 1, 2)$ and $w = (3, 1, 2)$ in R^3 .

Find scalars a, b and c such that $x = au + bv + cw$

2. Write v as a **linear combination** of u and w , if possible, where $u = (1, 2), w = (1, -1)$

a. $v = (0, 1)$ b. $v = (2, 3)$ c. $v = (1, 4)$ d. $v = (-5, 1)$

3. Determine whether the set S is **linearly independent or linearly dependent** in corresponding vector spaces.

a. $S = \{(-1, 2), (3, 1), (2, 1)\}$ b. $S = \{(-1, 2, 3), (1, 3, 5)\}$

c. $S = \{(1, -2, 2), (2, 3, 5), (3, 1, 7)\}$ d. $S = \{(-1, 2, 1), (2, 4, 0), (3, 1, 1)\}$

e. $S = \{(1, -2, 2, 1), (1, 2, 3, 5), (-1, 3, 1, 7)\}$

4. For which values of k is each set **linearly independent** in corresponding vector spaces.

a. $S = \{(-1, 2, 1), (k, 4, 0), (3, 1, 1)\}$ b. $S = \{(-1, k, 1), (1, 1, 0), (2, -1, 1)\}$

c. $S = \{(k, 1, 1), (1, k, 1), (1, 1, k)\}$ d. $S = \{(1, 2, 1, 0), (-2, 1, 1, -1), (-1, 3, 2, k)\}$

5. Find all values of m such that the set S is a **basis** of R^3

a. $S = \{(1, 2, 1), (m, 1, 0), (-2, 1, 1)\}$ b. $S = \{(-1, m, 1), (1, 1, 0), (m, -1, -1)\}$

6. Find a **basis** for and the **dimension** of the **subspace** U

a. $U = \{(2s - t, s, s + t) \mid s, t \in R\}$ b. $U = \{(s - t, s, t, s + t) \mid s, t \in R\}$

c. $U = \{(0, t, -t) \mid t \in R\}$ d. $U = \{(x, y, z) \mid x + y + z = 0\}$

e. $U = \{(x, y, z) \mid x + y + z = 0, x - y = 0\}$ f. $U = \text{span}\{(1, 2, 3), (2, 3, 4), (3, 5, 7)\}$

g. $U = \text{span}\{(1, 2, 4), (-1, 3, 4), (2, 3, 1)\}$ h. $U = \text{span}\{(1, 2, 1, 1), (2, 1, -1, 0), (3, 3, 0, 1)\}$

7. Find a basis for and the dimension of the solution space of the **homogeneous system of linear equations**.

$$\begin{array}{lll} \text{a. } \begin{cases} -x + y + z = 0 \\ 3x - y = 0 \\ 2x - 4y - 5z = 0 \end{cases} & \text{b. } \begin{cases} x + 2y - 4z = 0 \\ -3x - 6y + 12z = 0 \end{cases} & \text{c. } \begin{cases} x + y + z + t = 0 \\ 2x + 3y + z = 0 \\ 3x + 4y + 2z + t = 0 \end{cases} \end{array}$$

8. Find all values of m for which x lies in the subspace **spanned** by S

a. $x = (-3, 2, m)$ and $S = \{(-1, -1, 1), (2, -3, -4)\}$

b. $x = (4, 5, m)$ and $S = \{(1, -1, 1), (2, -3, 4)\}$

c. $x = (m+1, 5, m)$ and $S = \{(1, 1, 1), (2, 3, 1), (3, 4, 2)\}$

d. $x = (3, 5, 7, m)$ and $S = \{(1, 1, 1, -1), (1, 2, 3, 1), (2, 3, 4, 0)\}$

9. Find the dimension of the subspace

$$U = \text{span}\{(-2, 0, 3), (1, 2, -1), (-2, 8, 5), (-1, 2, 2)\}$$

10. Let $A = \begin{pmatrix} 1 & 2 & 2 & -1 \\ 3 & 6 & 5 & 0 \\ 2 & 2 & 1 & 2 \end{pmatrix}$. Find $\dim(\text{col } A)$ and $\dim(\text{row } A)$

$$\text{col } A = \text{Span}\{(1, 3, 2), (2, 6, 2), (2, 5, 1), (-1, 0, 2)\}$$

$$\text{row } A = \text{Span}\{(1, 2, 2, -1), (3, 6, 5, 0), (2, 2, 1, 2)\}$$

$$\dim(\text{col } A) = \dim(\text{row } A) = \text{rank}(A)$$

11. Which of the following are subspaces of \mathbb{R}^3 ?

(i) $U = \{(2+a, b-a, b) \mid a, b \in \mathbb{R}\}$

(ii) $V = \{(a+b, a, b) \mid a, b \in \mathbb{R}\}$

(iii) $W = \{(2a+b, 0, ab) \mid a, b \in \mathbb{R}\}$

12. Let $u = (1, -3, -2)$, $v = (-1, 1, 0)$ and $w = (1, 2, -3)$. Compute $\|u - v + w\|$

13. Let $u, v \in \mathbb{R}^3$ such that $\|u\| = 3, \|v\| = 4$ and $u \bullet v = -2$. Find

a. $\|u + v\|$ b. $\|2u + 3v\|$

Chapter 4: Vector Geometry

4.1. Vectors and Lines

Bài tập 4.1.2, 4.1.3, 4.1.4, 4.1.7, 4.1.10, 4.1.11, 4.1.12, 4.1.17, 4.1.18, 4.1.19, 4.1.22, 4.1.23, 4.1.24 (từ trang 222 – 225) sách **Linear Algebra**.

4.2. Projections and Planes

Bài tập 4.2.1, 4.2.2, 4.2.3, 4.2.4, 4.2.10, 4.2.11, 4.2.12, 4.2.13, 4.2.14, 4.2.15, 4.2.16, 4.2.19, 4.2.20, 4.2.21, 4.2.32 (từ trang 239 – 244) sách **Linear Algebra**.

4.3. More on Cross Product

Bài tập 4.3.3, 4.3.4, 4.3.5 (từ trang 248 – 249) sách **Linear Algebra**.

1. Find the equations of the line through the points $P_0(2, 0, 1)$ and $P_1(4, -1, 1)$.
2. Find the equations of the line through $P_0(3, -1, 2)$ parallel to the line with equations:

$$\begin{cases} x = -1 + 2t \\ y = 1 + t \\ z = -3 + 4t \end{cases}$$

3. Determine whether the following lines intersect and, if so, find the point of intersection.

$$\begin{cases} x = 1 - 3t \\ y = 2 + 5t \\ z = 1 + t \end{cases}, \begin{cases} x = -1 + s \\ y = 3 - 4s \\ z = 1 - s \end{cases}$$

4. Compute $\|v\|$ if v equals:

a. $(2, -1, 2)$ b. $2(1, 1, -1)$ c. $-3(1, 1, 2)$ d. $(1, 2, 3) - (4, 1, 2)$

5. Find a unit vector in the direction from $(3, -1, 4)$ to $(1, 3, 5)$.

6. Find $\|v - 3w\|$ when $\|v\| = 2$, $\|w\| = 1$, and $v \cdot w = 2$

7. Compute the angle between $u = (-1, 1, 2)$ and $v = (-1, 2, 1)$.

8. Show that the points $P(3, -1, 1)$, $Q(4, 1, 4)$, and $R(6, 0, 4)$ are the vertices of a right triangle.

9. Find the projection of $u = (2, -3, 1)$ on $d = (-1, 1, 3)$ and express $u = u_1 + u_2$ where u_1 is parallel to d and u_2 is orthogonal to d .

10. Find an equation of the plane through $P_0(1, -1, 3)$ with $n = (-3, -1, 2)$ as normal.

11. Find an equation of the plane through $P_0(3, -1, 2)$ that is parallel to the plane with equation $2x - 3y - z = 6$.
12. Find the shortest distance from the point $P(2, -1, -3)$ to the plane with equation $3x - y + 4z = 1$. Also find the point Q on this plane closest to P .
13. Find the equation of the plane through $P(1, 3, -2)$, $Q(1, 1, 5)$, and $R(2, -2, 3)$.
14. Find the shortest distance between the nonparallel lines

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} + t \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 4 \\ 1 \\ -1 \end{bmatrix} + s \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$$

15. Compute $u \cdot v$ where:
- a. $u = (2, -1, 3)$, $v = (-1, 1, 1)$ b. $u = (-2, 1, 4)$, $v = (-1, 5, 1)$
16. Find all real numbers x such that:
- a. $(3, -1, 2)$ and $(3, -2, x)$ are orthogonal.
- b. $(2, -1, 1)$ and $(1, x, 2)$ are at an angle of $\pi/3$.
17. Find the three internal angles of the triangle with vertices:
- a. $A(3, 1, -2)$, $B(3, 0, -1)$, and $C(5, 2, -1)$
- b. $A(3, 1, -2)$, $B(5, 2, -1)$, and $C(4, 3, -3)$
18. Find the equations of the line of intersection of the following planes.
- a. $2x - 3y + 2z = 5$ and $x + 2y - z = 4$.
- b. $3x + y - 2z = 1$ and $x + y + z = 5$.
19. Find the area of the triangle with vertices $P(2, 1, 0)$, $Q(3, -1, 1)$, and $R(1, 0, 1)$
20. Find the volume of the parallelepiped determined by the vectors $u = (1, 2, -1)$, $v = (3, 4, 5)$ and $w = (-1, 2, 4)$.
21. Find the reflection of the point P in the line $y = 1 + 2x$ in \mathbb{R}^2 if:
- a. $P = P(1, 1)$
- b. $P = P(1, 4)$
25. Find the angle between the following pairs of vectors.
- a. $u = (1, -1, 4)$, $v = (5, 2, -1)$
- b. $u = (2, 1, 5)$, $v = (0, 3, 1)$
22. In each case, compute the projection of u on v .

$$\text{a. } \mathbf{u} = \begin{bmatrix} 5 \\ 7 \\ 1 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix}$$

$$\text{b. } \mathbf{u} = \begin{bmatrix} 3 \\ -2 \\ 1 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} 4 \\ 1 \\ 1 \end{bmatrix}$$

$$\text{c. } \mathbf{u} = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}$$

$$\text{d. } \mathbf{u} = \begin{bmatrix} 3 \\ -2 \\ -1 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} -6 \\ 4 \\ 2 \end{bmatrix}$$

23. Find the shortest distance between the following pairs of nonparallel lines and find the points on the lines that are closest together.

$$\text{a. } \begin{cases} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + s \begin{bmatrix} 2 \\ 1 \\ -3 \end{bmatrix} \\ \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} + t \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \end{cases};$$

$$\text{b. } \begin{cases} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + s \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \\ \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix} + t \begin{bmatrix} 3 \\ 1 \\ 0 \end{bmatrix} \end{cases};$$

$$\text{c. } \begin{cases} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ -1 \end{bmatrix} + s \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} \\ \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} + t \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} \end{cases};$$

$$\text{d. } \begin{cases} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + s \begin{bmatrix} 2 \\ 0 \\ -1 \end{bmatrix} \\ \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ -1 \\ 0 \end{bmatrix} + t \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \end{cases};$$