Solutions to Linear Algebra Problems

Minh Le

April 14, 2026

Abstract

This note collects my solutions to selected problems in Linear Algebra. The goal is to present clear arguments and organized solutions for study and review. Books that I use for assignments sources include:

- Linear Algebra Done Right 4e, Sheldon Axler



Problem 1.

Prove that the union of two subspaces of V is a subspace of V if and only if one of the subspaces is contained in the other.

Source: Linear Algebra Done Right - Ex.12 p.25

Solution:

Let V_1, V_2 be two subspaces of V.
($\Rightarrow$): Suppose that $V_1 \cup V_2$ is a subspace of V. Suppose for contradiction that one of the subspaces V_1 or V_2 is not contained in the other. We can choose u, w such that $u \in V_1, u \notin V_2$ and $w \in V_2, w \notin V_1$.
Since $V_1 \cup V_2$ is a subspace, we have $u + w \in V_1 \cup V_2$. This means $u + w$ must be in V_1 or V_2.
If $u + w \in V_1$, since V_1 is a subspace and $-u \in V_1$, then $w = (u + w) - u \in V_1$, which is a contradiction. Similarly, if $u+w \in V_2$, since V_2 is a subspace and $-w \in V_2$, then $u = (u+w)-w \in V_2$, which is also a contradiction.
Therefore, $V_1 \subseteq V_2$ or $V_2 \subseteq V_1$.
($\Leftarrow$): If $V_1 \subseteq V_2$ or $V_2 \subseteq V_1$, then $V_1 \cup V_2 = V_2$ or $V_1 \cup V_2 = V_1$, which is a subspace.

Problem 2.

Suppose $v_1, \ldots, v_m$ is a list of vectors in V. For $k \in \{1, \ldots, m\}$, let

$$w_k = v_1 + \cdots + v_k.$$

Show that the list $v_1, \ldots, v_m$ is linearly independent if and only if the list $w_1, \ldots, w_m$ is linearly independent.

Source: Linear Algebra Done Right - Ex.14 p.38

Solution:

($\Rightarrow$): Suppose the list $v_1, \ldots, v_m$ is linearly independent. Consider a combination of $w_1, \ldots, w_m$ such that:

$$\begin{aligned} c_1w_1 + c_2w_2 + \cdots + c_mw_m &= 0 \\ \Rightarrow c_1v_1 + c_2(v_1 + v_2) + \cdots + c_m(v_1 + v_2 + \cdots + v_m) &= 0 \\ \Rightarrow (c_1 + c_2 + \cdots + c_m)v_1 + (c_2 + \cdots + c_m)v_2 + \cdots + c_mv_m &= 0 \end{aligned}$$

Since $v_1, \ldots, v_m$ are linearly independent, it follows that $c_1 = c_2 = \cdots = c_m = 0$. Therefore, the list $w_1, \ldots, w_m$ are linearly independent.
($\Leftarrow$): Suppose $w_1, \ldots, w_m$ are linearly independent. Consider a combination of $v_1, \ldots, v_m$ such

that:

$$
\begin{aligned}
c_1v_1 + c_2v_2 + \cdots + c_mv_m &= 0 \\
\Rightarrow c_1w_1 + c_2(w_2 - w_1) + c_3(w_3 - w_2) + \cdots + c_m(w_m - w_{m-1}) &= 0 \\
\Rightarrow (c_1 - c_2)w_1 + (c_2 - c_3)w_2 + \cdots + (c_{m-1} - c_m)w_{m-1} + c_mw_m &= 0
\end{aligned}
$$

Since $w_1, \ldots, w_m$ are linearly independent, it follows that $c_1 = c_2 = \cdots = c_m = 0$. Therefore, the list $v_1, \ldots, v_m$ are linearly independent.