

Linear Algebra

Chapter 4: Eigenvalues and Eigenvectors

University of Seoul
School of Computer Science
Minho Kim

Table of contents

Introduction: A Dynamical System on Graphs

Introduction to Eigenvalues and Eigenvectors

Determinants

Eigenvalues and Eigenvectors of $n \times n$ Matrices

Similarity and Diagonalization

Iterative Methods for Computing Eigenvalues

Applications and the Perron-Frobenius Theorem

Outline

Introduction: A Dynamical System on Graphs

Introduction to Eigenvalues and Eigenvectors

Determinants

Eigenvalues and Eigenvectors of $n \times n$ Matrices

Similarity and Diagonalization

Iterative Methods for Computing Eigenvalues

Applications and the Perron-Frobenius Theorem

Dynamical System

- ▶ For $A \in \mathbb{R}^{n \times n}$, $\lim_{k \rightarrow \infty} A^k = ?$
 - ▶ Try the Octave demos yourselves!
1. Complete graphs (K4.m)
 2. Petersen graph (Petersen.m)
 3. Cyclic graphs
 - 3.1 Odd number of nodes (C5.m)
 - 3.2 Even number of nodes (C6.m)
 4. Complete bipartite graphs (K3_3.m)
 - ▶ Steady state vector

Outline

Introduction: A Dynamical System on Graphs

Introduction to Eigenvalues and Eigenvectors

Determinants

Eigenvalues and Eigenvectors of $n \times n$ Matrices

Similarity and Diagonalization

Iterative Methods for Computing Eigenvalues

Applications and the Perron-Frobenius Theorem

Eigenvalue Problem

- ▶ For a square matrix A , are there nonzero vectors x such that Ax is just a scalar multiplication of x ? In other words, which nonzero vectors satisfy $Ax = \lambda x$? ($\lambda \in \mathbb{R}$) →
“**Eigenvalue problem**”
- ▶ eigen- [áigən]: “own” or “characteristic of”

Definition

Let A be an $n \times n$ matrix. A scalar λ is called an **eigenvalue** of A if there is a nonzero x such that $Ax = \lambda x$. Such a vector x is called an **eigenvector** of A corresponding to λ .

- ▶ Why are they important?
- ▶ Do they exist for *any* matrix?
- ▶ Is there only one eigenvector for an eigenvalue?
- ▶ Is there only one eigenvalue for an eigenvector?
- ▶ Given an eigenvalue, how can we find the corresponding eigenvectors? → Example 4.2
- ▶ How can we find eigenvalues?

Eigenspace

- ▶ Example 4.2

→ “The set of all eigenvectors corresponding to an eigenvalue λ of an $n \times n$ matrix A is just the set of *nonzero* vectors in $\text{null}(A - \lambda I)$.”

Definition

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A . The collection of all eigenvectors corresponding to λ , together with the zero vector, is called the **eigenspace** of λ and is denoted by E_λ .

Geometric Interpretation of Eigenvectors

- ▶ Ax and λx are parallel, i.e., x is an eigenvector of A iff T_A transforms x into a parallel vector.
- ▶ Examples: Scaling, reflection (Ex 4.4), rotation
- ▶ Only the direction of an eigenvector matters. (Why?)
→ Only unit vectors need to be considered. (Fig 4.7)

Finding Eigenvalues

- ▶ For $A \in \mathbb{R}^{n \times n}$, the eigenvectors of λ are the nonzero vectors satisfying $Ax = \lambda x$.
 - $Ax - \lambda Ix = \mathbf{0} \rightarrow (A - \lambda I)x = \mathbf{0}$
 - The eigenspace E_λ is the non-trivial null space of $A - \lambda I$.
- ▶ For 2×2 matrices,
 - a matrix has a non-trivial null space iff it is non-invertible.
 - a matrix is non-invertible iff its determinant is zero.
 - λ is an eigenvalue of A iff $\det(A - \lambda I) = 0$ (Example 4.5)
- ▶ Can be generalized to any square matrices.
 - What are the determinants for $n \times n$ matrices?

Outline

Introduction: A Dynamical System on Graphs

Introduction to Eigenvalues and Eigenvectors

Determinants

Eigenvalues and Eigenvectors of $n \times n$ Matrices

Similarity and Diagonalization

Iterative Methods for Computing Eigenvalues

Applications and the Perron-Frobenius Theorem

Determinants

- ▶ Notation: $\det A = |A|$
- ▶ 1×1 matrices

$$\det A = |a| = a \quad (\text{Not the absolute value})$$

- ▶ 2×2 matrices

$$\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} = a_{11}a_{22} - a_{12}a_{21}$$

- ▶ 3×3 matrices

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = ?$$

Determinant of a 3×3 Matrix

Definition

Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$. Then the **determinant** of A is the scalar

$$\det A = |A| = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

$$\bullet \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \rightarrow a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix}$$

$$\bullet \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \rightarrow a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix}$$

$$\bullet \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \rightarrow a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

Determinant of a 3×3 Matrix (cont'd)

- ▶ With A_{ij} defined as the submatrix of A obtained by deleting row i and column j ,

$$\begin{aligned}\det A &= a_{11} \det A_{11} - a_{12} \det A_{12} + a_{13} \det A_{13} \\ &= \sum_{j=1}^3 (-1)^{1+j} a_{1j} \det A_{1j}\end{aligned}$$

- ▶ $\det A_{ij}$ is called the (i, j) -**minor** of A .
- ▶ Computed with respect to the first row.
 - Why row not column? Why the first row?
 - Can be generalized to columns or other rows (The Laplace Expansion Theorem)
- ▶ Another method (See (2) on p.264)

$$\begin{aligned}|A| &= a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} \\ &\quad - a_{31}a_{22}a_{13} - a_{32}a_{23}a_{11} - a_{33}a_{21}a_{12}\end{aligned}$$

Determinants of $n \times n$ Matrices

Definition

Let $A = [a_{ij}]$ be an $n \times n$ matrix, where $n \geq 2$. Then the **determinant** of A is the scalar

$$\begin{aligned}\det A = |A| &= a_{11} \det A_{11} - a_{12} \det A_{12} + \cdots + (-1)^{1+n} a_{1n} \det A_{1n} \\ &= \sum_{j=1}^n (-1)^{1+j} a_{1j} \det A_{1j}\end{aligned}$$

- **Cofactor expansion along the first row:**

With (i, j) -**cofactor** of A defined as

$$C_{ij} = (-1)^{i+j} \det A_{ij},$$

the definition becomes

$$\det A = \sum_{j=1}^n a_{1j} C_{1j}.$$

The Laplace Expansion Theorem

Theorem 4.1: The Laplace Expansion Theorem

The determinant of an $n \times n$ matrix $A = [a_{ij}]$, where $n \geq 2$, can be computed as

$$\det A = a_{i1}C_{i1} + a_{i2}C_{i2} + \cdots + a_{in}C_{in} = \sum_{j=1}^n a_{ij}C_{ij}$$

(which is the **cofactor expansion along the i th row**) and also as

$$\det A = a_{1j}C_{1j} + a_{2j}C_{2j} + \cdots + a_{nj}C_{nj} = \sum_{i=1}^n a_{ij}C_{ij}$$

(the **cofactor expansion along the j th column**)

- ▶ Most useful when the matrix contains a row or column with lots of zeros. Why? (Example 4.11)

Determinants of Triangular Matrices

- ▶ The Laplace expansion theorem is particularly useful when the matrix is (upper or lower) triangular.

Theorem 4.2

The determinant of a triangular matrix is the product of the entries on its main diagonal. Specifically, if $A = [a_{ij}]$ is an $n \times n$ triangular matrix then

$$\det A = a_{11}a_{22} \cdots a_{nn}$$

- ▶ Why? (Example 4.12)

Computing Determinants

- ▶ Laplace expansion is very inefficient! (p.268)
- ▶ The determinant of a triangular matrix can be easily found.
- ▶ We can compute the determinant of a matrix efficiently from its reduced form.
 - “How does the determinant change after elementary row operations?”

Properties of Determinants

Theorem 4.3

Let $A = [a_{ij}]$ be a square matrix.

- a. If A has a zero row (column), then $\det A = 0$.
- b. If B is obtained by interchanging two rows (columns) of A , then $\det B = -\det A$.
- c. If A has two identical rows (columns), then $\det A = 0$.
- d. If B is obtained by multiplying a row (column) of A by k , then $\det B = k \det A$.
- e. If A , B , and C are identical except that the i th row (column) of C is the sum of the i th rows (columns) of A and B , then $\det C = \det A + \det B$.
- f. If B is obtained by adding a multiple of one row (column) of A to another row (column), then $\det B = \det A$.

Determinants of Elementary Matrices

- ▶ (b), (d) and (f) of the properties are related to elementary row operations.
- ▶ What are the determinants of elementary matrices?

Theorem 4.4

Let E be an $n \times n$ elementary matrix.

- If E results from interchanging two rows of I_n , then $\det E = -1$.
 - If E results from multiplying one row of I_n by k , then $\det E = k$.
 - If E results from adding a multiple of one row of I_n to another row, then $\det E = 1$.
- ▶ Determinants of all the elementary matrices are nonzero.

Determinants of Elementary Matrices (cont'd)

- ▶ If $B = EA$, $\det B = ?$

Lemma 4.5

Let B be an $n \times n$ matrix and let E be an $n \times n$ elementary matrix. Then

$$\det (EB) = (\det E)(\det B)$$

- ▶ How about $\det (AB)$ when A is NOT an elementary matrix? \rightarrow Theorem 4.8 (p.272)

Theorem 4.6

A square matrix A is invertible if and only if $\det A \neq 0$.

Determinants and Matrix Operations

How can we write the followings in terms of $\det A$ and $\det B$?

- ▶ $\det(kA) = ?$
- ▶ $\det(A + B) = ?$
- ▶ $\det(AB) = ?$
- ▶ $\det(A^{-1}) = ?$
- ▶ $\det(A^T) = ?$

$\det(kA)$ and $\det(A + B)$

- ▶ $\det(kA)$

Theorem 4.7

If A is an $n \times n$ matrix, then

$$\det(kA) = k^n \det A$$

- ▶ See Theorem 4.3(d).
- ▶ $\det(A + B)$
 - ▶ $\det(A + B) = \det A + \det B$? → No!
 - ▶ No general formula. (p.271)

$\det(AB)$, $\det(A^{-1})$ and $\det(A^T)$

- ▶ $\det(AB)$

Theorem 4.8

If A and B are $n \times n$ matrices, then

$$\det(AB) = (\det A)(\det B)$$

- ▶ $\det(A^{-1})$

Theorem 4.9

If A is invertible, then

$$\det(A^{-1}) = \frac{1}{\det A}$$

- ▶ $\det(A^T)$

Theorem 4.10

For any square matrix A ,

$$\det A = \det A^T$$

Cramer's Rule and the Adjoint

- ▶ What is the relation between determinants and the solution of a linear systems? → Cramer's rule (Theorem 4.11)
- ▶ What is the relation between determinants and the inverse of a matrix? → Adjoint (Theorem 4.12)
- ▶ Notation
 $A_i(\mathbf{b})$: the matrix obtained by replacing the i th column of A by \mathbf{b}

$$A_i(\mathbf{b}) = [\mathbf{a}_1 \quad \cdots \quad \mathbf{b} \quad \cdots \quad \mathbf{a}_n]$$

Cramer's Rule

Theorem 4.11: Cramer's Rule

Let A be an invertible $n \times n$ matrix and let \mathbf{b} be a vector in \mathbb{R}^n . Then the unique solution \mathbf{x} of the system $A\mathbf{x} = \mathbf{b}$ is given by

$$x_i = \frac{\det(A_i(\mathbf{b}))}{\det A} \quad \text{for } i = 1, \dots, n$$

- Requires to compute determinants
→ Computationally inefficient except small systems.

Adjoint

1. What is the formula of the inverse of a matrix in terms of determinants?
2. What is the solution of the equation $AX = I$?

$$A\mathbf{x}_1 = \mathbf{e}_1 \quad A\mathbf{x}_2 = \mathbf{e}_2 \quad \cdots \quad A\mathbf{x}_n = \mathbf{e}_n$$

3. By the Cramer's rule, $x_{ij} = \frac{\det(A_i(\mathbf{e}_j))}{\det A}$
4. $\det(A_i(\mathbf{e}_j)) = (-1)^{j+i} \det A_{ji} = C_{ji}$ (Why?)
5. $A^{-1} = X = \frac{1}{\det A} [C_{ji}] = \frac{1}{\det A} [C_{ij}]^T = \frac{1}{\det A} \operatorname{adj} A$

Adjoint (cont'd)

Theorem 4.12

Let A be an invertible $n \times n$ matrix. Then

$$A^{-1} = \frac{1}{\det A} \operatorname{adj} A$$

- ▶ $\operatorname{adj} A := [C_{ji}] = [C_{ij}]^T$: the **adjoint** (or **adjugate**) of A
 - ▶ $C_{ij} := (-1)^{i+j} \det A_{ij}$: (i, j) -cofactor of A

Outline

Introduction: A Dynamical System on Graphs

Introduction to Eigenvalues and Eigenvectors

Determinants

Eigenvalues and Eigenvectors of $n \times n$ Matrices

Similarity and Diagonalization

Iterative Methods for Computing Eigenvalues

Applications and the Perron-Frobenius Theorem

Finding Eigenvalues

- ▶ How to compute eigenvalues of a matrix?
- ▶ How many eigenvalues does a matrix have?
- ▶ λ is an eigenvalue of A iff $\det(A - \lambda I) = 0$.

The eigenvalues of a square matrix A are precisely the solutions λ of the equation

$$\det(A - \lambda I) = 0$$

- ▶ What does $\det(A - \lambda I)$ look like?
 - A polynomial in λ of degree n
(**Characteristic polynomial** of A)
 - At most n distinct eigenvalues

Finding Eigenvalues and Eigenvectors

Procedure

Let A be an $n \times n$ matrix.

1. Compute the characteristic polynomial $\det(A - \lambda I)$ of A .
2. Find the eigenvalues of A by solving the characteristic equation $\det(A - \lambda I) = 0$ for λ .
3. For each eigenvalue λ , find the null space of the matrix $A - \lambda I$. This is the eigenspace E_λ , the nonzero vectors of which are the eigenvectors of A corresponding to λ .
4. Find a basis for each eigenspace.
 - ▶ **Algebraic multiplicity** of an eigenvalue: multiplicity as a root of the characteristic equation.
 - ▶ **Geometric multiplicity** of an eigenvalue λ : $\dim E_\lambda$
 - ▶ What's the relation between the algebraic & geometric multiplicities? (Example 4.18 & 4.19)
→ Geometric multiplicity \leq Algebraic multiplicity
(Lemma 4.26 on p.303)

Eigenvalues of Triangular Matrices

- ▶ How does the characteristic equation look like if A is triangular? (See Theorem 4.2 on p.268)
 $\rightarrow (a_{11} - \lambda)(a_{22} - \lambda) \cdots (a_{nn} - \lambda) = 0$

Theorem 4.15

The eigenvalues of a triangular matrix are the entries on its main diagonal.

What Does Eigenvalues Tell Us?

Theorem 4.16

A square matrix A is invertible if and only if 0 is *not* an eigenvalue of A .

- ▶ Why?
- ▶ ...and there will be more (about the importance of eigenvalues).

Fundamental Theorem of Invertible Matrices: Ver. 3

Theorem 3.27

Let A be an $n \times n$ matrix. The following statements are equivalent:

- a. A is invertible.
- b. $A\mathbf{x} = \mathbf{b}$ has a unique solution for every \mathbf{b} in \mathbb{R}^n .
- c. $A\mathbf{x} = \mathbf{0}$ has only the trivial solution.
- d. The reduced row echelon form of A is I_n .
- e. A is a product of elementary matrices.
- f. $\text{rank}(A) = n$
- g. $\text{nullity}(A) = 0$
- h. The column vectors of A are linearly independent.
 - i. The column vectors of A span \mathbb{R}^n .
 - j. The column vectors of A form a basis for \mathbb{R}^n .
- k. The row vectors of A are linearly independent.
 - l. The row vectors of A span \mathbb{R}^n .
- m. The row vectors of A form a basis for \mathbb{R}^n .
- n. $\det A \neq 0$
- o. 0 is not an eigenvalue of A .

Eigenvalue of A^k and A^{-1}

Theorem 4.18

Let A be a square matrix with eigenvalue λ and corresponding eigenvector x .

- a. For any positive integer k , λ^k is an eigenvalue of A^k with corresponding eigenvector x .
 - b. If A is invertible, then $1/\lambda$ is an eigenvalue of A^{-1} with corresponding eigenvector x .
 - c. For any integer k , λ^k is an eigenvalue of A^k with corresponding eigenvector x .
- Application: Computing $A^k x$ where x is not an eigenvector of A . (Example 4.21)
- Is this possible for any x ?

Computing $A^k x$

Theorem 4.19

Suppose the $n \times n$ matrix A has eigenvectors v_1, v_2, \dots, v_m with corresponding eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_m$. If x is a vector in \mathbb{R}^n that can be expressed as a linear combination of these eigenvectors—say,

$$x = c_1 v_1 + c_2 v_2 + \dots + c_m v_m$$

then, for any integer k ,

$$A^k x = c_1 \lambda_1^k v_1 + c_2 \lambda_2^k v_2 + \dots + c_m \lambda_m^k v_m$$

- ▶ When does it work for any x ?

Theorem 4.20

Let A be an $n \times n$ matrix and let $\lambda_1, \lambda_2, \dots, \lambda_m$ be distinct eigenvalues of A with corresponding eigenvectors v_1, v_2, \dots, v_m . Then v_1, v_2, \dots, v_m are linearly independent.

Outline

Introduction: A Dynamical System on Graphs

Introduction to Eigenvalues and Eigenvectors

Determinants

Eigenvalues and Eigenvectors of $n \times n$ Matrices

Similarity and Diagonalization

Iterative Methods for Computing Eigenvalues

Applications and the Perron-Frobenius Theorem

Why Diagonalize Matrices?

- ▶ Triangular and diagonal matrices are good.
 - How can we relate a square matrix to a triangular or diagonal one keeping the eigenvalues?
- ▶ Gaussian elimination?
 - Eigenvalues are not preserved.
- ▶ Diagonalization

Similar Matrices

Definition

Let A and B be $n \times n$ matrices. We say that A **is similar to** B if there is an invertible $n \times n$ matrix P such that $P^{-1}AP = B$. If A is similar to B , we write $A \sim B$.

- ▶ Equivalent to “ $A = PBP^{-1}$ ” or “ $AP = PB$.”
- ▶ P depends on A and B . Is it unique?

Theorem 4.21

Let A , B and C be $n \times n$ matrices.

- $A \sim A$. (Reflexivity)
 - If $A \sim B$, then $B \sim A$. (Symmetry)
 - If $A \sim B$ and $B \sim C$, then $A \sim C$. (Transitivity)
- ▶ **Equivalent relation**

Similar Matrices (cont'd)

Theorem 4.22

Let A and B be $n \times n$ matrices with $A \sim B$. Then

- a. $\det A = \det B$.
 - b. A is invertible if and only if B is invertible.
 - c. A and B have the same rank.
 - d. A and B have the same characteristic polynomial.
 - e. A and B have the same eigenvalues.
- ▶ The converse is not necessarily true. (See Remark on p.300.)
 - ▶ Useful when showing two matrices are not similar. (How?)

Diagonalization

- ▶ Good if a square matrix is similar to a diagonal matrix.
- ▶ Is it always possible?
- ▶ How can we find P ?

Definition

An $n \times n$ matrix A is **diagonalizable** if there is a diagonal matrix D such that A is similar to D —that is, if there is an invertible $n \times n$ matrix P such that $P^{-1}AP = D$.

Diagonalization (cont'd)

- ▶ How can we find D and P ?

Theorem 4.23

Let A be an $n \times n$ matrix. Then A is diagonalizable if and only if A has n linearly independent eigenvectors.

More precisely, there exist an invertible matrix P and a diagonal matrix D such that $P^{-1}AP = D$ if and only if the columns of P are n linearly independent eigenvectors of A and the diagonal entries of D are eigenvalues of A corresponding to the eigenvectors in P in the same order.

- ▶ Is a non-invertible matrix diagonalizable? (Example 4.26)

Diagonalization (cont'd)

- ▶ How can we check if the eigenvectors are linearly independent? (See the 2nd remark below Example 4.26)

Theorem 4.24

Let A be an $n \times n$ matrix and let $\lambda_1, \lambda_2, \dots, \lambda_k$ be distinct eigenvalues of A . If \mathcal{B}_i is a basis for the eigenspace E_{λ_i} , then $\mathcal{B} = \mathcal{B}_1 \cup \mathcal{B}_2 \cup \dots \cup \mathcal{B}_k$ (i.e., the total collection of basis vectors for all the eigenspaces) is linearly independent.

- ▶ We don't have to check the linear independence of eigenvectors associated with different eigenvalues.

Theorem 4.25

If A is an $n \times n$ matrix with n distinct eigenvalues, then A is diagonalizable.

- ▶ The converse is not always true.

The Diagonalization Theorem

Lemma 4.26

If A is an $n \times n$ matrix, then the geometric multiplicity of each eigenvalue is less than or equal to its algebraic multiplicity.

- ▶ The sum of the algebraic multiplicities is always n .
Therefore, A is diagonalizable when...?

Theorem 4.27

Let A be an $n \times n$ matrix whose distinct eigenvalues are $\lambda_1, \lambda_2, \dots, \lambda_k$. The following statements are equivalent:

- A is diagonalizable.
 - The union \mathcal{B} of the bases of the eigenspaces of A (as in Theorem 4.24) contains n vectors.
 - The algebraic multiplicity of each eigenvalue equals its geometric multiplicity.
- ▶ Computing A^k (Example 4.29)
 $A^k = PD^kP^{-1}$ for all $k \geq 1$.

Computing $A^k \mathbf{x}$

- ▶ When \mathbf{x} is an eigenvector of A . (Theorem 4.18)

$$A^k \mathbf{x} = \lambda^k \mathbf{x}$$

- ▶ When \mathbf{x} is a linear combination of the eigenvectors of A . (Theorem 4.19)

$$\mathbf{x} = \sum_{j=1}^m c_j \mathbf{v}_j \rightarrow A^k \mathbf{x} = \sum_{j=1}^m (c_j A^k \mathbf{v}_j) = \sum_{j=1}^m (c_j \lambda_j^k \mathbf{v}_j)$$

- ▶ When A is diagonalizable. (Example 4.29)

$$A^k \mathbf{x} = (PDP^{-1})^k \mathbf{x} = PD^k P^{-1} \mathbf{x}$$

- ▶ Otherwise... Good luck!

Outline

Introduction: A Dynamical System on Graphs

Introduction to Eigenvalues and Eigenvectors

Determinants

Eigenvalues and Eigenvectors of $n \times n$ Matrices

Similarity and Diagonalization

Iterative Methods for Computing Eigenvalues

Applications and the Perron-Frobenius Theorem

Outline

Introduction: A Dynamical System on Graphs

Introduction to Eigenvalues and Eigenvectors

Determinants

Eigenvalues and Eigenvectors of $n \times n$ Matrices

Similarity and Diagonalization

Iterative Methods for Computing Eigenvalues

Applications and the Perron-Frobenius Theorem

Markov Chains

$$\mathbf{x}_{k+1} = P\mathbf{x}_k \quad \rightarrow \quad \mathbf{x}_k = P^k \mathbf{x}_0$$

- ▶ P : transition matrix
- ▶ All the components in each column of P add up to 1.
(Why?)
- ▶ \mathbf{x}_k : **state vector** cf) **probability vector**
- ▶ Here, we will see...
 - ▶ Steady state vector \mathbf{x} : $P\mathbf{x} = \mathbf{x}$.
→ “Every Markov chain has a unique steady state vector.”
(p.231)
 - ▶ $\lim_{k \rightarrow \infty} \mathbf{x}_k = \lim_{k \rightarrow \infty} P^k \mathbf{x}_0 = ?$
→ $\lim_{k \rightarrow \infty} P^k \mathbf{x}_0 = \mathbf{x}$
: \mathbf{x}_k converges to \mathbf{x} regardless of \mathbf{x}_0 .

Markov Chains (cont'd)

Theorem 4.30

If P is the $n \times n$ transition matrix of a Markov chain, then 1 is an eigenvalue of P .

- ▶ There always exists a vector x such that $Px = x$.
→ There always exists a steady state vector. But is it unique?

Theorem 4.31

Let P be an $n \times n$ transition matrix with eigenvalue λ .

- $|\lambda| \leq 1$
- If P is regular and $\lambda \neq 1$, then $|\lambda| < 1$.
 - ▶ **Positive matrix:** All the entries are positive.
 - ▶ **Regular matrix:** P^k is positive for some k .
 - ▶ If P is regular, -1 cannot be an eigenvalue.

Markov Chains (cont'd)

Lemma 4.32

Let P is a regular $n \times n$ transition matrix. If P is diagonalizable, then the dominant eigenvalue $\lambda_1 = 1$ has algebraic multiplicity 1.

- ▶ There is only one eigenvector (and its scalar multiplications) such that $Px = x$.
→ the steady state vector is unique.

Markov Chains (cont'd)

Theorem 4.33

Let P be a regular $n \times n$ transition matrix. Then as $k \rightarrow \infty$, P^k approaches an $n \times n$ matrix L whose columns are identical, each equal to the same vector x . This vector x is a steady state probability vector for P .

- ▶ $\lim_{k \rightarrow \infty} P^k = \begin{bmatrix} x & \cdots & x \end{bmatrix} =: L$
→ $\lim_{k \rightarrow \infty} P^k$ converges. But what is $\lim_{k \rightarrow \infty} P^k x_0 = ?$

Theorem 4.34

Let P be a regular $n \times n$ transition matrix, with x the steady state probability vector for P , as in Theorem 4.33. Then, for any initial probability vector x_0 , the sequence of iterates x_k approaches x .

- ▶ $\lim_{k \rightarrow \infty} P^k x_0 = x$ for any x_0 (initial probability vector)

Population Growth

$$L = \begin{bmatrix} b_1 & b_2 & b_3 & \cdots & b_{n-1} & b_n \\ s_1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & s_2 & 0 & \cdots & 0 & 0 \\ 0 & 0 & s_3 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & s_{n-1} & 0 \end{bmatrix}$$

- ▶ Leslie matrix
- ▶ “The proportion of the population in each class is approaching a steady state.” (p.235)
 - There exists a vector such that $Lx = \lambda x$ where $\lambda > 0$.

Theorem 4.35

Every Leslie matrix has a unique positive eigenvalue and a corresponding eigenvector with positive components.