

Topics in Computer Graphics
Chap 7: Polynomial Curve Constructions
fall, 2011

University of Seoul
School of Computer Science
Minho Kim

Table of contents

Aitken's Algorithm

Lagrange Polynomials

The Vandermonde Approach

Limits of Lagrange Interpolation

Cubic Hermite Interpolation

Quintic Hermite Interpolation

Point-Normal Interpolation

Least Squares Approximation

Smoothing Equations

Designing with Bézier Curves

The Newton Form and Forward Differencing

Motivation

- ▶ Given n points in 2D/3D,
 - ▶ Interpolation: find a (polynomial) curve that passes through the data exactly.
 - ▶ Aitken's algorithm (7.1)
 - ▶ Lagrange polynomials (7.2)
 - ▶ The Vandermonde approach (7.3)
 - ▶ The Newton form and forward differencing (7.11)
 - ▶ Approximation: find a (polynomial) curve that goes through the data closely in a reasonable way.
 - ▶ Least squares approximation (7.8)
 - ▶ Smoothing equations (7.9)
- ▶ Normal/tangent data in addition to position data.
 - ▶ Cubic/quintic Hermite interpolation (7.5&7.6)
 - ▶ Point-normal interpolation (7.7)

Outline

Aitken's Algorithm

Lagrange Polynomials

The Vandermonde Approach

Limits of Lagrange Interpolation

Cubic Hermite Interpolation

Quintic Hermite Interpolation

Point-Normal Interpolation

Least Squares Approximation

Smoothing Equations

Designing with Bézier Curves

The Newton Form and Forward Differencing

Aitken's Algorithm

- ▶ Problem: Given points $\{\mathbf{p}_i\}_{i=0}^n$ with corresponding parameter values $\{t_i\}_{i=0}^n$, find a curve $\mathbf{p}(t)$ such that

$$\mathbf{p}(t_i) = \mathbf{p}_i, \quad i = 0, \dots, n.$$

Aitken's Algorithm (cont'd)

1. Base case: Let

$$\mathbf{p}_i^1(t) = \frac{t_{i+1} - t}{t_{i+1} - t_i} \mathbf{p}_i + \frac{t - t_i}{t_{i+1} - t_i} \mathbf{p}_{i+1}, \quad i = 0, \dots, n-1.$$

- ▶ $\mathbf{p}_i^1(t)$ is a line segment for $t_i \leq t \leq t_{i+1}$.
- ▶ $\mathbf{p}_i^1(t)$ interpolates \mathbf{p}_i and \mathbf{p}_{i+1} : $\mathbf{p}_i^1(t_i) = \mathbf{p}_i$ and $\mathbf{p}_i^1(t_{i+1}) = \mathbf{p}_{i+1}$.

2. Let $\mathbf{p}_0^{n-1}(t)$ is a polynomial of degree $n-1$ that interpolates $\mathbf{p}_0, \dots, \mathbf{p}_{n-1}$ and $\mathbf{p}_1^{n-1}(t)$ is one that interpolates $\mathbf{p}_1, \dots, \mathbf{p}_n$.

3. Then

$$\mathbf{p}_0^n(t) = \frac{t_n - t}{t_n - t_0} \mathbf{p}_0^{n-1}(t) + \frac{t - t_0}{t_n - t_0} \mathbf{p}_1^{n-1}(t)$$

- ▶ is a polynomial curve of degree n and
- ▶ interpolates $\mathbf{p}_0, \dots, \mathbf{p}_n$.

Aitken's Algorithm (cont'd)

With $\mathbf{p}_i^0(t) := \mathbf{p}_i$, the interpolating curve $\mathbf{p}_0^n(t)$ is defined recursively as

$$\mathbf{p}_i^r(t) = \frac{t_{i+r} - t}{t_{i+r} - t_i} \mathbf{p}_i^{r-1}(t) + \frac{t - t_i}{t_{i+r} - t_i} \mathbf{p}_{i+1}^{r-1}(t), \quad \begin{cases} r = 1, \dots, n \\ i = 0, \dots, n - r. \end{cases}$$

Properties

- ▶ Affine invariance
- ▶ Linear precision
- ▶ No convex hull property
 - Is it possible for a smooth interpolating curve has a convex hull property?
- ▶ No variation diminishing property
- ▶ Does not answer if such polynomial curve is unique.
- ▶ Not a closed form

Outline

Aitken's Algorithm

Lagrange Polynomials

The Vandermonde Approach

Limits of Lagrange Interpolation

Cubic Hermite Interpolation

Quintic Hermite Interpolation

Point-Normal Interpolation

Least Squares Approximation

Smoothing Equations

Designing with Bézier Curves

The Newton Form and Forward Differencing

Lagrange Polynomials

The interpolating curve is given by

$$\mathbf{p}(t) = \sum_{i=0}^n \mathbf{p}_i L_i^n(t)$$

where $L_i^n(t)$ are *Lagrange polynomials* defined as

$$L_i^n(t) := \frac{\prod_{\substack{j=0 \\ j \neq i}}^n (t - t_j)}{\prod_{\substack{j=0 \\ j \neq i}}^n (t_i - t_j)}$$

- ▶ Lagrange polynomials are *cardinal*:

$$L_i^n(t_j) = \delta_{i,j}$$

where $\delta_{i,j}$ is the Kronecker delta.

- ▶ $\{L_i^n(t)\}_{i=0}^n$ form a basis of all polynomials of degree n
- ▶ Interpolating polynomial (of degree n) is unique
- ▶ $\sum_{i=0}^n L_i^n(t) \equiv 1(t)$

Outline

Aitken's Algorithm

Lagrange Polynomials

The Vandermonde Approach

Limits of Lagrange Interpolation

Cubic Hermite Interpolation

Quintic Hermite Interpolation

Point-Normal Interpolation

Least Squares Approximation

Smoothing Equations

Designing with Bézier Curves

The Newton Form and Forward Differencing

The Vandermonde Approach

What is the monomial form

$$\mathbf{p}^n(t) = \sum_{j=0}^n \mathbf{a}_j t^j$$

of the interpolating polynomial?

The Vandermonde Approach (cont'd)

1.

$$\mathbf{p}^n(t_0) = \mathbf{p}_0 = \mathbf{a}_0 + \mathbf{a}_1 t_0 + \cdots + \mathbf{a}_n t_0^n$$

$$\mathbf{p}^n(t_1) = \mathbf{p}_1 = \mathbf{a}_0 + \mathbf{a}_1 t_1 + \cdots + \mathbf{a}_n t_1^n$$

$$\vdots$$

$$\mathbf{p}^n(t_n) = \mathbf{p}_n = \mathbf{a}_0 + \mathbf{a}_1 t_n + \cdots + \mathbf{a}_n t_n^n$$

$$2. \quad \begin{bmatrix} \mathbf{p}_0 \\ \mathbf{p}_1 \\ \vdots \\ \mathbf{p}_n \end{bmatrix} = \begin{bmatrix} 1 & t_0 & \cdots & t_0^n \\ 1 & t_1 & \cdots & t_1^n \\ \vdots & \vdots & \ddots & \vdots \\ 1 & t_n & \cdots & t_n^n \end{bmatrix} \begin{bmatrix} \mathbf{a}_0 \\ \mathbf{a}_1 \\ \vdots \\ \mathbf{a}_n \end{bmatrix} \rightarrow \mathbf{p} = T\mathbf{a}$$

3. If $\{t_j\}_{j=0}^n$ are distinct, $\det T \neq 0$. (Why?)

→ $\det T$ is the Vandermonde of the interpolation problem

4. Solution: $\mathbf{a} = T^{-1}\mathbf{p}$

Generalized Vandermonde

For some basis $\{F_j^n(t)\}_{j=0}^n$, (e.g. Bernstein basis) the interpolating polynomial is expressed as

$$\mathbf{p}^n(t) = \sum_{j=0}^n \mathbf{c}_j F_j^n(t)$$

and the coefficients $\{\mathbf{c}_j\}_{j=0}^n$ can be found as

$$\rightarrow \begin{bmatrix} \mathbf{c}_1 \\ \mathbf{c}_2 \\ \vdots \\ \mathbf{c}_n \end{bmatrix} = \begin{bmatrix} F_0^n(t_0) & F_1^n(t_0) & \cdots & F_n^n(t_0) \\ F_0^n(t_1) & F_1^n(t_1) & \cdots & F_n^n(t_1) \\ \vdots & \vdots & \ddots & \vdots \\ F_0^n(t_n) & F_1^n(t_n) & \cdots & F_n^n(t_n) \end{bmatrix}^{-1} \begin{bmatrix} \mathbf{p}_0 \\ \mathbf{p}_1 \\ \vdots \\ \mathbf{p}_n \end{bmatrix}$$

Outline

Aitken's Algorithm

Lagrange Polynomials

The Vandermonde Approach

Limits of Lagrange Interpolation

Cubic Hermite Interpolation

Quintic Hermite Interpolation

Point-Normal Interpolation

Least Squares Approximation

Smoothing Equations

Designing with Bézier Curves

The Newton Form and Forward Differencing

Limits of Lagrange Interpolation

- ▶ Small change of input results in large change in output
 - ▶ The process is *ill-conditioned*.
 - ▶ Polynomial interpolation is not *shape preserving*.
 - ▶ Runge phenomenon
- ▶ Expensive to solve
 - ▶ for construction (solving dense linear system)
 - ▶ for evaluation (for high n)
- ▶ Piecewise schemes are better! (e.g. spline interpolation)

Outline

Aitken's Algorithm

Lagrange Polynomials

The Vandermonde Approach

Limits of Lagrange Interpolation

Cubic Hermite Interpolation

Quintic Hermite Interpolation

Point-Normal Interpolation

Least Squares Approximation

Smoothing Equations

Designing with Bézier Curves

The Newton Form and Forward Differencing

Cubic Hermite Interpolation

Find the cubic polynomial that interpolates the following:

$$\mathbf{p}(0) = \mathbf{p}_0 \in \mathbb{E}^3$$

$$\dot{\mathbf{p}}(0) = \mathbf{m}_0 \in \mathbb{R}^3$$

$$\dot{\mathbf{p}}(1) = \mathbf{m}_1 \in \mathbb{R}^3$$

$$\mathbf{p}(1) = \mathbf{p}_1 \in \mathbb{E}^3,$$

Easier to work with Bézier form

$$\mathbf{p}(t) = \sum_{j=0}^3 \mathbf{b}_j B_j^3(t)$$

1. $\mathbf{b}_0 = \mathbf{p}_0$ and $\mathbf{b}_3 = \mathbf{p}_1$
2. $\dot{\mathbf{p}}(0) = 3\Delta\mathbf{b}_0 = \mathbf{m}_0$ and $\dot{\mathbf{p}}(1) = 3\Delta\mathbf{b}_2 = \mathbf{m}_1$
 $\rightarrow \mathbf{b}_1 = \mathbf{p}_0 + \frac{1}{3}\mathbf{m}_0$ and $\mathbf{b}_2 = \mathbf{b}_1 - \frac{1}{3}\mathbf{m}_1$

Cubic Hermite Interpolation: Cardinal Form

$$\mathbf{p}(t) = \mathbf{p}_0 B_0^3(t) + \left(\mathbf{p}_0 + \frac{1}{3} \mathbf{m}_0 \right) B_1^3(t) + \left(\mathbf{p}_1 - \frac{1}{3} \mathbf{m}_1 \right) B_2^3(t) + \mathbf{p}_1 B_3^3(t)$$

$$\rightarrow \mathbf{p}(t) = \mathbf{p}_0 H_0^3(t) + \mathbf{m}_0 H_1^3(t) + \mathbf{m}_1 H_2^3(t) + \mathbf{p}_1 H_3^3(t)$$

\rightarrow

$$H_0^3(t) = B_0^3(t) + B_1^3(t)$$

$$H_1^3(t) = \frac{1}{3} B_1^3(t)$$

$$H_2^3(t) = -\frac{1}{3} B_2^3(t)$$

$$H_3^3(t) = B_2^3(t) + B_3^3(t)$$

Cubic Hermite Interpolation: Cardinal Form

$$\begin{array}{cccc} H_0^3(0) = 1 & \frac{d}{dt}H_0^3(0) = 0 & \frac{d}{dt}H_0^3(1) = 0 & H_0^3(1) = 0 \\ H_1^3(0) = 0 & \frac{d}{dt}H_1^3(0) = 1 & \frac{d}{dt}H_1^3(1) = 0 & H_1^3(1) = 0 \\ H_2^3(0) = 0 & \frac{d}{dt}H_2^3(0) = 0 & \frac{d}{dt}H_2^3(1) = 1 & H_2^3(1) = 0 \\ H_3^3(0) = 0 & \frac{d}{dt}H_3^3(0) = 0 & \frac{d}{dt}H_3^3(1) = 0 & H_3^3(1) = 1 \end{array}$$

- ▶ $H_0^3(t) + H_3^3(t) \equiv 1(t)$ (Why?)
- ▶ Not invariant under affine domain transformations
- ▶ Not symmetric

Outline

Aitken's Algorithm

Lagrange Polynomials

The Vandermonde Approach

Limits of Lagrange Interpolation

Cubic Hermite Interpolation

Quintic Hermite Interpolation

Point-Normal Interpolation

Least Squares Approximation

Smoothing Equations

Designing with Bézier Curves

The Newton Form and Forward Differencing

Outline

Aitken's Algorithm

Lagrange Polynomials

The Vandermonde Approach

Limits of Lagrange Interpolation

Cubic Hermite Interpolation

Quintic Hermite Interpolation

Point-Normal Interpolation

Least Squares Approximation

Smoothing Equations

Designing with Bézier Curves

The Newton Form and Forward Differencing

Outline

Aitken's Algorithm

Lagrange Polynomials

The Vandermonde Approach

Limits of Lagrange Interpolation

Cubic Hermite Interpolation

Quintic Hermite Interpolation

Point-Normal Interpolation

Least Squares Approximation

Smoothing Equations

Designing with Bézier Curves

The Newton Form and Forward Differencing

Least Squares Approximation

- ▶ $P + 1$ number of data points can be interpolated by a polynomial of degree n if $d \geq P$.
- ▶ What if $n < P$? \rightarrow approximation required
- ▶ How to measure the closeness of the approximating curve and the data points?
- ▶ *least squares approximation*: To find a polynomial curve $\mathbf{x}(t)$ that minimizes

$$f(\mathbf{c}_0, \dots, \mathbf{c}_n) = \sum_{j=0}^P \|\mathbf{x}(t_j) - \mathbf{p}_j\|^2.$$

Least Squares Approximation (cont'd)

1. The approximating curve is expressed as

$$\mathbf{x}(t) = c_0 C_0^n(t) + \cdots + c_n C_n^n(t)$$

for some basis $\{C_j^n\}_{j=0}^n$.

2. Assuming $\mathbf{x}(t)$ interpolates all the points $\{\mathbf{p}_j\}_{j=0}^P$, we get a linear system

$$\begin{bmatrix} C_0^n(t_0) & \cdots & C_n^n(t_0) \\ \vdots & \ddots & \vdots \\ C_0^n(t_P) & \cdots & C_n^n(t_P) \end{bmatrix} \begin{bmatrix} \mathbf{c}_0 \\ \vdots \\ \mathbf{c}_n \end{bmatrix} = \begin{bmatrix} \mathbf{p}_0 \\ \vdots \\ \mathbf{p}_P \end{bmatrix} \quad \rightarrow \quad M\mathbf{C} = \mathbf{P}$$

where

$$M \in \mathbb{R}^{(P+1) \times (n+1)}, \mathbf{C} \in \mathbb{R}^{(n+1) \times 3}, \mathbf{P} \in \mathbb{R}^{(P+1) \times 3}, \quad P > n$$

- Linear systems for each component
- underdetermined (no solution in general)

Least Squares Approximation (cont'd)

3. Can be converted to a solvable linear system by multiplying M^T :

$$M^T M \mathbf{C} = M^T \mathbf{P}$$

- ▶ $M^T M$ is always invertible. (Why?)
- ▶ The solution minimizes

$$f(\mathbf{c}_0, \dots, \mathbf{c}_n) = \sum_{j=0}^P \|\mathbf{x}(t_j) - \mathbf{p}_j\|^2.$$

(Why?)

The solution satisfies

$$\frac{\partial f}{\partial \mathbf{c}_k^d} = 0, \quad k = 0, \dots, n, d = 1, 2, 3.$$

Outline

Aitken's Algorithm

Lagrange Polynomials

The Vandermonde Approach

Limits of Lagrange Interpolation

Cubic Hermite Interpolation

Quintic Hermite Interpolation

Point-Normal Interpolation

Least Squares Approximation

Smoothing Equations

Designing with Bézier Curves

The Newton Form and Forward Differencing

Smoothing Equations

- ▶ Minimizing the least square error does not guarantee a good curve. (Fig 7.8)
- ▶ We want a curve that does not “wobble” much. (How?)
- ▶ If the control polygon behaves nicely, then so does the curve.
- ▶ How to minimize wiggling (of the polygon)? → minimize the 2nd differences $\Delta^2 \mathbf{b}_i$

$$\mathbf{b}_0 - 2\mathbf{b}_1 + \mathbf{b}_2 = 0$$

$$\vdots$$

$$\rightarrow S\mathbf{B} = \mathbf{0}$$

$$\mathbf{b}_{n-2} - 2\mathbf{b}_{n-1} + \mathbf{b}_n = 0$$

Smoothing Equations (cont'd)

- ▶ Combined with the least squares problem,

$$\begin{bmatrix} M \\ S \end{bmatrix} \mathbf{B} = \begin{bmatrix} \mathbf{P} \\ \mathbf{0} \end{bmatrix}$$

- ▶ Balancing the error and shape

$$\begin{bmatrix} (1 - \alpha)M \\ \alpha S \end{bmatrix} \mathbf{B} = \begin{bmatrix} (1 - \alpha)\mathbf{P} \\ \mathbf{0} \end{bmatrix} \quad 0 \leq \alpha \leq 1$$

- ▶ Larger α for noisier data

Outline

Aitken's Algorithm

Lagrange Polynomials

The Vandermonde Approach

Limits of Lagrange Interpolation

Cubic Hermite Interpolation

Quintic Hermite Interpolation

Point-Normal Interpolation

Least Squares Approximation

Smoothing Equations

Designing with Bézier Curves

The Newton Form and Forward Differencing

Designing with Bézier Curves

- ▶ “What if we want to move a point on the curve to change the shape?”
- ▶ Given a Bézier curve

$$\mathbf{x}(t) = \sum_{i=0}^n \mathbf{b}_i B_i^n(t),$$

if we change a point $\mathbf{x}(\hat{t})$ to \mathbf{y} , how should we change $\{\mathbf{b}_i\}$ to $\{\hat{\mathbf{b}}_i\}$ such that the curve passes through \mathbf{y} at $t = \hat{t}$?

$$\mathbf{y} = \sum_{i=0}^n \hat{\mathbf{b}}_i B_i^n(\hat{t})$$

- ▶ Assume \mathbf{b}_0 and \mathbf{b}_n are unchanged.
- ▶ Underdetermined: 1 equation and $n - 1$ unknowns

Designing with Bézier Curves (cont'd)

1.

$$\begin{bmatrix} B_1^n(\hat{t}) & \cdots & B_{n-1}^n(\hat{t}) \end{bmatrix} \begin{bmatrix} \hat{\mathbf{b}}_1 \\ \vdots \\ \hat{\mathbf{b}}_{n-1} \end{bmatrix} = \mathbf{y} - \mathbf{b}_0 B_0^n(\hat{t}) - \mathbf{b}_n B_n^n(\hat{t}) =: \mathbf{z}$$
$$\rightarrow A\hat{\mathbf{B}} = \mathbf{z}$$

2. Assume all the points $\mathbf{b}_1, \dots, \mathbf{b}_{n-1}$ move along (unknown) \mathbf{c} direction:

$$\hat{\mathbf{B}} = \mathbf{B} + A^T \mathbf{c} \rightarrow A\hat{\mathbf{B}} = A\mathbf{B} + AA^T \mathbf{c} \rightarrow \mathbf{z} - A\mathbf{B} = AA^T \mathbf{c}$$

3. Since AA^T is a scalar, with $a := AA^T$

$$\mathbf{c} = \frac{1}{a}(\mathbf{y} - \mathbf{x}) \rightarrow \hat{\mathbf{B}} = \mathbf{B} + \frac{1}{a}A^T(\mathbf{y} - \mathbf{x}).$$

Outline

Aitken's Algorithm

Lagrange Polynomials

The Vandermonde Approach

Limits of Lagrange Interpolation

Cubic Hermite Interpolation

Quintic Hermite Interpolation

Point-Normal Interpolation

Least Squares Approximation

Smoothing Equations

Designing with Bézier Curves

The Newton Form and Forward Differencing

The Newton Form of Interpolating Polynomial

- ▶ For equally spaced parameter intervals: $t_{i+1} = t_i + h$
- ▶ For cubic case,

$$\begin{aligned}\mathbf{p}(t) = & \mathbf{p}_0 + \frac{1}{h}(t - t_0)\Delta\mathbf{p}_0 + \frac{1}{2!h^2}(t - t_0)(t - t_1)\Delta^2\mathbf{p}_0 \\ & + \frac{1}{3!h^3}(t - t_0)(t - t_1)(t - t_2)\Delta^3\mathbf{p}_0\end{aligned}$$

where

$$\Delta^i \mathbf{p}_j := \Delta^{i-1} \mathbf{p}_{j+1} - \Delta^{i-1} \mathbf{p}_j \quad \text{and} \quad \Delta^0 \mathbf{p}_j = \mathbf{p}_j.$$

- ▶ With $g := 1/h$,

\mathbf{p}_0			
\mathbf{p}_1	$g\Delta\mathbf{p}_0$		
\mathbf{p}_2	$g\Delta\mathbf{p}_1$	$g^2\Delta^2\mathbf{p}_0$	
\mathbf{p}_3	$g\Delta\mathbf{p}_2$	$g^2\Delta^2\mathbf{p}_1$	$g^3\Delta^3\mathbf{p}_0$

Fast Evaluation for Plotting

- ▶ Efficient way to evaluate $\mathbf{p}(t_2 + h), \mathbf{p}(t_2 + 2h), \dots$

$$\begin{array}{cccc} \mathbf{p}_0 & & & \\ \mathbf{p}_1 & g\Delta\mathbf{p}_0 & & \\ \mathbf{p}_2 & g\Delta\mathbf{p}_1 & g^2\Delta^2\mathbf{p}_0 & \\ \mathbf{p}_3 & g\Delta\mathbf{p}_2 & g^2\Delta^2\mathbf{p}_1 & g^3\Delta^3\mathbf{p}_0 \\ \mathbf{p}_4 & g\Delta\mathbf{p}_3 & g^2\Delta^2\mathbf{p}_2 & g^3\Delta^3\mathbf{p}_1 \\ \mathbf{p}_5 & g\Delta\mathbf{p}_4 & g^2\Delta^2\mathbf{p}_3 & g^3\Delta^3\mathbf{p}_2 \\ \vdots & \vdots & \vdots & \vdots \end{array}$$

- ▶ $\{g\Delta^3\mathbf{p}_j\}$ are all equal. (Why?)
- ▶ From right to left

$$\begin{aligned} g^2\Delta^2\mathbf{p}_2 &= g^3\Delta^3\mathbf{p}_1 + g^2\Delta^2\mathbf{p}_1 \rightarrow g\Delta\mathbf{p}_3 = g^2\Delta^2\mathbf{p}_2 + g\Delta\mathbf{p}_2 \\ &\rightarrow \mathbf{p}_4 = g\Delta\mathbf{p}_3 + \mathbf{p}_3 \end{aligned}$$

- ▶ In general, with $\mathbf{q}_j^i := g^i\Delta^i\mathbf{p}_j$,

$$\mathbf{q}_j^i = \mathbf{q}_{j-1}^{i+1} + \mathbf{q}_{j-1}^i, \quad i = 2, 1, 0$$

and $\mathbf{p}_j = \mathbf{q}_j^0$.

Fast for Plotting (cont'd)

- ▶ No multiplication involved! (after start-up phase) → extremely fast
- ▶ Given a polynomial in cubic Bézier form,
 1. Compute initial four points.
 2. Evaluate sequence of points and plot using line segments.
- ▶ Roundoff error is accumulated → The curve may deviates more and more...