

Topics in Computer Graphics

Chap 3: Linear Interpolation

fall, 2011

University of Seoul
School of Computer Science
Minho Kim

Table of contents

Linear Interpolation

Piecewise Linear Interpolation

Menelaos' Theorem

Blossoms

barycentric Coordinates in the Plane

Tessellations

Linear Interpolation

Let $\mathbf{a}, \mathbf{b} \in \mathbb{E}^3$. The set of all points $\mathbf{x} \in \mathbb{E}^3$ of the form

$$\mathbf{x} = \mathbf{x}(t) = (1 - t)\mathbf{a} + t\mathbf{b}, \quad t \in \mathbb{R}$$

is called the *straight line* through \mathbf{a} and \mathbf{b} .

- ▶ For $t = 0$, $\mathbf{x}(0) = \mathbf{a}$: the line passes through \mathbf{a} .
- ▶ For $t = 1$, $\mathbf{x}(1) = \mathbf{b}$: the line passes through \mathbf{b} .
- ▶ For $0 \leq t \leq 1$, the point \mathbf{x} is between \mathbf{a} and \mathbf{b} .
- ▶ For $t < 0$ or $t > 1$, the point is outside.
- ▶ \mathbf{x} is represented as a barycentric combination of two points in \mathbb{E}^3 .
 - The three points $\mathbf{a}, \mathbf{x}, \mathbf{b}$ in \mathbb{E}^3 are an affine map of the three 1D points $0, t, 1$.
 - Linear interpolation is an affine map of the real line onto a straight line in \mathbb{E}^3 .

Linear Interpolation (cont'd)

- ▶ Linear interpolation is affinely invariant.

$$\Phi \mathbf{x} = \Phi ((1 - t)\mathbf{a} + t\mathbf{b}) = (1 - t)\Phi \mathbf{a} + t\Phi \mathbf{b}$$

- ▶ Can be applied to vectors as well : The vector $\vec{v} := d - c \in \mathbb{R}$ is mapped to the vector $\mathbf{l}(\vec{v})\mathbf{l}(d) - \mathbf{l}(c) \in \mathbb{R}^3$ by the linear interpolation \mathbf{l} .

Linear Interpolation and Barycentric Combination

- ▶ For any three conlinear points $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{E}^3$, the barycentric coordinates of \mathbf{b} w.r.t. \mathbf{a} and \mathbf{c} is

$$\mathbf{b} = \alpha \mathbf{a} + \beta \mathbf{c}, \quad \alpha = \frac{\text{vol}_1(\mathbf{b}, \mathbf{c})}{\text{vol}_1(\mathbf{a}, \mathbf{c})}, \beta = \frac{\text{vol}_1(\mathbf{a}, \mathbf{b})}{\text{vol}_1(\mathbf{a}, \mathbf{c})}.$$

- ▶ vol_1 : signed distance between two points
- ▶ $\text{ratio}(\mathbf{a}, \mathbf{b}, \mathbf{c}) = \frac{\text{vol}_1(\mathbf{a}, \mathbf{b})}{\text{vol}_1(\mathbf{b}, \mathbf{c})} = \frac{\beta}{\alpha}$
- ▶ The barycentric coordinates of a point do not change under affine maps.

$$\text{ratio}(\Phi \mathbf{a}, \Phi \mathbf{b}, \Phi \mathbf{c}) = \frac{\beta}{\alpha}$$

→ Affine maps are ratio preserving.

→ *Every map that takes straight lines to straight lines and its ratio preserving is an affine map.*

Affine Domain Transformation

The straight line

$$\mathbf{x}(t) = (1 - t)\mathbf{a} + t\mathbf{b}$$

for $t \in [0, 1]$ is the same as the straight line

$$\mathbf{x}(u) = \frac{b - u}{b - a}\mathbf{a} + \frac{u - a}{b - a}\mathbf{b}$$

for $u \in [a, b]$ with $t = (u - a)/(b - a)$.

→ Linear interpolation is invariant under affine domain transformation.

- ▶ Affin domain transformation: An affine map of the real line onto itself.

Piecewise Linear Interpolation

- ▶ Let $\mathbf{b}_0, \dots, \mathbf{b}_n \in \mathbb{E}^3$ form a polygon \mathbf{B} .
→ \mathbf{B} is the *piecewise linear interpolant* \mathcal{PL} to the points \mathbf{b}_i .
- ▶ If the points \mathbf{b}_I lie on a curve \mathbf{c}
→ \mathbf{B} is a piecewise linear interpolant to \mathbf{c} :

$$\mathbf{B} = \mathcal{PL}\mathbf{c}.$$

- ▶ Piecewise linear interpolation is affinely invariant.:

$$\mathcal{PL}\Phi\mathbf{c} = \Phi\mathcal{PL}\mathbf{c}.$$

- ▶ *Varoatopm diminishing property:*

$$\text{cross}(\mathcal{PL}\mathbf{c}) \leq \text{cross } \mathbf{c}.$$

Menelaos' Theorem

Let

$$\mathbf{b}[0, t] = (1 - t)\mathbf{b}_0 + t\mathbf{b}_1$$

$$\mathbf{b}[s, 0] = (1 - s)\mathbf{b}_0 + s\mathbf{b}_1$$

$$\mathbf{b}[1, t] = (1 - t)\mathbf{b}_1 + t\mathbf{b}_2$$

$$\mathbf{b}[s, 1] = (1 - s)\mathbf{b}_1 + s\mathbf{b}_2$$

and

$$\mathbf{b}[s, t] = (1 - t)\mathbf{b}[s, 0] + t\mathbf{b}[s, 1]$$

$$\mathbf{b}[t, s] = (1 - s)\mathbf{b}[0, t] + s\mathbf{b}[t, 1].$$

Then

$$\mathbf{b}[s, t] = \mathbf{b}[t, s].$$

- [Menelaus' Theorem @Wikipedia](#)

Blossoms

A blossom is an n -variate function $\mathbf{b}[t_1, \dots, t_n]$ from \mathbb{R}^n into \mathbb{E}^2 or \mathbb{E}^3 satisfying the following three properties:

- ▶ Symmetry:

$$\mathbf{b}[t_1, \dots, t_n] = \mathbf{b}[\pi(t_1, \dots, t_n)]$$

where $\pi(t_1, \dots, t_n)$ denotes a permutation of the arguments t_1, \dots, t_n .

→ The order of the arguments does not matter

→ Menelaos' theorem

- ▶ Multiaffinity

$$\mathbf{b}[(\alpha r + \beta s), *] = \alpha \mathbf{b}[r, *] + \beta \mathbf{b}[s, *], \quad \alpha + \beta = 1$$

→ Affine w.r.t. *any* argument.

- ▶ Diagonality

$$\mathbf{b}[t, \dots, t] = \mathbf{b}[t^{<n>}]$$

When all the n arguments are the same, it traces out a polynomial curve of degree n .

Blossoms with Vector Argument

With $\vec{h} := b - a$,

$$\mathbf{b}[\vec{h}, *] = \mathbf{b}[b - a, *] = \mathbf{b}[b, *] - \mathbf{b}[a, *]$$

→ If (at least) one of the blossom arguments is a vector, then the blossom value is a vector.

Leibniz Formula

$$\mathbf{b}[(\alpha r + \beta s)^{<n>}] = \sum_{i=0}^n \binom{n}{i} \alpha^i \beta^{n-i} \mathbf{b}[r^{<i>}, s^{<n-i>}]$$

Alternative formula

$$\mathbf{b}[(\alpha r + \beta s)^{<n>}] = \sum_{\substack{i+j=n \\ i,j \geq 0}} \binom{n}{i,j} \alpha^i \beta^{n-i} \mathbf{b}[r^{<i>}, s^{<j>}]$$

where

$$\binom{n}{i,j} := \frac{n!}{i!j!}.$$

Barycentric Coordinates in the Plane

Considering a triangle with vertices $\mathbf{a}, \mathbf{b}, \mathbf{c}$ in \mathbb{E}^2 , *any* point $\mathbf{p} \in \mathbb{E}^2$ can be represented as a barycentric combination of $\mathbf{a}, \mathbf{b}, \mathbf{c}$:

$$\mathbf{p} = u\mathbf{a} + v\mathbf{b} + w\mathbf{c}, \quad \text{where } u + v + w = 1.$$

- ▶ Is (u, v, w) is unique? \rightarrow The coefficients $\mathbf{u} := (u, v, w)$ is the *barycentric coordinates* of \mathbf{p} w.r.t. $\mathbf{a}, \mathbf{b}, \mathbf{c}$.
- ▶ Applying the Cramer's rule,

$$u = \frac{\text{area}(\mathbf{p}, \mathbf{b}, \mathbf{c})}{\text{area}(\mathbf{a}, \mathbf{b}, \mathbf{c})}, \quad v = \frac{\text{area}(\mathbf{a}, \mathbf{p}, \mathbf{c})}{\text{area}(\mathbf{a}, \mathbf{b}, \mathbf{c})}, \quad w = \frac{\text{area}(\mathbf{a}, \mathbf{b}, \mathbf{p})}{\text{area}(\mathbf{a}, \mathbf{b}, \mathbf{c})}.$$

\rightarrow Requires $\text{area}(\mathbf{a}, \mathbf{b}, \mathbf{c}) \neq 0$.

- ▶ Barycentric coordinates are *affinely invariant*.
- ▶ Ceva's theorem (Fig. 3.5)
- ▶ Location of \mathbf{p} according to the signs of its barycentric coordinates \rightarrow Fig. 3.6

Bivariate Linear Interpolation

Given three points $\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3 \in \mathbb{E}^3$, any point of the form

$$\mathbf{p} = \mathbf{p}(\mathbf{u}) = \mathbf{p}(u, v, w) = u\mathbf{p}_1 + v\mathbf{p}_2 + w\mathbf{p}_3, \quad u + v + w = 1$$

lies in the plane spanned by $\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3$.

- ▶ Can be generalized to higher dimensions, e.g., a barycentric coordinates w.r.t. a tetrahedron.

Tessellations

- ▶ Bivariate piecewise linear interpolation requires **triangulation** of a plane.
→ related to the concept of **tessellation**
- ▶ **Dirichlet tessellation** (a.k.a. Voronoi diagram)
“...we associate with each point \mathbf{p}_k a tile T_k consisting of all points \mathbf{p} that are closer to \mathbf{p}_k than to any other point \mathbf{p}_i .
The collection of all these tiles is called the Dirichlet tessellation of the given point set.”

Triangulations

- ▶ A triangulation \mathcal{T} of a set of 2D points $\{\mathbf{p}_i\}$ is a collection of triangles such that
 - ▶ The vertices of the triangles consist of the \mathbf{p}_i .
 - ▶ The interiors of any two triangles do not intersect.
 - ▶ If two triangles are not disjoint, then they share either a vertex or an edge.
- ▶ Delaunay triangulation
 - ▶ Dual of the Voronoi diagram
 - ▶ Satisfies the maxmin criterion.
 - ▶ Given a point set, is its Delaunay triangulation unique?
- ▶ Piecewise linear interpolation on a plane