

CCC '20 J2 - Epidemiology

Time Limit: 1.0s **Memory Limit:** 512M
Python 3: 3.0s

Canadian Computing Competition: 2020 Stage 1, Junior #2

People who study epidemiology use models to analyze the spread of disease. In this problem, we use a simple model.

When a person has a disease, they infect exactly R other people but only on the very next day. No person is infected more than once. We want to determine when a total of more than P people have had the disease.

(This problem was designed before the current coronavirus outbreak, and we acknowledge the distress currently being experienced by many people worldwide because of this and other diseases. We hope that including this problem at this time highlights the important roles that computer science and mathematics play in solving real-world problems.)

Input Specification

There are three lines of input. Each line contains one positive integer. The first line contains the value of P . The second line contains N , the number of people who have the disease on Day 0. The third line contains the value of R . Assume that $P \leq 10^7$ and $N \leq P$ and $R \leq 10$.

Output Specification

Output the number of the first day on which the total number of people who have had the disease is greater than P .

Sample Input 1

```
750
1
5
```

Output for Sample Input 1

```
4
```

Explanation of Output for Sample Input 1

The 1 person on Day 0 with the disease infects 5 people on Day 1. On Day 2, exactly 25 other people are infected. On Day 3, exactly 125 other people are infected. A total of $1 + 5 + 25 + 125 + 625 = 781$ people have had the disease by the end of Day 4 and $781 > 750$.

Sample Input 2

10
2
1

Output for Sample Input 2

5

Explanation of Output for Sample Input 2

There are 2 people on Day 0 with the disease. On each other day, exactly 2 other people are infected. By the end of Day 4, a total of exactly 10 people have had the disease and by the end of Day 5, more than 10 people have had the disease.