

# Hydrodynamic processes in protoplanetary disks


Min-Kai Lin

Steward Theory Fellow  
University of Arizona

June 14 2016

# A new era for planet formation


Planets form in protoplanetary accretion disks around young stars



(ALMA Partnership, 2015, HL Tau)


# A new era for planet formation

Planets form in protoplanetary accretion disks around young stars




(TW Hydrae, Andrews et al., 2016)

# Asymmetric transition disks



(HD142527, Casassus et al., 2015)

# Directly imaged planets



(51 Eridani, Macintosh et al., 2015)

# Planet formation through accretion disk theory

## Fundamental gas dynamics of protoplanetary accretion disks

- How do disks transport angular momentum and accrete?
  - ▶ Turbulent transport?
  - ▶ Spiral arms from self-gravity?
  - ▶ Magnetic winds?

# Planet formation through accretion disk theory

## Fundamental gas dynamics of protoplanetary accretion disks

- How do disks transport angular momentum and accrete?
  - ▶ Turbulent transport?
  - ▶ Spiral arms from self-gravity?
  - ▶ Magnetic winds?
- What are the sources of turbulence in disks?
  - ▶ Hydrodynamic, gravitational, or magneto-hydrodynamic?
  - ▶ Where and when do these operate?

# Planet formation through accretion disk theory

## Fundamental gas dynamics of protoplanetary accretion disks

- How do disks transport angular momentum and accrete?
  - ▶ Turbulent transport?
  - ▶ Spiral arms from self-gravity?
  - ▶ Magnetic winds?
- What are the sources of turbulence in disks?
  - ▶ Hydrodynamic, gravitational, or magneto-hydrodynamic?
  - ▶ Where and when do these operate?
- Origin of large-scale structures? Rings, gaps, asymmetries/vortices, spirals
  - ▶ Planet induced?
  - ▶ Fluid instabilities?

# Planet formation through accretion disk theory

## Fundamental gas dynamics of protoplanetary accretion disks

- How do disks transport angular momentum and accrete?
  - ▶ Turbulent transport?
  - ▶ Spiral arms from self-gravity?
  - ▶ Magnetic winds?
- What are the sources of turbulence in disks?
  - ▶ Hydrodynamic, gravitational, or magneto-hydrodynamic?
  - ▶ Where and when do these operate?
- Origin of large-scale structures? Rings, gaps, asymmetries/vortices, spirals
  - ▶ Planet induced?
  - ▶ Fluid instabilities?
- How do large-scale structures affect planet formation?
  - ▶ Dust-trapping mechanisms (enhance planetesimal formation)?
  - ▶ Dynamical interaction with planets?

# Hydrodynamic activities in all stages of PPD evolution


Protostellar disks: gravitational instabilities

disk evolution

MMSN disk: vertical shear instability

planet formation

Planet-induced vortices/dust-traps



(Lin, Fargo sims.)

- Going beyond Toomre and Lin-Shu analyses (Lin & Kratter, 2016)

# Hydrodynamic activities in all stages of PPD evolution


Protostellar disks: gravitational instabilities

disk evolution

MMSN disk: vertical shear instability

planet formation

Planet-induced vortices/dust-traps



(Nelson et al., 2013)

- Does it occur in realistic PPDs?  
(Lin & Youdin, 2015)

# Hydrodynamic activities in all stages of PPD evolution


Protostellar disks: gravitational instabilities

disk evolution

MMSN disk: vertical shear instability

planet formation


Planet-induced vortices/dust-traps




(Lin, 2012b)

- 3D effects on vortex evolution (Lin, in prep.)

# Wide orbit planets/brown dwarfs & disk instability theory



(HR8799 Marois et al., 2010)



(Lin, Fargo sims., log dens.)

- Young, massive PPDs can fragment by self-gravity

# Fragmentation conditions

- Massive disk

$$Q \equiv \frac{c_s \Omega}{\pi G \Sigma} \lesssim 2 \text{ or } M_{\text{disk}} \gtrsim 0.1 M_*$$

- Fast cooling

$$t_{\text{cool}} \Omega \lesssim 3$$

The cooling criterion is empirical!

**WARNING**

Critical cooling depends on the numerical simulation!

(resolution, 2D/3D, local/global, particle-based or grid-based simulations)

# Fragmentation conditions

- Massive disk

$$Q \equiv \frac{c_s \Omega}{\pi G \Sigma} \lesssim 2 \text{ or } M_{\text{disk}} \gtrsim 0.1 M_*$$

- Fast cooling

$$t_{\text{cool}} \Omega \lesssim 3$$

The cooling criterion is empirical!

**WARNING**

Critical cooling depends on the numerical simulation!

(resolution, 2D/3D, local/global, particle-based or grid-based simulations)

Motivation 1:

Assess disk fragmentation without input from hydrodynamic simulations

# Beyond classical gravitational instability

Modern simulations (c. 2010)

- Cooling physics, e.g.

$$\frac{\partial E}{\partial t} = -\frac{E}{t_{\text{cool}}}$$

- Turbulent/viscous, e.g.

$$\nu = \alpha \frac{c_s^2}{\Omega}$$

Analytic toolbox (c. 1960)

- Isothermal/adiabatic (no cooling)

$$\frac{\partial E}{\partial t} = 0$$

- Laminar (inviscid)

$$\nu = 0$$

Lin-Shu dispersion relation, Toomre  $Q$

$$\omega^2 = \kappa^2 - 2\pi G \Sigma |k| + c_s^2 k^2$$

$$Q \equiv \frac{c_s \kappa}{\pi G \Sigma}$$

# Beyond classical gravitational instability

Modern simulations (c. 2010)

- Cooling physics, e.g.

$$\frac{\partial E}{\partial t} = -\frac{E}{t_{\text{cool}}}$$

- Turbulent/viscous, e.g.

$$\nu = \alpha \frac{c_s^2}{\Omega}$$

Analytic toolbox (c. 1960)

- Isothermal/adiabatic (no cooling)

$$\frac{\partial E}{\partial t} = 0$$

- Laminar (inviscid)

$$\nu = 0$$

- $\omega^2(k; Q) = \kappa^2 - 2\pi G \Sigma |k| + c_s^2 k^2$


## Motivation 2:

Generalize analytic treatment of GI to include cooling, irradiation and viscosity

$$\omega = \omega(k; Q, t_{\text{cool}}, \alpha)$$

Cooling/viscosity  $\rightarrow$  can be unstable even for  $Q > 1$  (cf.  $Q < 1$  for classic GI)


# Cooling-driven gravitational instability



(Lin & Kratter, 2016)  $T_{\text{irr}}$  : irradiation temp.

# Understanding simulations


Cooling timescale to remove pressure over a lengthscale  $\sim H$



Simulations: Gammie (2001); Rice et al. (2005, 2011); Paardekooper (2012)

## Viscous gravitational instability


- Viscosity/friction can remove rotational stabilization  
(Lynden-Bell & Pringle, 1974)



(Lin & Kratter, 2016)

## Putting it all together

- Input physical disk model **with cooling and viscosity** — get **growth timescales**




(Lin & Kratter, 2016)

- High  $\dot{M}$  disk fragments  $\gtrsim 60\text{AU}$ , growth times  $\sim$  one orbit

# What's next for disk GI theory?

- Global effects with cooling and viscosity
  - ▶ Mass infall
  - ▶ Disks with radial structure
  - ▶ Large-scale spiral instabilities
- Magnetic effects : good or bad for stability?
  - ▶ Extend Lin (2014) to include cooling/viscosity
- Direct simulations : include actual turbulence (cf. viscosity)
  - ▶ Hydrodynamic turbulence due to VSI? Next topic!

## Astrophysical disks have vertical shear



cylind. rad. from star,  $r$

$$\frac{\partial \Omega}{\partial z} \neq 0$$

(Because  $\nabla P \times \nabla \rho \neq 0$ )


- Vertically isothermal thin-disk with  $T \propto r^q$ ,

$$r \frac{\partial \Omega}{\partial z} \simeq \left( \frac{qz}{2H} \right) \times \frac{H}{r} \Omega_{\text{Kep}}$$

- $H/r \sim 0.05$  in PPDs

# Vertical shear instability

$\partial_z \Omega \neq 0 \Rightarrow$  free energy  $\rightarrow$  instability?



- Change in kinetic energy:


$$\Delta E \sim l_r^2 \left( \Omega^2 + \frac{l_z}{l_r} \cdot r \frac{\partial \Omega^2}{\partial z} \right)$$

- Vertical shear is weak, **BUT**

$\Delta E < 0$  if  $|l_z| \gg |l_r|$   
 $\Rightarrow$  **INSTABILITY**

- Energy released for vertically elongated disturbances.

# VSI needs to fight buoyancy in real disks



- Vertical shear is weak,  $r\partial_z\ln\Omega \sim O(h) \ll 1$  (so need  $l_z/l_r \gg 1$ )
- Vertical buoyancy is strong,  $N_z/\Omega \sim O(1)$

# Ultra-fast cooling can overcome buoyancy forces

(Lin & Youdin, 2015) : quasi-global analyses of the VSI

- Including energy equation with finite cooling timescale  $t_{\text{cool}}$

For  $T \propto r^q$  PPD, find that VSI requires:


$$t_{\text{cool}}\Omega_K < \frac{h|q|}{\gamma - 1} \ll 1$$

- $h|q|$ : vertical shear ( $h \equiv H/r \ll 1$ ) — destabilizing
- $\gamma - 1$ : vertical buoyancy — stabilizing
- As seen in high res. numerical simulations  
e.g. Nelson et al. (2013) find VSI only for  $t_{\text{cool}}\Omega_K \lesssim 0.06$

# Do protoplanetary disks actually develop VSI?

Cooling via dust-opacity ( $\propto T^2$ ) in the Minimum Mass Solar Nebula

$$\beta_{\text{crit}} \equiv \frac{h|q|}{\gamma - 1}$$



# What's next for VSI theory?

## Vertical shear in dusty disks

$$r \frac{\partial \Omega^2}{\partial z} = \frac{c_s^2}{\left[1 + \left(\frac{\rho_{\text{dust}}}{\rho_{\text{gas}}}\right)\right]^2} \left[ \frac{\partial \ln \rho_{\text{gas}}}{\partial z} \frac{\partial}{\partial r} \left( \frac{\rho_{\text{dust}}}{\rho_{\text{gas}}} \right) - \frac{\partial \ln \rho_{\text{gas}}}{\partial r} \frac{\partial}{\partial z} \left( \frac{\rho_{\text{dust}}}{\rho_{\text{gas}}} \right) \right]$$

Is there a 'dusty VSI'?

# What's next for VSI theory?

## Vertical shear in dusty disks

$$r \frac{\partial \Omega^2}{\partial z} = \frac{c_s^2}{\left[1 + \left(\frac{\rho_{\text{dust}}}{\rho_{\text{gas}}}\right)\right]^2} \left[ \frac{\partial \ln \rho_{\text{gas}}}{\partial z} \frac{\partial}{\partial r} \left( \frac{\rho_{\text{dust}}}{\rho_{\text{gas}}} \right) - \frac{\partial \ln \rho_{\text{gas}}}{\partial r} \frac{\partial}{\partial z} \left( \frac{\rho_{\text{dust}}}{\rho_{\text{gas}}} \right) \right]$$

Is there a 'dusty VSI'?

Not for perfectly coupled dust particles in an isothermal disk

# Isothermal, well-coupled dusty gas $\equiv$ adiabatic pure gas

## Isothermal dusty gas

- Small dust particles follow gas

$$\frac{D}{Dt} \left( \frac{\rho_{\text{dust}}}{\rho_{\text{gas}}} \right) \simeq 0$$

- RHS  $\equiv 0$  for  $t_{\text{stop}} \rightarrow 0$

## Textbook hydrodynamics

- Entropy follows gas

$$\frac{DS}{Dt} \simeq 0$$

- RHS  $\equiv 0$  for  $t_{\text{cool}} \rightarrow \infty$

Exploit this analogy to

- Provide **thermodynamical interpretations** of known results in dusty fluids
  - ▶ E.g. streaming instabilities
- Explore **dusty analogs** of known phenomena in standard hydrodynamics
  - ▶ E.g. dusty Solberg-Hoiland criterion  $\Rightarrow$  a limit on dust edges
- Study **asymmetric dust-trapping**, next topic!

# Isothermal, well-coupled dusty gas $\equiv$ adiabatic pure gas

## Isothermal dusty gas

- Small dust particles follow gas

$$\frac{D}{Dt} \left( \frac{\rho_{\text{dust}}}{\rho_{\text{gas}}} \right) \simeq 0$$


- RHS  $\equiv 0$  for  $t_{\text{stop}} \rightarrow 0$

## Textbook hydrodynamics

- Entropy follows gas

$$\frac{DS}{Dt} \simeq 0$$

- RHS  $\equiv 0$  for  $t_{\text{cool}} \rightarrow \infty$



- Dusty-gas sims. from Lorén-Aguilar & Bate (2015)
- What is this?

# Isothermal, well-coupled dusty gas $\equiv$ adiabatic pure gas

## Isothermal dusty gas

- Small dust particles follow gas

$$\frac{D}{Dt} \left( \frac{\rho_{\text{dust}}}{\rho_{\text{gas}}} \right) \simeq 0$$

- RHS  $\equiv 0$  for  $t_{\text{stop}} \rightarrow 0$

## Textbook hydrodynamics

- Entropy follows gas


$$\frac{DS}{Dt} \simeq 0$$

- RHS  $\equiv 0$  for  $t_{\text{cool}} \rightarrow \infty$

Exploit this analogy to

- Provide **thermodynamical interpretations** of known results in dusty fluids
  - ▶ E.g. streaming instabilities
- Explore **dusty analogs** of known phenomena in standard hydrodynamics
  - ▶ E.g. dusty Solberg-Hoiland criterion  $\Rightarrow$  a limit on dust edges
- Study **asymmetric dust-trapping**, next topic!

# Transition disk asymmetries: vortices?




(Oph IRS 48, van der Marel et al., 2013)

# Transition disk asymmetries: vortices?




Jupiter's Great Red Spot

# Transition disk asymmetries: vortices?




(Oph IRS 48, van der Marel et al., 2013)

## Dust-trapping at pressure maxima



- Drag forces cause dust to accumulate at pressure bumps

# Dust distribution in disk vortices




$$\rho_d(a) \propto \exp\left(-\frac{a^2}{2H_v^2}\right),$$

- $a$ : distance from the vortex center  
(Lyra & Lin, 2013)

$$H_v(\chi, \delta, \text{St}) = \frac{H_g}{f(\chi)} \sqrt{\frac{\delta}{\delta + \text{St}}}.$$

- $\chi$ : vortex aspect-ratio
- $\delta$ : turbulence in the vortex
- St: Stokes number (dust-gas friction)
- $H_g$ : gas scale height


# Application to observations



(SAO 206462, Pérez et al., 2014)

$\chi_{\text{obs}} \sim 7$ , model+ data  $\rightarrow v_{\text{turb}} \sim 0.22 c_s$ .

# Gap edges as sites for vortex formation




(Credits: UA grad. student M. Hammer, 'Vortices and orbital migration')

- Disk-planet interaction → gaps
- Surface density maxima at gap edges, or PV minima due to strong shear
- Rossby wave instability → edges 'roll up' into vortices  
(Li et al., 2001)


# Slowly-growing planets make weaker vortices

(Hammer, Kratter & Lin, 2016, in prep.)



- Vortex lifetime/observability depends on planet formation time-scale


Basic theory is 2D, but PPDs are 3D



Take a look in the  $(r, z)$  plane through the vortex

# Rossby vortices are vertically global


$$\Delta\rho/\rho$$



- Global 3D Zeus simulations (Lin, 2012b)
- Consistent with 3D linear theory (Lin, 2012a, 2013a,b)
- Vortex evolution is sensitive to disk vertical structure (Lin, 2014)


# Elliptic instability of 3D vortices

- A 3D instability that weakens/destroys vortices (Lesur & Papaloizou, 2009)
- In plane ( $v_z = 0$ ) **elliptical flow** about the vortex center



- **Instability**  $\rightarrow$  small-scale 3D turbulence ( $v_z \neq 0$ )


$$t=800P_0, 10^2 \langle v_z^2 \rangle^{1/2} / c_s$$




Athena sims. (Lin, in prep.)

# Elliptic instability of 3D vortices

Density pert.




Vort./ $2\Omega$



Athena simulations (Lin, in prep.)

## 3D vortex evolution with self-gravity (Lin, in prep.)




- Vortex collapse correlates with increasing internal turbulence


# 3D vortex evolution with self-gravity (Lin, in prep.)

Density perturbation


$t=1600P_0$ ,  $z=0.0H$ ,  $\Delta\rho/\rho$



$t=800P_0$ ,  $z=0.0H$ ,  $\Delta\rho/\rho$





# 3D vortex evolution with self-gravity (Lin, in prep.)



# What's next on disk vortex dynamics?

- Accretion onto ellipsoidal density distributions
- Gap-opening by vortices



- Dust dynamics in elliptical flows
  - ▶ Explicit 3D sims. to study the effect of hydro. turbulence

# Summary

## Generalized gravitational instability

- Cooling: reduces thermal support
- Viscosity: reduces rotational support
- Fragmentation: GI due to cooling and/or turbulent stresses (viscosity)

## Vertical shear instability

- Feeds off free energy in  $\partial_z \Omega \neq 0$
- Enabled by ultra-fast cooling in PPDs
- VSI possible in the outer PPD between 10—100AU

## Vortices in 3D PPDs

- Can moderate self-gravity enhance vortex stability?
- How can we use the observed structure of vortices to infer disk properties?

# References

- ALMA Partnership 2015, *ApJL*, 808, L3
- Andrews S. M., Wilner D. J., Zhu Z., Birnstiel T., Carpenter J. M., Perez L. M., Bai X.-N., Oberg K. I., Hughes A. M., Isella A., Ricci L., 2016, ArXiv e-prints
- Casassus S., Wright C. M., Marino S., Maddison S. T., Wootten A., Roman P., Pérez S., Pinilla P., Wyatt M., Moral V., Ménard F., Christiaens V., Cieza L., van der Plas G., 2015, *ApJ*, 812, 126
- Chiang E., Youdin A. N., 2010, *Annual Review of Earth and Planetary Sciences*, 38, 493
- Gammie C. F., 2001, *ApJ*, 553, 174
- Lesur G., Papaloizou J. C. B., 2009, *A&A*, 498, 1
- Li H., Colgate S. A., Wendorff B., Liska R., 2001, *ApJ*, 551, 874
- Lin M.-K., 2012a, *ApJ*, 754, 21
- Lin M.-K., 2012b, *MNRAS*, 426, 3211
- Lin M.-K., 2013a, *MNRAS*, 428, 190
- Lin M.-K., 2013b, *ApJ*, 765, 84
- Lin M.-K., 2014, *MNRAS*, 437, 575
- Lin M.-K., Kratter K. M., 2016, ArXiv e-prints
- Lin M.-K., Youdin A. N., 2015, *ApJ*, 811, 17
- Lorén-Aguilar P., Bate M. R., 2015, *MNRAS*, 453, L78
- Lynden-Bell D., Pringle J. E., 1974, *MNRAS*, 168, 603
- Lyra W., Lin M.-K., 2013, *ApJ*, 775, 17
- Macintosh et al. B., 2015, *Science*, 350, 64
- Marois C., Zuckerman B., Konopacky Q. M., Macintosh B., Barman T., 2010, *Nature*, 468, 1080
- Nelson R. P., Gressel O., Umurhan O. M., 2013, *MNRAS*, 435, 2610
- Paardekooper S.-J., 2012, *MNRAS*, 421, 3286
- Pérez L. M., Isella A., Carpenter J. M., Chandler C. J., 2014, *ApJL*, 783, L13
- Rice W. K. M., Armitage P. J., Mamatsashvili G. R., Lodato G., Clarke C. J., 2011, *MNRAS*, 418, 1356
- Rice W. K. M., Lodato G., Armitage P. J., 2005, *MNRAS*, 364, L56
- van der Marel N., van Dishoeck E. F., Bruderer S., Birnstiel T., Pinilla P., Dullemond C. P., van Kempen T. A., Schmalzl M., Brown J. M., Herczeg G. J., Matthews G. S., Geers V., 2013, *Science*, 340, 1199