Paper 1 · the whole draft

The title and the authors' credit are left to the end. This is the current state of everything else, regenerated from the code and its data on every change; the making is public at four.mino.mobi/papers.

Abstract

Gauss's law gives charges in four spatial dimensions a $1/r^2$ potential, under which no atom has a ground state. We ask what atoms exist if the law between charges is instead pinned at $1/r$, the only power law under which orbitals of different angular momentum share a shell. In four dimensions $1/r$ is not harmonic, and its price is the shell theorem: an inner cloud screens the nucleus only partly, and a neutral atom is felt from outside. We solve every neutral atom through $Z = 140$ self-consistently with local exchange, cross-checked by numerical Hartree–Fock, each method tested first on our own world. The table has rows of 2, 10, 10, 28, 28, 60 elements and noble gases at 2, 12, 22, 50, 78, 138, as the generalized Madelung rule predicts, and its larger subshells make room for 52 elements with no counterpart in ours. Solving it again with the shell theorem forced back on separates the geometry from the law: the rows are the geometry's, but what fills inside them from the first d element on is the law's. Partial screening holds outer electrons a median 2.54 times harder, so the hardest-held electron belongs to a full d shell rather than a noble gas; neutral atoms attract with a permanent $-3Q_AQ_B/64R^5$ that outranges dispersion; and four-dimensional space breaks Hund's first rule for d^2, d^{16}, $4f^3$ and $4f^{29}$.

1 · Introduction

Why does space have three dimensions? Ehrenfest (1918, 1920) turned the question into physics. He asked what in the laws of physics would change with the number of dimensions, and found, among other things, that under the inverse-power force Gauss's law gives in D dimensions, no planetary orbit is stable for $D > 3$. Tangherlini (1963) carried the argument into general relativity and into quantum mechanics: the hydrogen atom with Gauss's potential has no stable states above three dimensions. The usual conclusion is that a world with more than three large dimensions has no atoms. Callender (2005) points out what that conclusion leans on, that the laws carry over to other dimensions unchanged.

They needn't. The literature on the hydrogen atom in D dimensions (Nieto 1979; Al-Jaber 1998) and on dimensional scaling (Herschbach, Avery, and Goscinski 1993; Loeser and Herschbach 1986a, 1986b; Ghosh, Kais, and Herschbach 2020) keeps the potential at $1/r$ in every D, mostly as a convention that keeps the problem solvable. Négadi and Kibler (1996) used the D-dimensional hydrogen atom to extend Madelung's filling rule (Madelung 1936) to any dimension; applied to four, it gives rows of 2, 10, 10, 28, 28, 60. Pyykkö and Zhao (1991) solved the atoms of a two-dimensional world by Hartree–Fock. Above three dimensions there are self-consistent atoms, but not a table: Hartree–Fock for two-electron atoms in any D (Loeser and Herschbach 1986a), the Thomas–Fermi atom in n dimensions (Kventsel and Katriel 1981), and many-electron models from the limit of large D (Loeser 1987; Loeser et al. 1994). To our knowledge this paper is the first shell-resolved self-consistent survey of the neutral atoms through $Z = 140$ in four Euclidean dimensions, for a pairwise $1/r$ interaction and electrons with two spin states, with local exchange, no correlation and a first-order correction for open subshells' terms, cross-checked by Hartree–Fock. Nor, to our knowledge, has anyone asked what the choice of $1/r$ itself does to the atoms.

This paper does both. Part I treats the law as a choice: why one has to be made, why $1/r$ exactly, and what it costs in four dimensions (section 2). Part II solves every neutral atom to $Z = 140$ under it (sections 3 and 4) and finds that four-dimensional space breaks Hund's first rule (section 5). Part III solves the table again with the shell theorem forced back on, to find which of its features the law decides (section 6), and gives the pull between neutral atoms that later work on molecules starts from (section 7). The appendices hold the arguments and methods in full, every check of every method against our own world, and the starting data for chemistry: valence states, the atoms' response, and a test of the valence rule. Every number in the text is recomputed from the code and its data whenever anything changes, and checked against a ledger of 40 claims. The solvers, with their known-answer tests and a script that solves the paper's atoms again, are released under the MIT licence at tangled.org/morphyxmino.bsky.social/four-atom; the data and the paper's making are public at four.mino.mobi/papers.

Part I · The law

2 · The law between charges

Ask what the atoms are in a world with four dimensions of space and there is no answer yet, because the question leaves out the law between charges. In ours that law isn't a free choice. Gauss's law, which a local, linear field theory of charge gives, spreads a point charge's field over the sphere around it, whose area grows as r^{D-1}, so the potential falls as $r^{-(D-2)}$. In three dimensions that is $1/r$. In four it is $1/r^2$, and with $1/r^2$ there are no atoms.

2.1 Gauss's law leaves no atom. One electron in the pull $V = -\alpha/r^2$ has the Hamiltonian $H = -\frac{1}{2}\nabla^2 - \alpha/r^2$, and nothing in it is a length. Stretch a wavefunction, $\psi(x) \to \psi(\lambda x)$,

and both terms scale by λ^2, so a bound level at $E < 0$ would bring one at $\lambda^2 E$ for every λ. Either nothing binds or the levels run down without end. Hardy's inequality (Barbatis 2024) decides which: for $\alpha \le \frac{1}{2}$ the kinetic energy always outweighs the pull and nothing binds, and for $\alpha > \frac{1}{2}$ the wavefunction changes sign infinitely often on the way in and the electron falls to the centre (Case 1950; Landau and Lifshitz 1977, sec. 35) (appendix F.1). Our one-electron solver shows both (figure 1a). For $\alpha = 0.45$ it finds no level on any grid. For $\alpha = 0.7$ the lowest level is -5.87×10^3, -5.87×10^5 and -5.87×10^7 hartree as the grid's inner end moves from 10^{-3} to 10^{-5} bohr: set by where the calculation stops, not by the electron. There is no Bohr radius, and classically no stable orbit either (Ehrenfest 1918).




Figure 1. The choice of law. (a) Gauss's law in four dimensions gives a potential $-\alpha/r^2$. One electron in it, solved on grids whose inner end r_0 is moved in: for every α above ½ the lowest level scales as $1/r_0^2$, set by where the calculation stops rather than by the electron (the fall to the centre); at or below ½ there is no level on any grid (Hardy's inequality). Either way there is no Bohr radius and no atom. (b) With a pull $-1/r^\eta$ instead, the splitting between 2p and 2s (and 3d and 3s) as a fraction of the shell's binding vanishes only at $\eta = 1$, to 10^{-11}: the hidden SO(5) symmetry of 1/r, which no other exponent has (section 3.1). Shells of $2(l + 1)^2$, and the rows of figure 2, need exactly 1/r.

2.2 The way out taken. Something has to give: the squeeze $\hbar^2/2m_e r^2$ that resists the pull, or the pull. An electron whose energy grows faster than p^2 binds under Gauss's law, at the cost of Galilean relativity for all matter. A small fourth dimension gives back three-dimensional atoms. This universe keeps the electron and four large dimensions and changes the law: the potential between charges falls as $1/r$. The choice between the first route and this one is a judgement, stated as one (appendix F.2): the electron route keeps our Faraday cages, the law route keeps our mechanics.

2.3 Only 1/r keeps a shell together. Any pull $-1/r^\eta$ with $0 < \eta < 2$ binds and has a length of its own. What singles out $\eta = 1$ is a symmetry. Under $1/r$ every bound orbit is a closed ellipse (Santos, Soares, and Tort 2007), so the Runge–Lenz vector $A_i = \sum_j L_{ij} p_j - Z x_i/r$ is conserved, and in quantum mechanics the rotations SO(D) together with $\mathbf{A}$ generate SO($D + 1$) (Fock 1935; Alliluev 1958; Bander and Itzykson 1966). A level of hydrogen is then the set of degree-$(n - 1)$ harmonics on a sphere one dimension up, so every l below n shares the energy, in four dimensions $-Z^2/2(n + \frac{1}{2})^2$. With $(l + 1)^2$ orbitals to each l, shell n holds 2, 10, 28, 60 electrons: a group of levels with a gap above it. Without the symmetry nothing holds the l together. On the same solver 2p sits −6.2 % from 2s at $\eta = 0.8$, 5×10^{-10} % at $\eta = 1$, +15.4 % at 1.2 and +64.0 % at 1.5, as a fraction of the shell's binding (figure 1b). Rows would still form under another exponent, but they would have to be found anew for each, and the generalized Madelung rule, which starts from hydrogen's degeneracy, wouldn't apply. Under $1/r$ the rows come from counting harmonics, which is what lets Part III tell the geometry's share of the table from the law's (appendix F.3).

2.4 What 1/r costs: the shell theorem. In four dimensions $\nabla^2(1/r) = -1/r^3$ away from the charge, so there is no Gauss's law for it: the flux through a sphere round a charge grows with the sphere (appendix F.4). A thin spherical shell of charge with radius s, felt at radius r, gives $\kappa(t)/\max(r, s)$, $t = \min(r, s)/\max(r, s)$, with κ the law's $1/|x - y|$ averaged over the shell. In ours $\kappa = 1$, the shell theorem, because $1/r$ is harmonic and a harmonic function averages over a sphere to its value at the centre. Here the Laplacian is negative, so $\kappa < 1$: it falls from 1 to $\kappa(1) = 8/3\pi = 0.848826$ at the shell (appendix A, figure A1). Three things follow. An inner cloud screens the nucleus from an outer electron only partly. A hollow conductor's centre sits at $3\pi/8 = 1.178097$ of its wall's potential, so there is no Faraday cage. And a neutral atom is felt from outside, as $-Q/8R^3$ at a distance R, with $Q = \sum \langle r^2 \rangle$ over its electrons (section 7). Everything distinctive in Part III comes from here.

2.5 Where 1/r comes from, and what is declared. A potential $1/r^\eta$ in D dimensions is the Fourier transform of $1/k^{D-\eta}$, so the pinned law needs $1/k^3$ where Maxwell's statics give $1/k^2$. No local field equation gives it. Of the five routes we examined (appendix F.5), the one that survives is a vacuum whose permittivity grows with wavenumber, $\varepsilon_0\, \ell\, k$ with ℓ a length (Tarasov and Trujillo 2013). With the permeability $\mu_0/\ell\, k$, the fields $\mathbf{D}/\varepsilon_0$ and $\mu_0 \mathbf{H}$ obey four-dimensional Maxwell's equations exactly, and what pushes a charge is that light seen through a lens, $\mathbf{E} = (\mathbf{D}/\varepsilon_0)/\ell\, k$; we call it the **lens vacuum**. Its price is relativity, for the force between charges only: the static $1/r$ acts across space at once. This paper uses only its statics, which are all an atom's ground state needs. The free electron keeps its own relativity, which the lens vacuum gives up only for the force between charges, and that fixes how many electrons an orbital holds. In 4 + 1 dimensions the smallest Dirac electron has four components, the product of the four spatial Dirac matrices serves as time's, and a slow electron keeps the two components of one chirality, which turn with one half of a four-dimensional rotation only: two spin states, two to an orbital (appendix F.6). A four-state electron is possible in principle and would double every shell. What stays declared is that the electron is the smallest spinor and that it obeys Pauli's exclusion, which in ours the spin-statistics theorem derives from a local, relativistic theory, as the force here is not. Nuclei are point charges stable to $Z = 140$, and the fine-structure constant is small, so atoms are solved without relativity. Single atoms don't collapse (their energies follow the Thomas–Fermi $Z^{5/2}$ (Kventsel and Katriel 1981), fitted exponent 2.527), but the stability of matter in bulk,

whose proofs in ours lean on the shell theorem, is open (appendix F.7).

Part II · The table

3 · Methods

Everything here is computed by code written for this paper, JavaScript with no outside libraries; nothing is taken from a quantum-chemistry package, and two independent codes check it, Hartree–Fock in a Gaussian basis for any D (McMurchie and Davidson 1978) and, in ours, PySCF (Sun et al. 2020) (appendix E.2). An atom is a point nucleus of charge Z and Z electrons, non-relativistic, with every pair of charges interacting through $q_1 q_2 / r$. Each electron moves in the direction-averaged field of the nucleus and the other electrons' clouds, so the problem is radial. With $\psi = r^{-(D-1)/2}\, u(r)\, Y$, Y a harmonic of degree l on the $(D-1)$-sphere,

$$-\tfrac{1}{2}\, u'' + \left[\frac{L(L+1)}{2r^2} + V(r)\right] u = \varepsilon\, u, \qquad L = l + \frac{D-3}{2},$$

so four dimensions differ from ours by the half-integer shift of L and by the degeneracies, $2(l+1)^2$ electrons to a subshell. Four-dimensional hydrogen's ground state is −2/9 hartree; the solver gives −0.22222. Every electron–electron term goes through the kernel κ of section 2.4, which is where the missing shell theorem enters the atom. Exchange is local, from the uniform electron gas of the same dimension and law, $\varepsilon_x = -A_D\, \rho^{1/D}$ with $A_4 = 16/15\sqrt{\pi}$ (Loos and Gill 2016) (appendix B), each electron's own share removed (Perdew and Zunger 1981); there is no correlation term.

The radial equation is solved by Numerov's method on a logarithmic grid, each level by Sturm bisection, and the field iterated to self-consistency. The ground configuration is searched, not assumed: from the hydrogenic filling, refilled by the orbital energies, then every move of one or two electrons among the open subshells, a candidate counting only if its field settles. First ionization energies are differences of two self-consistent energies, the ion searched for its own ground state. Hund's first rule, which the search assumes, is checked on every open subshell by diagonalizing its terms, and the table is corrected where it fails (section 5). Two further calculations bound the table. The counterfactual solves every element again with the shell theorem forced on, $\kappa \to 1$, everything else unchanged (section 6). Numerical Hartree–Fock, with exact exchange, ranks the same candidates a second way: where it agrees with local exchange a configuration is firm, where it doesn't, open (section 4.3, appendix E.6). The grid, the stop and the quadratures are converged well past what the text quotes (appendix C), every method has a known-answer test run on every change, and appendix G gives the methods in full.

4 · The table

4.1 Rows and noble gases. The table's rows hold 2, 10, 10, 28, 28, 60 elements (and 2 of the next by element 140), and its noble gases are elements 2, 12, 22, 50, 78, 138, each a peak of the first ionization energy (figures 2 and 3). Those are the rows the generalized Madelung rule of Négadi and Kibler (1996) gives in four dimensions; to our knowledge this is the first self-consistent table that confirms them. The rule is a filling order, and like Madelung's in ours it doesn't promise each atom's ground configuration. Read that way, 103 of the 140 are exactly the rule's. Each of the other 37 moves one or two electrons between two subshells, 34 of them from the outer s into the d or f below it, as our chromium, copper and palladium do. The solver finds more of these departures than there are: over our first 36 elements it makes 6 where the measured configurations make 2, from its lean toward d (appendix E.3). So the 4D count is likely too high, and these departures, not the rows, are where the table is least sure.




Figure 2. The periodic table of four dimensions, Z = 1 to 140, each atom solved self-consistently under the pinned 1/r. Rows close at the noble gases 2, 12, 22, 50, 78 and 138, and hold 2, 10, 10, 28, 28 and 60 elements, as Négadi and Kibler's generalized Madelung rule gives when applied to four dimensions. A subshell holds $2(l+1)^2$ electrons, so the p block is 8 wide, the d block 18 and row 6's f block 32 (drawn below, as ours is). An element takes our element's name with its first letter moved on where its place answers to one of ours; the rest (·) have no counterpart. Dots: the elements in figure 4. Hover a cell for its configuration and ionization.




Figure 3. First ionization energy against Z across the four-dimensional table. The noble gases close their rows, but the hardest-held electron isn't a noble gas's: element 128, a full 18-electron d shell with nothing outside it (our palladium's kind), holds its outermost electron at 29.6 eV against its row's noble gas at 26.7; element 68, the same kind a row up, comes within a volt of its row's. In ours palladium holds far less hard than xenon. Across row 6's 32-wide f block (81–112) the binding barely moves.

4.2 The hardest-held electron isn't a noble gas's. Element 128 has an 18-electron d shell, full, with nothing outside it. It holds its outermost electron at 29.56 eV, harder than its row's noble gas 138 (26.71 eV) and harder than any other atom in the table. Element 68, the same kind a row up, comes within a volt of its noble gas 78 (22.32 eV against 23.10). Ours goes the other way: palladium's full $4d^{10}$ gives up an electron at 8.3

eV, xenon at 12.1 (Sansonetti and Martin 2005). Searching each ion for its own ground state leaves 128's ionization where it was. Section 6 shows the law does it.

4.3 How firm each configuration is. Hartree–Fock and local exchange agree on 126 of the 140 ground configurations, and these are firm. The 14 they don't agree on are open: 30, 31, 37, 38, 40, 58, 61 in the d rows, where local exchange puts one electron more in d than Hartree–Fock, as it does in ours; 107, 108, 109, 110, 111 at the end of row 6's f block, where it puts one more in f; and 2 at the start of a block (53, 83). Our own table is the calibration (appendix E.6): there Hartree–Fock gets 36 of 36 configurations right against local exchange's 32, and where the two agree they are right. In row 6's f block both methods overbind f, and our lanthanides measure by how much: Hartree–Fock makes f too stable by at most 4.1 eV. Four-dimensional f electrons are bound so deeply that 17 of the block's 32 elements hold against that bias. The interactive table gives each element's verdict.

4.4 Every element in profile. Figure 4 draws six atoms' radial densities, and figure 5 what their outermost orbitals look like on the 3-sphere. The same two pictures for every element are on the interactive table, where any subshell can be cut by any of the six planes of four-dimensional space (four.mino.mobi/table; `#Z` opens on element Z). The configurations, energies, ionizations and radii are in `data/table4.csv`. What chemistry starts from, each atom's valence states and a direct test of the valence rule on darbon's hydride DH_5, is in appendix I.




Figure 4. Six four-dimensional atoms in profile: each subshell's electrons per unit ln r against distance, from the solved atoms; the area under a curve is its electron count, and the outermost subshell is drawn thick. Darbon (7), our carbon's place, five electrons of a ten-electron shell; element 8, which answers to nothing of ours; oeon (12), the first noble gas past helium, its p shell holding eight; uitanium (26), where the d block starts; element 101, 21 electrons into an f shell that holds 32; and element 128, the hardest-held electron in the table. Every element's profile: four.mino.mobi/table/#Z.




Figure 5. Orbitals four dimensions has and ours doesn't, cut through the nucleus by a plane, on the solved atoms' own radial functions. (a) The fourth p orbital, p_w, in the x–w plane: a dumbbell along w, so any three-dimensional slice through the nucleus misses it entirely. (b–d) One of the nine 4D d orbitals, $x^2 + y^2 - z^2 - w^2$, in three planes: positive all the way round the x–y plane, negative all the way round the z–w plane, and a four-leaf clover in a plane that mixes them. The x–y and z–w planes are completely perpendicular and meet only at the nucleus; nothing in three dimensions is positive round one plane and negative round another. Any element's orbitals, in any plane: four.mino.mobi/table/#Z; hydrogen's exact ones, turned live: kata.mino.mobi/orbital.

5 · Hund's first rule in four dimensions

The solver gives each open subshell the most parallel spins it can hold. That is Hund's first rule, the known answer for the ground terms of free atoms in ours. In four dimensions it is an assumption, and for some fillings it is wrong. To test it, every arrangement of each open subshell's electrons is diagonalized under the law, on the atom's own orbital (appendix H.1). In ours this reproduces Racah's term separations to the wavenumber, and maximum spin is lowest in all 23 open subshells of our table.

5.1 d^2, d^{16}, $4f^3$ and $4f^{29}$ break the rule. In four dimensions all 36 open p subshells keep the rule, and so do the d fillings d^1, d^3–d^{12}, d^{14}, d^{15}, d^{17}. The exceptions are d^2 and its mirror d^{16}, in all three d rows: element 26 ($3d^2$, 0.40 eV), element 39 ($3d^{16}$, 0.77 eV), element 54 ($4d^2$, 0.30 eV), element 66 ($4d^{16}$, 0.58 eV), element 114 ($5d^2$, 0.36 eV), element 126 ($5d^{16}$, 0.63 eV). In each the ground term is a spin singlet of ten states, and the triplet the solver assumed lies 0.30–0.77 eV above it. In the f block $4f^3$ (element 83) fails too, and so does its mirror $4f^{29}$ (element 108), each with a ground term of spin ½, 0.15 and 0.26 eV below the quartet. No other f filling was found to break the rule, though for the fullest blocks only the highest spins could be run (appendix H.2).

5.2 The four-dimensional shell breaks it, not the law. On element 26's own orbital Gauss's law $1/r^2$ gives the same order, the singlet 0.82 eV below the triplet against 0.40 eV under $1/r$. The cause is the shell. For two electrons in a d subshell each term's energy is exactly $F^0 + a_2F^2 + a_4F^4$: the F^n are the radial parts of the repulsion and depend on the orbital, and the $a_n = \langle C_n(\cos\gamma)\rangle$ over the term depend only on the term (γ the angle between the two electrons seen from the nucleus). The ten-state singlet, $(2,0) \oplus (0,2)$ of SO(4), holds the electrons nearest right angles, $\langle \cos^2\gamma \rangle = 0.125$ against random's 0.250 and the best triplet's 0.188, and lies below every triplet whenever $F^4/F^2 < 5/3$. Element 26's orbital has 0.39 under the law and 0.65 under Gauss's, so the rule breaks either way. In ours the triplet ^{3}F is the term that keeps the electrons nearest right angles, and it is the lowest for every F^4/F^2. Pauli

keeps same-spin electrons apart, which is the usual reason for Hund's rule; four-dimensional space gives paired spins a better way. Every four-dimensional rotation splits into a left and a right half, a d orbital carries one unit of each, and the singlet adds the two electrons' left halves and cancels their right ones, or the other way round. Three-dimensional rotations don't split, so ours has no such state (appendix H.3; interactive at four.mino.mobi/atom/hund.html).

5.3 The correction moves energies and ionizations, no configuration. Where the rule fails, the solver's energy is the maximum-spin state's, too high by the gap. Every d- and f-block element was searched again with each candidate's gaps added (appendix H.4). **The terms change no ground configuration.** They lower the eight Hund-breaking atoms' energies and move first ionizations in pairs, by 0.30–0.67 eV: up for an atom whose own ground term gains the gap, down for the next, whose ion lands on the broken filling. The wider search found one configuration the bake had missed, element 82's $4f^1\ 5d^1\ 6s^2$, our cerium's kind. The table of section 4 is the corrected one (`data/table4.csv`); the Hund-spin table is supplement S1. In a ligand field the singlet survives on the 16-cell's eight axes and gives way to a triplet, barely, at a tesseract's sixteen corners (appendix H.5).

Part III · What the law does

6 · What the law decides

To see which of the table's features the law decides, every 4D element was solved a second time with one thing changed: the clouds' potential taken as if the shell theorem held ($\kappa = 1$), and everything else, the four-dimensional kinetic energy, the degeneracies, exchange, the search, left as it was. Each search was seeded with the real table's configuration, and the real law was re-solved in every counterfactual configuration, so a difference is the law's, not a weaker search (appendix G.10). Of the 140 elements, 112 settle, every one up to element 88; the rest, 89–116, have no settled ground state under the shell theorem from any start tried, so what follows about the rows holds through row 5.

6.1 The rows are the geometry's. With the shell theorem forced on, the first 24 elements keep every configuration, and the noble gases close the same rows at elements 2, 12, 22, 50, 78, 138 (figure 6). The rows' lengths, 2, 10, 10, 28, 28, come from four-dimensional space's $2(l + 1)^2$ to a subshell, not from the law, so long as the law keeps a shell together (section 2.3).




Figure 6. What the law does. Every 4D atom solved a second time with the shell theorem forced on (the clouds' pull as if each sat at the centre), everything else held. (a) The table: grey, the same ground configuration under both laws; orange, changed; a question mark in place of the element's number, no settled ground state under the shell theorem (row 6's f block, 89–116, after a second, wider search). Every atom up to 88 settles. The first 24 elements and all six noble gases are unchanged: the rows are the geometry's, at least through row 5. Every change starts at the first d element: without the law, p beats d inside each row. (b) First ionization under both: the law holds outer electrons a median 2.5 times harder, and element 128 nearly ten times (29.6 eV against 3.1); with the shell theorem the noble gases hold hardest, as ours do.

6.2 What fills inside the rows is the law's. Of the settled atoms, 57 change configuration, the first at element 25, the first d element. Under the shell theorem p beats d: element 25 takes $4s^2\ 4p^1$ where the law gives $3d^1\ 4s^2$, and the change runs on through the d and f blocks.

6.3 Partial screening holds outer electrons harder. In ours an inner cloud screens the nucleus from an outer electron whole. Here it screens only partly, by $\kappa(t)$, so outer electrons see more of the nucleus. Across the 110 atoms past helium that settle under both, the law holds the outermost electron a median 2.54 times as hard as the shell theorem would, from 1.06 to 9.55; among the noble gases the ratio grows from 1.09 (helium) to 2.90 (element 138). The largest is element 128's: its full $5d^{18}$ shell holds its electron at 29.56 eV under the law and at 3.1 eV under the shell theorem, where the noble gases are again the hardest-held atoms, as ours are. The table's hardest-held electron is partial screening at its strongest, a full shell whose own cloud hides the nucleus from it least.

6.4 Negative ions. Partial screening holds an extra electron harder too. One electron short of a full p shell (p^7 in four dimensions), atoms bind an extra electron by 3.4, 5.7, 7.8, 10.3, 12.9 eV going down the group, where ours stay level (F 3.401, Cl 3.613, Br 3.364 eV, measured). 80 of the 140 bind one by more than 0.7 eV, the level above which the method held in ours (appendix E.4); the light noble gases don't hold one at all.

7 · Neutral atoms aren't silent

Under the shell theorem a neutral atom is invisible from outside its cloud. Here it isn't. A shell of charge q and radius s, felt at a distance R outside it, gives $q\kappa(s/R)/R$ with $\kappa = 1 - t^2/8 - \dots$ (appendix A). Summed over the nucleus and the electrons the leading terms cancel, and what remains, for an electron at distance R, is a pull $-Q/8R^3$ with $Q = \sum \langle r^2 \rangle$. The solved noble gases follow it: at 30 bohr the full cloud's potential and the far form agree to 0.22 %. Between two atoms the same failure gives a permanent attraction $-3Q_A Q_B/64R^5$, with no dipole needed, which outranges London's R^{-6}.

For two like atoms the permanent pull's coefficient is $C_5 = 3Q^2/64$, and $R^* = C_6/C_5$ is the distance inside which dispersion is the larger, with C_6 from the solver's coupled response by Casimir and Polder's integral (appendix J.1, table 2; in ours its noble gases' polarizabilities are off by −8 % to +4 % against Derevianko, Porsev, and Babb (2010)). For hydrogen and helium R^* is 28 and 9.26 bohr, for the alkali-like and alkaline-earth-like atoms 4.9–94, and for every other atom in the table 0.59–1.5, inside the atom. Past contact, for most of the table, the permanent term is the leading pull between neutral atoms at every distance where a far-field form applies. Pulled apart by Hartree–Fock, two closed-shell atoms follow it (1.02–1.06 times the far form at 16 bohr), so the atoms the molecular codes separate into are this paper's (appendix J.2).

That sets a requirement on every later model of four-dimensional matter. In ours the trace of a charge distribution's second moment is dropped from multipole expansions, because $1/r$ is harmonic and the trace carries no field. Here it is a neutral atom's leading field, so a four-dimensional force field has to keep it; the later papers' do, as a cloud term on every atom.

8 · Limitations and conclusion

Limitations. The force acts instantly (section 2.5), so only statics are safe, and nothing here moves. Heavy atoms would need relativity, which the law has given up for the force between charges, so the table past about element 100 is a first pass; round a point nucleus the innermost Dirac level fails at $Z \approx 205$ in four dimensions (Katsura and Aoki 2006; GU, MA, and DONG 2002), beyond the table. Nuclei are declared stable to $Z = 140$. The method has local exchange and no correlation, which in ours favours d by about a volt (appendix E.3), every subshell's cloud averaged over the sphere, and open subshells' terms resolved to first order on the solver's own orbitals. The counterfactual leaves row 6's f block unsettled. Nothing here claims that four-dimensional chemistry looks like ours; the next papers take up bonds, liquids and crystals under the same law.

Conclusion. A world with four large dimensions of space has to choose its law between charges, because Gauss's law gives it no atoms. Pinned at $1/r$, the only power law that keeps a shell together, it has atoms, and they have a periodic table: rows of 2, 10, 10, 28, 28, 60, noble gases at 2, 12, 22, 50, 78, 138, as the generalized Madelung rule predicts. The rows are the geometry's. What fills inside them from the first d element on is the law's, because in four dimensions $1/r$ isn't harmonic and an inner cloud screens only partly: outer electrons are held about two and a half times harder, the hardest-held electron belongs to a full d shell rather than a noble gas, and a neutral atom pulls from afar. And four-dimensional space breaks Hund's first rule for d^2, d^{16}, $4f^3$ and $4f^{29}$, under both laws tested.

Appendices

Notation

Atomic units throughout: $\hbar = m_e = e = 1$, and the law's constant is 1, so two charges q_1 and q_2 a distance r apart have the energy $q_1 q_2/r$ in hartree with r in bohr. "Ours" is our own three-dimensional world. The subshell letters s, p, d, f (upright) are $l = 0, 1, 2, 3$, and in four dimensions a subshell of angular momentum l holds $2(l + 1)^2$ electrons. Bold letters are vectors.

symbol	meaning	from
D	the number of dimensions of space: 3 in ours, 4 here	§2
r	distance from the nucleus, or between two charges	§2
x, y; x_i	points of D-dimensional space; the coordinates of x	§2.1
$V(r)$	the potential energy of one electron	§2.1
a	the strength of Gauss's pull $-a/r^2$ in four dimensions	§2.1
H, ψ	a one-electron Hamiltonian and its wavefunction	§2.1
∇, ∇^2	gradient and Laplacian in D dimensions	§2.1
λ	a scale factor, $x \to \lambda x$	§2.1
σ	the power r^σ of a wavefunction close to the nucleus	App. F.1
ε	a one-electron (orbital) energy, in hartree	§3
E	a total energy; $\mathbf{E}$ (bold) is the electric field	§2.1
r_0	the inner end of a calculation's radial grid	App. F.1
p, $\mathbf{p}$	momentum	§2.2
$\hbar, m_e$	Planck's constant over 2π and the electron's mass (both 1 in atomic units)	§2.2
η	the exponent of a trial law $V = -1/r^\eta$ ($\eta = 1$ is the pinned law, $\eta = D - 2$ Gauss's)	§2.3
Z	the nuclear charge, the element's number	§2.3
L_{ij}, $\mathbf{A}$	angular momentum in the plane ij; the Runge–Lenz vector	§2.3
k	wavenumber (Fourier space); k_F is the Fermi wavenumber	§2.5
n, l	the principal and angular quantum numbers	§2.3
s	the radius of a spherical shell of charge	§2.4
t	$\min(r, s)/\max(r, s)$, the ratio of the two radii	§2.4
$\kappa(t)$	the shell kernel: a shell's potential over the shell theorem's, $\kappa = 1$ in ours	§2.4
R	the distance from an atom to an outside charge, or between two atoms	§2.4
Q	an atom's $\sum \langle r^2 \rangle$ over its electrons; Q_A, Q_B for atoms A and B	§2.4
$\alpha_p(\mathrm{i}\omega)$	an atom's dipole polarizability at imaginary frequency; $\alpha_p(0)$ the static one	App. J.1
C_6, C_5	the coefficients of two like atoms' dispersion $-C_6/R^6$ and permanent pull $-C_5/R^5$, $C_5 = 3Q^2/64$	§7
R^*	C_6/C_5, the distance inside which dispersion is the larger	§7
Δ	the spread of a d subshell's one-electron levels in a ligand field	App. H.5
φ	an electric potential	App. F.5
ω, c	angular frequency; the speed of light	App. F.5
ρ	a density: of charge in §2.5, of electrons in an atom from §3	§2.5
ε_0, μ_0	the vacuum's permittivity and permeability	§2.5
ℓ	the lens vacuum's length: permittivity $\varepsilon_0\ \ell\ k$, permeability $\mu_0 / \ell\ k$	§2.5
β	the exponent of the nonlinear vacuum, $\nabla \cdot (\lvert\mathbf{E}\rvert^{\beta-2}\mathbf{E}) = \rho$	App. F.5
$\mathbf{E}$, $\mathbf{D}$, $\mathbf{H}$	electric field, displacement, magnetic field	§2.5
N	a number of electrons, or of atoms	App. F.7
$u(r)$	the radial function: $\psi = r^{-(D-1)/2}\, u(r)\, Y$ with Y a harmonic of degree l	§3
L	$l + (D - 3)/2$, the angular momentum the radial equation sees	§3
ε_x, v_x, A_D	exchange energy per electron, exchange potential, and the exchange constant: $\varepsilon_x = -A_D\, \rho^{1/D}$	§3
h	the grid's step in $\ln r$	App. G.5
S, M_S	an open subshell's total spin and its projection	App. H.1
F^n	the radial (Slater) integrals of the repulsion within a subshell, $n = 0, 2, 4, \ldots$	§5.2
C_n	the Gegenbauer polynomial $C_n^{(1)}$, the 3-sphere's analogue of Legendre's P_n	§5.2
γ	the angle between two electrons, seen from the nucleus	§5.2
a_n	$\langle C_n(\cos\gamma)\rangle$ over a term, its share of F^n	§5.2
θ	the polar angle on a sphere	App. A
K, E (upright)	complete elliptic integrals of the first and second kind	App. A
P_n	the Legendre polynomial	App. A
Ω_D	the volume of the unit ball in D dimensions ($\pi^2/2$ for $D = 4$)	App. B

A · The 4D kernel

A thin spherical shell of unit charge and radius s, felt at radius r, gives the potential $\kappa(t)/\max(r,s)$, where $t = \min(r,s)/\max(r,s)$ and κ is the law's $1/|x-y|$ averaged over the shell. On the 3-sphere the average over directions carries the weight $\frac{2}{\pi}\sin^2\theta\,d\theta$, so

$$\kappa(t) = \frac{2}{\pi}\int_0^\pi \frac{\sin^2\theta\,d\theta}{\sqrt{1+t^2-2t\cos\theta}}.$$

In ours the same average, with weight $\frac{1}{2}\sin\theta\,d\theta$, is exactly 1 for every t: the shell theorem. In four dimensions it isn't. The integral has a closed form in the complete elliptic integrals K and E of modulus t,

$$\kappa(t) = \frac{4\left[(1+t^2)\,\mathrm{E}(t) - (1-t^2)\,\mathrm{K}(t)\right]}{3\pi t^2},$$

which gives $\kappa(1) = 8/3\pi$ (since E(1) = 1), and the series

$$\kappa(t) = 1 - \frac{t^2}{8} - \frac{t^4}{64} - \frac{5t^6}{1024} - \frac{35t^8}{16384} - \cdots$$

Each term is t^n times the average of the Legendre polynomial $P_n(\cos\theta)$ over the 3-sphere; the odd ones vanish. The solver evaluates κ by Simpson's rule in θ with 400 intervals, tabulated at 2001 points of t and interpolated. The closed form agrees with that quadrature to 2.7×10^{-15} for t up to 0.9 (ledger `kernelclosed`); nearer 1 the quadrature's own error dominates, a few parts in 10^7.

The series' first correction is the far pull of section 7. A cloud of N electrons around a nucleus $+Z$, seen from a distance R outside it, gives $(Z - N + \sum_i \langle r_i^2\rangle/8R^2 + \ldots)/R$. A neutral atom therefore acts on an electron through $-Q/8R^3$ with $Q = \sum\langle r^2\rangle$, and two neutral atoms on each other through $-3Q_AQ_B/64R^5$ (section 7). Figure A1 draws κ, and the potential inside and outside a shell.




Figure A1. The pull of a thin shell of charge on a point at another radius, over a point charge's at the shell's centre, as κ(t) of the ratio t of the two radii. The shell theorem makes it 1; under the pinned 1/r in four dimensions it falls to 8/3π at the shell itself.

B · The 4D electron gas

The solver's exchange is the local form of the uniform electron gas of the same dimension and the same law. In D dimensions a spin-unpolarized Fermi sea of density ρ has $\rho = 2\,\Omega_D k_F^D/(2\pi)^D$, with Ω_D the volume of the unit ball ($\pi^2/2$ for $D = 4$). Its kinetic energy per volume is $\frac{D}{D+2}\rho\,k_F^2/2$: $\frac{3}{10}(3\pi^2)^{2/3}\rho^{5/3}$ in ours (Thomas–Fermi's 2.8712), $\frac{4\pi}{3}\rho^{3/2}$ in four dimensions. Its exchange energy per volume is $-A_D\,\rho^{1+1/D}$. In ours $A_3 = \frac{3}{4}(3/\pi)^{1/3} = 0.738559$, Dirac's constant. In four dimensions under the pinned $1/r$, with the Fermi sea's density matrix built on the Bessel function J_2 and the integral $\int_0^\infty J_2(x)^2\,x^{-2}\,dx = 4/15\pi$, it is $A_4 = 16/15\sqrt{\pi} = 0.601802$. These agree with Loos and Gill's expressions for the D-dimensional gas (2016). The selftest checks the textbook values in ours and the 4D integral (blocks 4 and 18). The potential each electron feels is $v_x = -(1+1/D)\,A_D\,\rho^{1/D}$, spin-polarized as in section 3 (each spin's own density doubled), with each electron's own share taken out.

C · Numerical convergence

The solver's stop. The self-consistent field stops when the total energy changes by less than 10^{-7} hartree between iterations. Re-solving 19 atoms (4D 2, 4D 7, 4D 12, 4D 13, 4D 22, 4D 26, 4D 50, 4D 51, 4D 68, 4D 78, 4D 101, 4D 128, 4D 138, ours 2, ours 10, ours 11, ours 18, ours 26, ours 36) with the stop at 10^{-12} moves no total energy by more than 2.7×10^{-7} hartree and no first ionization by more than 2.8×10^{-7} (ledger `converged`).

The radial grid. The baked grid has 700 points from 10^{-5} to 150 bohr, logarithmic with step 0.02364. 11 atoms (4D 2, 4D 12, 4D 13, 4D 22, 4D 50, 4D 78, 4D 128, 4D 138, ours 10, ours 18, ours 36) were re-solved on six other grids, each changing one parameter (ledger `grid`):

change from the baked grid	largest |ΔE| (hartree)	largest |Δ ionization| (hartree)
N 500	2.0×10^{-2}	1.4×10^{-4}
N 1000	1.0×10^{-2}	7.2×10^{-5}
r_0 10^{-4}	6.7×10^{-2}	3.8×10^{-5}
r_0 10^{-6}	5.4×10^{-3}	4.2×10^{-5}
r_max 100	1.0×10^{-3}	6.9×10^{-6}
r_max 250	1.3×10^{-3}	8.9×10^{-6}

The total energies of the heaviest atoms move by up to 6.7×10^{-2} hartree, 2.4×10^{-5} of themselves, with the innermost point. The ionization energies, the differences the paper's chemistry uses, move by at most 1.4×10^{-4} hartree.

D · The law calculations

Section 2's arguments use a separate one-electron solver (`laws.js`), so that the grid's inner end can be moved. It uses the radial equation with $L = l + (D - 3)/2$ on a logarithmic grid of 6000 points, counts the nodes of an outward Numerov solution at a trial energy, and bisects on the count to 10^{-13} relative; a level that falls with the inner end of the grid is a fall to the centre. Its check is 4D hydrogen, −0.22222222 hartree against −2/9.

Gauss's $-\alpha/r^2$. At α = 0.45 there is no bound level on any grid. At α = 0.7 the lowest level is -5.87×10^3, -5.87×10^5 and -5.87×10^7 hartree with the inner end at 10^{-3}, 10^{-4} and 10^{-5} bohr: $\varepsilon r_0^2 = -0.00587$ and −0.00587, a fall to the centre set by r_0 (ledger `gauss`). The boundary between the two is Hardy's constant $((D-2)/2)^2 = 1$ (Barbatis 2024), which with the kinetic energy $\frac{1}{2}|\nabla\psi|^2$ puts the critical α at $\frac{1}{2}$ (Landau and Lifshitz 1977, sec. 35).

Power laws $-1/r^\eta$. The splitting of 2p from 2s, as a fraction of the shell's binding, is −6.2 % at η = 0.8, 5×10^{-10} % at η = 1, +15.4 % at η = 1.2 and +64.0 % at η = 1.5. For 3d against 3s it is −9.2 %, +21.9 % and +78.3 % (ledger `uniqueness`). Only η = 1 keeps a shell together.

E · Checks against our world

E.1 Hydrogen. One electron needs no search and no exchange, and four-dimensional hydrogen has an exact answer: $-Z^2/2(n+\frac{1}{2})^2$ hartree, so −2/9 for the ground state. The solver gives −0.22222.

E.2 Two electrons. Helium-like atoms in four dimensions are checked against an independent code, Hartree–Fock in a Gaussian basis written for any dimension (`gauss.js`), and the same atoms in ours against PySCF at the MP2 level (selftest blocks 17 and 43). The two codes share no part of the radial solver. From outside, the only accurate four-dimensional numbers we have been able to read are correlated variational energies for helium-like ions (Escobar-Ruiz et al. 2021). Hartree–Fock (`hf.js`, appendix G.12) lies above them by 0.012–0.014 hartree for Z = 2–10, as it must, since they include correlation; that is the size and sign of a correlation energy, a consistency check rather than a benchmark. A four-dimensional Hartree–Fock limit to test against (Loeser and Herschbach 1986a) is in tables we have not obtained.

E.3 Our own table, and the solver's lean toward d. The same solver, the same law (in ours 1/r is Gauss's) and the same search, run in three dimensions, gives 32 of 36 of our measured ground configurations. The four it misses, titanium, vanadium, cobalt and nickel, each take one d electron too many against 4s, the known lean of local exchange toward d. The misses have one cause. Moving an electron from 4s into 3d comes out too cheap in every 3d atom we can compare (eV, the solver against measured: Sc 0.14 against 1.43, Ti −0.89 against 0.81, Fe 0.51 against 0.87, Co −0.35 against 0.42, Ni −1.16 against −0.03), short by 0.37–1.70 eV and a mean 1.05, always in d's favour. The measured values are differences of each configuration's lowest term, its levels averaged with weights $2J+1$, from NIST's level tables (Sansonetti and Martin 2005), because the solver has no spin–orbit coupling and lands each configuration in its lowest term. Taking the lowest level of each instead moves them by at most 0.06 eV and turns nickel's sign (+0.03 rather than −0.03). That is local exchange without correlation. Resolving every open subshell's whole term structure doesn't remove it. With that correction (the lowest term less the average of the maximum-spin arrangements, which is what a spherically averaged field stands for), the bias stays at 1.05 eV and the configurations get worse, 26 of 32 against 28 of 32. So the paper corrects only Hund's first rule (section 5), the one thing a spin-polarized averaged field cannot represent. Its first ionization energies are off by a median 4.3 % and at most 16.2 % (figure A2). That is the accuracy to expect of the 4D table: a method that gets our transition metals mostly right and our ionizations to a few percent, not a spectroscopic one.




Figure A2. The solver in our three dimensions. (a) Computed against measured first ionization energies (NIST), with the four elements whose ground configuration it misses (Ti, V, Co, Ni: one d electron too many against 4s, the known lean of local exchange). (b) The percentage error against Z.

E.4 Ions. Each ion of ours searched for its own ground state, rather than made by taking one electron from the atom's configuration, changes no first ionization. Searched the same way, our +2 and +3 ions give second and third ionization energies within a median 2.0 % and 1.4 % of measured ones (Kramida et al. 2026) (worst V, 7 % and C, 4 %), closer than the first

ionizations' 4.3 %, and they lose their 4s electrons before 3d, as measured (Fe^{+} $3d^6 4s$, Fe^{2+} $3d^6$, Fe^{3+} $3d^5$). In four dimensions the 3d row's ions do the same. Elements in the first half of the 4d and 5d blocks, and the two that open row 6's f block with a 5d electron, instead give up a d electron at +1 or +2 while keeping the outer s (53–56, 58, 61, 81–82, 114–118, 120, 122), as our yttrium and lutetium do (Y^{+} $5s^2$, Lu^{+} $6s^2$). The same search for anions gives our electron affinities where the extra electron is bound by the field and exchange: the p-block acceptors (C 1.32 against 1.262, Si 1.30 against 1.39, Ge 1.25 against 1.233, Al 0.40 against 0.433, F 2.80 against 3.401, Cl 3.15 against 3.613, O 0.71 against 1.461 eV, measured second). It also finds every anion that should be unbound to be unbound: the noble gases, beryllium, magnesium and nitrogen. It fails where correlation does the binding (H -0.33 against 0.754, Li -0.08 against 0.618, Na -0.02 against 0.548, P -0.10 against 0.746). In ours, then, the method's affinities hold above about 0.7 eV and not below (the measured values are from Rienstra-Kiracofe et al. (2002)). That is a calibration in our world, not an error bar for four dimensions: an affinity near threshold there would need its own checks, a wider radial grid for the extra electron, the anion's and the atom's fields both converged, and correlation.

E.5 Ionizations are converged in stop and grid. Tightening the solver's stop or changing its grid moves no ionization energy by a millihartree (section 3 and appendix C).

E.6 Hartree–Fock as a second method. Hartree–Fock (appendix G.12) on the same candidates gets 36 of 36 ground configurations of ours right, against local exchange's 32. The two agree on 32, and every one of those is right; they differ on Ti, V, Co and Ni, and on each of these Hartree–Fock is right. The promotion of a 4s electron into 3d shows what each method does, against the same measured centroids as appendix E.3 (eV, error of local exchange and of Hartree–Fock: Sc −1.29/−0.42, Ti −1.70/−0.27, V −2.09/−0.12, Cr −1.73/−0.26, Mn −0.72/+1.19, Fe −0.37/+0.93, Co −0.76/+1.11, Ni −1.13/+1.31, Cu −0.44/+1.12). Local exchange makes d too cheap in all nine. Hartree–Fock makes it a little too cheap up to half filling (Sc–Cr, −0.42 to −0.12) and too dear after it (Mn–Cu, 0.93 to 1.31). So the two do not bracket the measured value; from scandium to chromium both lie on the same side of it. Their difference is a measure of how much an answer depends on the method, not an error bar: neither method bounds the other, and both omit the correlation that would move each toward the measured values.

In four dimensions the two agree on 126 of the 140 elements, and these are the table's firm configurations. They differ on 14, which are open: 7 in the d rows (30, 31, 37, 38, 40, 58, 61), where local exchange puts one electron more in d than Hartree–Fock, as in ours; 5 at the end of row 6's f block (107, 108, 109, 110, 111), where it puts one more in f; and 2 at the start of a block (53, 83). For the d rows, ours gives a lean. Before half filling (30, 31, 58) Hartree–Fock's error in ours was small and in the same direction as local exchange's, so its preference for s^2 should hold. After it (37, 38, 40, 61) its margins, 0.10 to 1.08 eV, are inside the volt by which it overprices d in ours. In the f row the first comparison was incomplete. Its candidates lay within the window of local exchange's ground, and local exchange overbinds f, so configurations with one f electron fewer fell outside it. A window of 10 eV and a climb through Hartree–Fock's own neighbourhood (appendix G.12) close that gap, and our lanthanides calibrate what remains. Against the measured levels (Sansonetti and Martin 2005), Hartree–Fock gets 15 of 18 lanthanide ground configurations right (lanthanum to hafnium) and local exchange 12; where the two agree they are right 12 times in 14. In each of the 17 measured states that change the f count, Hartree–Fock makes f too stable, by 0.15 to 4.1 eV. So a configuration with one f electron fewer than Hartree–Fock's ground could truly lie below it only if Hartree–Fock puts it less than B = 4.1 eV up; one with more f only moves further up. In four dimensions f is bound far more deeply. The cheapest move of an f electron out of Hartree–Fock's ground costs 5.8 eV in element 90, 12.6 in 100 and 19.9 in 109, against 0.9 eV in our terbium, where the measured cost is 0.04. Hence four verdicts on row 6's f block:

- **Firm** where the methods agree and every competitor with fewer f lies more than B up, or the ground has no f: 81, 88–100 and 102–104.
- **Open though the methods agree** where such a competitor lies within B: 82 (1.53 eV), 84 (0.40), 85 (1.12), 86 (2.21) and 87 (3.86). Element 84 is the case of our cerium, whose measured $4f^1 5d^1 6s^2$ both methods miss, agreeing on $4f^2 6s^2$.
- **Leaning to Hartree–Fock** where local exchange's choice holds one f electron more: 83 and 107–111. Local exchange overbinds f by more still, so its extra f electron is the less likely.
- **Untested** where no configuration with fewer f was solved: 101, 105, 106 and 112.

So 17 of the block's 32 are firm, because in four dimensions the f shell's binding outruns the calibration's error; the open cases sit at the block's two ends, where f competes with d and s at the start and is about to close at the end. In the d rows no Hartree–Fock ground configuration lies near the window's edge. Every element's verdict, both methods' rankings and the margins are in `data/hftable.json`.

F · The law in full

F.1 The fall to the centre. In $H = -\frac{1}{2}\nabla^2 - a/r^2$ nothing is a length (section 2.1), so either nothing binds or the levels run down without end. Which of the two happens is decided at the origin. Close in, an s wavefunction goes as r^σ with

$$\sigma(\sigma + D - 2) + 2a = 0, \qquad \sigma = -1 \pm \sqrt{1 - 2a} \quad (D = 4).$$

For $a \le \frac{1}{2}$ the roots are real, and Hardy's inequality (Barbatis 2024)

$$\int |\nabla\psi|^2 \;\ge\; \left(\frac{D-2}{2}\right)^2 \int \frac{|\psi|^2}{r^2} \;=\; \int \frac{|\psi|^2}{r^2} \quad (D = 4)$$

says the kinetic energy $\frac{1}{2}\int |\nabla\psi|^2$ always outweighs the pull: $H \ge 0$, and nothing binds. For $a > \frac{1}{2}$ the roots are complex and

the wavefunction near the origin is $r^{-1}\cos(\sqrt{2\alpha-1}\,\ln r+\delta)$. It changes sign infinitely often on the way in, and every node below a level means another level beneath it, so there is no lowest one: the electron falls to the centre (Case 1950; Landau and Lifshitz 1977, sec. 35).

A calculation shows both. Our one-electron solver (`laws.js`, appendix D) integrates the radial equation of section 3 by Numerov's method on a logarithmic grid that stops at an inner radius r_0, and finds each level by bisecting on the number of nodes. For $\alpha = 0.45$ it finds no bound level on any grid. For $\alpha = 0.7$ the lowest level is -5.87×10^{3}, -5.87×10^{5} and -5.87×10^{7} hartree with $r_0 = 10^{-3}$, 10^{-4} and 10^{-5} bohr. It grows as $1/r_0^2$ ($\varepsilon r_0^2 = -0.00587, -0.00587$), set by where the calculation stops rather than by the electron (figure 1a). Either way there is no Bohr radius, no size an atom could settle at. Ehrenfest made the classical version of this argument (Ehrenfest 1918): under Gauss's law in four dimensions no planetary orbit is stable either.

F.2 The ways out, and what each costs. Something has to give. The squeeze that resists the pull costs $\hbar^2/2m_e r^2$, because the electron's energy grows as the square of its momentum p. The pull goes as $1/r^2$ because of Gauss. Change either and a length comes back. One way is to change the electron: an energy growing as p^3 or p^4 outgrows a $1/r^2$ pull close in, so atoms bind at every strength. It keeps Gauss's law, and with it the shell theorem and the Faraday cage. But it gives up Galilean relativity for all matter: a moving laboratory and a still one see the same physics only if energy is quadratic in momentum. Another is a small fourth dimension, the way extra dimensions could hide in ours. But atoms larger than it are three-dimensional atoms and smaller ones fall again, so it doesn't give four-dimensional chemistry. The third is to change the law: keep the electron, keep four large dimensions, and make the potential between charges fall as $1/r$. This universe takes the third, and section 2 says why $1/r$ and what it costs. The choice between the first and the third is a judgement, stated as one: the electron route keeps our Faraday cages, the law route keeps our mechanics.

F.3 The symmetry behind the shells. Classically, under $1/r$ every bound orbit is a closed ellipse. By Bertrand's theorem only $1/r$ and r^2 do that (Santos, Soares, and Tort 2007), and r^2 confines rather than falling off; asking for closed orbits is a criterion for choosing the law, not something every atom needs. An orbit in a central potential lies in the plane of its starting position and velocity in any number of dimensions (FIORANI 2009), so the planar theorem holds in four. A closed ellipse keeps a fixed direction, its long axis. So the motion conserves a vector besides the angular momentum, the Runge–Lenz vector $A_i = \sum_j L_{ij}\,p_j - Zx_i/r$, where $L_{ij} = x_i p_j - x_j p_i$ is the angular momentum in the plane ij. Under any other exponent the ellipse precesses, its axis turns, and $\mathbf{A}$ isn't conserved.

In quantum mechanics the rotations of D-dimensional space, SO(D), together with the D components of $\mathbf{A}$ generate the rotations of a space with one dimension more, SO($D+1$). Fock found this by carrying momentum space onto a sphere one dimension up (Fock 1935), Alliluev carried the construction to D dimensions (Alliluev 1958), and Bander and Itzykson worked out the group theory (Bander and Itzykson 1966). Why it happens only for $1/r$ can be put in one line. The Schrödinger equation in momentum space contains the potential's Fourier transform. For $1/r^{\eta}$ in D dimensions that is $1/k^{D-\eta}$, which for $\eta = 1$ is $1/k^{D-1}$. Carried onto the sphere, $1/k^{D-1}$ becomes, up to factors the projection itself supplies, Gauss's potential of a point charge in $D+1$ dimensions, and the natural functions of that are the sphere's own harmonics. For any other η the transform carries over to something else, and the larger symmetry is lost.

The symmetry does the work. States that the larger rotations carry into one another must have the same energy. A level of hydrogen is the set of degree-$(n-1)$ harmonics on that sphere, and the ordinary rotations split this set into $l = 0, 1, \ldots, n-1$. So every l below n shares the energy, which in four dimensions is $-Z^2/2(n+\frac{1}{2})^2$. With $(l+1)^2$ orbitals to each l, shell n holds $\sum_{l<n} 2(l+1)^2$ electrons: 2, 10, 28, 60. That degeneracy is what makes a shell, a group of levels with a gap above it.

Without the symmetry nothing holds the l together. On the same solver, 2p sits −6.2 % from 2s at $\eta = 0.8$, 5×10^{-10} % at $\eta = 1$, +15.4 % at $\eta = 1.2$ and +64.0 % at $\eta = 1.5$ (as a fraction of the shell's binding; 3d against 3s goes the same way, +78.3 % at $\eta = 1.5$; figure 1b). Below 1 the electron with more angular momentum is the more bound, above it the less. So under $\eta = 1.5$ a one-electron atom's 2p is bound little more than a third as hard as its 2s, and each l is a level of its own. A table would still have rows, since many electrons split the l in ours too. But the rows would be set by how the exponent treats electrons that come near the nucleus, and would have to be found again for every η. The generalized Madelung rule, which starts from hydrogen's degeneracy, wouldn't apply. Under $1/r$ the rows come from counting harmonics, and that is what lets part III tell the geometry's share of the table from the law's.

The literature on hydrogen in D dimensions (Nieto 1979; Al-Jaber 1998) and on dimensional scaling (Herschbach, Avery, and Goscinski 1993) takes $1/r$ in every D, mostly as a convention that keeps the problem solvable. Négadi and Kibler's generalized Madelung rule (1996), which gives the four-dimensional rows when it is applied to $D = 4$, builds on the same D-dimensional hydrogen atom. That is why the convention is more than one: it is the only power law under which the question "what are the shells?" has the answer the rule needs.

F.4 No Gauss's law for 1/r. In three dimensions $\nabla^2(1/r) = 0$ away from the charge, and that one fact is Gauss's law. In D dimensions $\nabla^2 r^{-\eta} = \eta(\eta+2-D)\,r^{-\eta-2}$, which vanishes only for Gauss's own $\eta = D-2$. In four,

$$\nabla^2 \frac{1}{r} = -\frac{1}{r^3} \qquad (r \neq 0),$$

which the solver's finite differences reproduce (−1.00000 at $r = 1$, −0.12500 at $r = 2$). By the divergence theorem the flux of the field through a sphere round a charge is the integral of $-\nabla^2(1/r)$ over the ball inside it. The integrand is $1/r^3$ and the ball's volume element $2\pi^2 r^3\, dr$, so the flux grows linearly with the sphere's radius, as if empty space were charged with a density $\propto 1/r^3$. There is no Gauss's law for it.

F.5 Five field theories that give 1/r, and the one taken. A potential is what a field theory's static equation gives for a point charge, and the equation is simplest in Fourier space, where each pair of derivatives becomes a factor k^2 of the wavenumber k. Maxwell's statics give a point charge the potential $\varphi \propto 1/k^2$, and that transforms back to $1/r^{D-2}$ in space, so $1/r^2$ in four dimensions. A potential $1/r^\eta$ in D dimensions is the transform of $1/k^{D-\eta}$, so the pinned $1/r$ needs $1/k^3$. There are five ways to get it (the site's force chapter, four.mino.mobi/atom/force.html, works through each). Four fail something we ask of the law, and each failure is stated with what it assumes. The third survives with a flaw that can be removed, and it is the one this paper takes.

1. *A local, relativistic field:* impossible for a quadratic field equation with finitely many derivatives. Such an equation is a polynomial in k^2, a whole number of pairs of derivatives, and $k^3 = (k^2)^{3/2}$ isn't one. (A non-local kernel can arise from integrating out other fields; that is route 2 or 3.)
2. *Relativistic but non-local:* the potential obeys $(k^2 - \omega^2/c^2)^{3/2}\,\varphi \propto \rho$, Maxwell's wave operator raised to the power 3/2, which in space and time is an integral over all of both. This is the non-local electrodynamics $FD^{s-2}F$ of Heydeman et al. (2020) with $s = 3$ in $d = 5$ spacetime dimensions, a case their analysis covers ($d = s + 2 \geq 5$). There the gauge-invariant field strength has scaling dimension 2, below the 3 that unitarity requires of it in a conformal theory, and they expect time evolution not to be unitary: states of negative probability. Fractional and non-local gauge theories are not non-unitary as a class (Marino et al. 2014; La Nave, Limtragool, and Phillips 2019); it is this one, in the framework where the bound applies, that fails.
3. *A vacuum whose permittivity grows with wavenumber,* $\varepsilon_0\, \ell\, k$ with ℓ a length. The static equation becomes $\varepsilon_0\, \ell\, k \cdot k^2 \varphi = \rho$, which is $1/k^3$, the pinned $1/r$. With the permeability left at μ_0, light obeys $k^2 = \ell\, k\, \omega^2/c^2$, so $\omega = c\sqrt{k/\ell}$. It disperses like deep-water waves, and there is no speed of light. A permittivity rising as a power of k, and the fractional Poisson equation it gives, is known from media with power-law spatial dispersion (Tarasov and Trujillo 2013; Tarasov 2016); their power $\alpha = 3$ without screening is this one, applied here in four dimensions. This is the route we take, and the flaw is removed below.
4. *A nonlinear vacuum,* $\nabla \cdot (|\mathbf{E}|^{\beta-2}\mathbf{E}) = \rho$. It keeps a Gauss's law, for the flux of $|\mathbf{E}|^{\beta-2}\mathbf{E}$ instead of $\mathbf{E}$. A point charge then has $|\mathbf{E}|^{\beta-1} r^{D-1}$ constant, and a $1/r$ potential ($|\mathbf{E}| \propto 1/r^2$) needs $\beta = (D+1)/2 = 2.5$, the p-Laplacian with $p = 5/2$, whose point solution is exactly that (Peter Lindqvist 2017). It is a consistent theory, but fields no longer add: the field of two charges isn't the sum of each one's, so there is no pair potential, which an atom's Hamiltonian as written here needs.
5. *A small fourth dimension:* $1/r$ far away, but then atoms are three-dimensional (section 2.2).

Route 1 doesn't exist under its assumptions, route 2 has states of negative probability in the framework where that can be shown, route 4 gives up a pair potential, and route 5 gives back three-dimensional atoms. That leaves route 3. This is a choice among the routes found, not a proof that no other theory gives the law. The permittivity fixes the statics, and the statics are all that the pinned law asks of it. The permeability is still free, and it decides only how fields move. Left at μ_0 it gives the dispersive light above. We choose $\mu_0/\ell\, k$, and call the result the **lens vacuum**. The product of permittivity and permeability is then $\varepsilon_0\mu_0$ at every wavenumber, so light keeps the one speed c. The fields $\mathbf{D}/\varepsilon_0$ and $\mu_0\mathbf{H}$ obey Maxwell's equations of empty four-dimensional space exactly, so light is ordinary four-dimensional light at every wavelength, and the energy is never negative. What pushes a charge is $\mathbf{E} = (\mathbf{D}/\varepsilon_0)/\ell\, k$, that light seen through a lens. Dividing by k in Fourier space is, in space, spreading over a kernel $1/r^3$ (the transform of $1/k$ in four dimensions), so $\mathbf{E}$ at a point is a blur of $\mathbf{D}$ over all of space. Its statics are $1/\ell\, k^3$, the pinned $1/r$. The price is relativity, for the force between charges only. The blur acts across all of space at once, so there is an absolute "now", and a charge feels another's change instantly, a little, everywhere. We have not found this pair of constitutive laws in the literature we read. Light at one speed and an energy that is never negative do not by themselves make the lens vacuum a causal or quantum theory; that is for the later paper.

This paper uses only route 3's statics: charges at rest and the $1/r$ between them, which is all an atom's ground state needs. Every choice of permeability gives the same statics, so nothing here depends on that choice, on ℓ, on how the lens vacuum's fields move and radiate, or on the absolute now. Those dynamics, of the same lens vacuum, are the subject of a later paper.

F.6 Electrons: what is derived and what is declared; nuclei and the fine-structure constant. An electron's spin is a representation of the rotations, and in four dimensions they split: so(4) = so(3) $\oplus$ so(3), a turn's left half and right half. The spinors of Spin(4) = SU(2) × SU(2) come as $(\frac{1}{2}, 0)$, two states that turn with the left half only, $(0, \frac{1}{2})$, two that turn with the right, and their sum, four. The smallest have two states. The free electron can keep its relativity, which the lens vacuum gives up only for the force between charges. Its Dirac equation in 4 + 1 dimensions has four components, and the product of the four spatial Dirac matrices squares to one and anticom-

mutes with each of them, so it serves as time's (the site's self-test builds them). A slow electron keeps the two components on which that matrix is +1, and they carry one half of the rotations only: two spin states, and a handed spin that a right-isoclinic turn leaves alone (four.mino.mobi/atom/motion.html#spin). A four-state electron, $(\frac{1}{2},0)\oplus(0,\frac{1}{2})$, would need two such Dirac fields with the same charge and mass, two species in all but name. It would hold four to an orbital: the hydrogen shells would hold 4, 20, 56 and 120, every row of the table would be twice as long, and the noble gases would sit elsewhere. The vector $(\frac{1}{2},\frac{1}{2})$ also has four states, but a full turn leaves it unchanged; it is integer spin, not an electron's. So two states is what the smallest relativistic electron gives, and that is the choice made. Two things remain declared. One is that the electron is that smallest spinor. The other is Pauli's exclusion: in ours the spin-statistics theorem derives it from relativity and locality together, and the force here is neither, so the theorem can't be borrowed. The exclusion is what the proofs of stability in ours lean on (Dyson and Lenard 1967; Lieb and Thirring 1975), so without it bulk matter would have no claim to stability here either. Nuclei are point charges $+Z$, stable for every Z to 140 (declared: nuclear physics in four dimensions is another paper). The fine-structure constant is small, so atoms are solved without relativity (section 8). No other long-range field acts on charges.

F.7 Stability: shown for single atoms, open for matter in bulk. In ours it is a theorem that N atoms have an energy bounded below by a constant times N (Dyson and Lenard 1967; Lieb and Thirring 1975), whatever the nuclei's positions. The usual proofs lean on the shell theorem, which fails here. That doesn't close the question: decompositions of radial interactions that need no shell theorem exist in any dimension (Hainzl and Seiringer 2002), and Riesz gases, of which a $1/r$ law in four dimensions is one, have their own mathematics (Lewin 2022), though neither is a proof for electrons and nuclei here. So we give evidence, not a proof, and the evidence is narrower than the theorem. Single atoms don't collapse. Thomas–Fermi scaling gives total energies growing as $Z^{3-2/D}$, $Z^{7/3}$ in ours and $Z^{5/2}$ in four dimensions (Kventsel and Katriel 1981), and the solved table follows it, with a fitted exponent of 2.527 over Z = 10–140 (ours: 2.387 over 10–36; both a little above the asymptotic value, as finite-Z corrections make them). Separated neutral atoms don't run away either. Every force between them here (the permanent $-3Q_AQ_B/64R^5$ of section 7, dispersion's R^{-6}) falls faster than R^{-4}, so summed over four-dimensional space each atom's share of the far field is finite. That says nothing about atoms that overlap, trade charge or crowd together, which is where a proof of stability has its work. Stability of matter here is open.

G · The methods in full

G.1 The code. Everything in this paper is computed by code written for it: JavaScript with no outside libraries, run under Node.js, and the same files run in a browser (four.mino.mobi/atom). The atom is `atom/js/atom.js`, with the shell kernel in `kernel.js`. One electron in any law is `papers/p1/laws.js`, and the terms of open subshells are `atom/tools/hund.mjs` and `termtable.mjs`. Nothing is taken from a quantum-chemistry package. Two independent codes check it (appendix E.2): `atom/js/gauss.js`, Hartree–Fock in a Gaussian basis (Boys 1950) for any D, with the integrals of McMurchie and Davidson (1978), and in ours PySCF (Sun et al. 2018, 2020). Every number quoted is read from the code's output by the paper's build (appendix K).

G.2 The law inside an atom. An atom here is a point nucleus of charge Z and Z electrons, non-relativistic, with every pair of charges interacting through the pinned potential q_1q_2/r in four spatial dimensions. The same law acts between an electron and the nucleus and between two electrons; nothing in the method borrows Gauss's law. Atomic units throughout (hartree, bohr).

Each electron moves in the field of the nucleus and of the other electrons' clouds, averaged over directions, so the problem is radial. In D dimensions a wavefunction $\psi = r^{-(D-1)/2}\,u(r)\,Y$, with Y a harmonic of degree l on the $(D-1)$-sphere, has a radial function u that obeys

$$-\tfrac{1}{2}u'' + \left[\frac{L(L+1)}{2r^2} + V(r)\right]u = \varepsilon\,u, \qquad L = l + \frac{D-3}{2},$$

so four dimensions differ from ours by the half-integer shift of L ($L = l + \frac{1}{2}$) and by the degeneracies: a subshell of angular momentum l holds $2(2l+1)$ electrons in ours and $2(l+1)^2$ in four dimensions. With $V = -Z/r$ the levels are $-Z^2/2(n+(D-3)/2)^2$, so 4D hydrogen's ground state is $-2/9$ hartree; the solver gives -0.22222 (appendix E.1).

G.3 The clouds' potential. A spherical shell of charge at radius s, felt at radius r, gives in ours $1/\max(r,s)$ (the shell theorem). In four dimensions under $1/r$ it gives $\kappa(t)/\max(r,s)$, $t = \min/\max$, with κ the $1/r$ kernel averaged over a 3-sphere,

$$\kappa(t) = \frac{2}{\pi}\int_0^\pi \frac{\sin^2\theta\,d\theta}{\sqrt{1+t^2-2t\cos\theta}},$$

which falls from $\kappa(0) = 1$ to $\kappa(1) = 8/3\pi = 0.848826$ (appendix A; Simpson's rule on 400 intervals, then a table of 2001 points with linear interpolation). Every electron–electron term in the self-consistent field, its energy and the ionized atom's goes through this kernel. That is where the missing shell theorem enters the atom: an inner cloud screens the nucleus from an outer electron only partly.

G.4 Exchange. Local exchange from the uniform electron gas of the same dimension and the same law, $\varepsilon_x(\rho) = -A_D\,\rho^{1/D}$ per electron for an electron density ρ, with $A_3 = \frac{3}{4}(3/\pi)^{1/3} = 0.738559$ and $A_4 = 16/15\sqrt{\pi} = 0.601802$, as in the general-$D$ formulas of Loos and Gill (2016) for $1/r$ jellium (appendix B). Spins are polarized by Hund's rule (a subshell fills one spin first), and each electron's own share is removed from both its

Hartree and its exchange potential: the Hartree term by its own orbital's cloud, the exchange term by the fully polarized exchange of that orbital alone, in the manner of the self-interaction correction of Perdew and Zunger (1981). There is no correlation term. That is an omission, not a gap in what is known: the arbitrary-dimensional exchange is old (Glasser and Boersma 1983), and the correlation of the $1/r$ electron gas has been computed in any dimension, four included, by the random-phase approximation (Schlesier, Benavides-Riveros, and Marques 2020) and beyond it (Duc Pham et al. 2023). We have not implemented or tested a four-dimensional correlation functional; the later paper's numbers carry a correction to the earlier one's coefficients that would have to be settled first.

G.5 Solving one configuration. The radial equation is solved on a logarithmic grid of 700 points from $r_0 = 10^{-5}$ to 150 bohr (step $h = 0.02364$ in $\ln r$) by Numerov's method on $y = u/\sqrt{r}$, with each level found by Sturm bisection on the node count of the outward solution (90 halvings), the outward integration stopped where $h^2 f/12 > \frac{1}{2}$ (for $y'' = fy$) so that Numerov cannot fake nodes in the forbidden region, and the orbital matched to an inward solution at the turning point.

For a fixed configuration (occupations of each subshell, every orbital of a subshell sharing one radial function) the field is iterated to self-consistency: each orbital solved in the potential of the nucleus, the others' clouds and exchange; the new clouds mixed into the old at 0.4; up to 80 iterations, stopping when the total energy changes by less than 10^{-7} hartree between iterations.

G.6 The stop, against one 10^5 times tighter. A stop on the energy can leave the far density less converged than the energy is; the plane-wave work met exactly that in its pseudopotentials' all-electron atom, where exchange carried an unconverged tail into the potential. So 19 atoms across every row of both worlds were re-solved in their baked configurations and then driven on from that field to a stop of 10^{-12} hartree (appendix C, `audit-scf.mjs`). Total energies and first ionization energies moved by at most 2.7×10^{-7} and 2.8×10^{-7} hartree, the outermost orbital energy by 1.7×10^{-6}, and $Q = \sum \langle r^2 \rangle$ (the far pull's, and the forces between atoms') by 1.9×10^{-5} of itself; the baked $\langle r^2 \rangle$ agree with the tight ones to 7.0×10^{-5}. The density beyond about 15 bohr is not converged relatively, but it is noise there: off by large factors only where it is 10^{-13} to 10^{-17} electrons per bohr, and converged to 10^{-6} where an atom has a loosely held s electron that puts real charge out that far. Nothing the paper quotes depends on it; a calculation that builds a potential from the far density through exchange must converge the density pointwise, as the pseudopotentials now do.

G.7 The radial grid, against six others. 11 atoms from every row of both worlds were re-solved in their baked configurations on six other grids, changing one of the number of points (500, 1000), the inner end (10^{-4}, 10^{-6} bohr) and the outer end (100, 250 bohr) at a time (appendix C, `convergence.mjs`). First ionization energies moved by at most 1.4×10^{-4} hartree on any of them. Total energies moved by at most 6.7×10^{-2} hartree, 2.4×10^{-5} of themselves, the largest for our own krypton, whose 1s sits deeper than any four-dimensional atom's and is the grid's hardest case. The paper quotes ionization energies and energy differences, not total energies.

G.8 The ground configuration. For each Z the ground configuration is searched, not assumed. The search starts from the hydrogenic filling (subshells in order of n, then l), solves it, and refills by the orbital energies of every subshell, occupied or not, in the field it has just found, up to 6 rounds or until the filling stops changing. It then tries moving one or two electrons between the open subshells, the two highest filled ones and the two lowest empty ones, solving each candidate from the current field. A candidate counts only if its field settles: an unsettled field can report an energy far too low, since with exchange a barely bound orbital can drift for hundreds of iterations. The lowest settled energy is the ground state.

For 7 elements of the 4D table (88, 101–103 and 105–107) the first search landed on an unsettled field. They were searched again seeded from both neighbours' configurations with one or two electrons added or taken, each seed solved from scratch, and the lowest settled one kept.

The first ionization energy is ΔSCF, a difference of two self-consistent energies: the ion's own ground state with an electron taken from whichever of the 3 least-bound occupied subshells leaves it lowest, relaxed from the atom's field, minus the atom's energy. Koopmans' orbital energies are not used.

The table is 140 elements in four dimensions and 36 in ours, baked once (on GitHub's runners past $Z \approx 80$) into a single generated file, which a selftest re-solves in sample on every change.

G.9 The terms of open subshells. The search above gives every open subshell Hund spins. Section 5 finds where that's wrong, and the table is corrected for it. Each d- and f-block element is searched again, seeded as the repair pass seeds, and every settled candidate's energy gets its open subshells' term correction: the lowest term less the lowest maximum-spin term. That comes from the subshell's own repulsion on that candidate's own orbital (`atom/tools/termtable.mjs`). It is zero wherever Hund's first rule holds. The ionization is corrected the same way on both sides. Fillings whose spin blocks are too large to solve per candidate take the survey's verdict (section 5.1), which for them is that the rule holds.

G.10 The counterfactual. To separate what the table owes to the law from what it owes to the geometry, every 4D element is solved a second time with the shell theorem forced on: the clouds' potential taken as $1/\max(r, s)$ (our kernel) while everything else, the 4D kinetic operator, the degeneracies, exchange, the search, stays as it was. Each element's search is seeded with the real table's configuration and its neighbours' ± one electron, so the counterfactual isn't a weaker search; and the real law is re-solved in each counterfactual configuration, to check that the real table's configuration is still the lower one under the law. Where no candidate's field settled, a sec-

ond, wider search started from every configuration in reach (the real table's for the element and its neighbours ± one electron, the unsettled filling, and the counterfactual's own neighbours ± one), each solved from scratch with the clouds mixed at 0.15 for up to 1500 iterations. A counterfactual atom counts as settled only if its field converged and its first ionization energy is positive (a negative one means an electron would rather leave: the search missed a lower configuration). By that rule 112 of 140 settle, every atom up to 88; the rest (89–116) are row 6's f block and the first elements after it.

G.11 One electron, any law. The arguments of section 2 use a separate solver for one electron in any central potential (`laws.js`): the same radial equation and Numerov–Sturm method on a grid whose inner end can be moved, which is what tests a fall to the centre. For $V = -\alpha/r^2$ it finds no level below $\alpha = \frac{1}{2}$ on any grid and, above it, a lowest level that scales as $1/r_0^2$ (section 2.1); for $V = -1/r^n$ it gives the splittings within each shell (section 2.3). Its check is 4D hydrogen: −0.22222222 against −2/9.

G.12 A second method: Hartree–Fock. To see which configurations depend on the choice of method, every element is solved a second way, by numerical Hartree–Fock (`hf.js`): one radial function per subshell on its own logarithmic grid, exact exchange instead of local exchange, and the energy the average over the subshells' maximum-spin arrangements, to which each open subshell's whole term correction (its lowest term less that average, appendix G.9's machinery on the Hartree–Fock orbitals) is added. Exchange between subshells is expanded in the Gegenbauer coefficients of $1/|\mathbf{x}-\mathbf{y}|$, each radial kernel a convolution on the logarithmic grid; in four dimensions the coefficients are tabulated by quadrature, a_0 being the shell kernel κ. The candidates are the local-exchange search's settled configurations within 4 eV of its ground, so both methods rank the same set. For row 6's f block and our lanthanides, where local exchange's bias toward f can push the right configuration out of that window, the window is 10 eV, and Hartree–Fock then climbs from its own ground through every configuration one electron away, those more than 6 eV up ranked without the term correction. A configuration with an empty subshell below an occupied one of the same l (sodium's $4s^1$ with $3s$ empty) is left out of the Hartree–Fock ranking: the variational solve drops the electron into the empty lower subshell and returns the compact configuration's energy under the other name. The energies carry an $O(h^2)$ error from the kernel's kink at $r = r'$, so absolute energies are extrapolated from steps 0.02 and 0.01; differences converge at 0.02, where the ranking is done. Checks (block 71): our He, Ne, N ^{4}S, C ^{3}P and Fe ^{5}D against the numerical Hartree–Fock limits to 2×10^{-5} hartree, and four-dimensional closed shells against the Gaussian code.

G.13 A known-answer test for every method. Every method above has a known-answer test in the selftest, run on every change: hydrogen in both worlds (block 1), the radial solver (2), the kernel against direct integration (3), exchange against the electron-gas constants (4, 18), the ground search (5, 7), two-electron atoms against an independent Gaussian Hartree–Fock code in four dimensions and MP2 against PySCF in ours (17, 43), Hartree–Fock against its known limits (71), our table against measured values (6: 32 of 36 configurations; ionization to a 4.3 % median error), and this paper's own claims recomputed from the bake (67).

H · Hund's first rule in full

H.1 Every term, from the open subshell's determinants. For each open subshell of each ground configuration, its orbitals are the degree-l harmonics on the 3-sphere, $(l+1)^2$ of them (4 for p, 9 for d, 16 for f). The repulsion between two of its electrons is the law's $1/|x-y|$, projected exactly onto the harmonics of degree 0 to $2l$, with the radial parts integrated over the atom's own orbital. Every arrangement of the subshell's electrons with a given M_S is set up and the lowest energy found (Lanczos, matrix-free; a 4D d^9 block has 15 876 arrangements). A term of spin S appears in every block with $M_S \le S$, so the ground term's spin is the largest M_S whose block still reaches the lowest energy. In ours this reproduces Racah's term separations for d^2 and d^3 to the wavenumber, and the known ground terms of Cr^{3+}, Mn^{2+}, Pr^{3+} and Gd^{3+}. Over all 23 open subshells of our table, maximum spin is lowest, as it should be.

H.2 How complete the f survey is. No other f filling was found to break the rule, but the search is not complete. For f^2, f^4, f^5, f^{28}, f^{30}, f^{31} every spin was checked. For f^6–f^{25} only the highest two or three were, because the lower-spin blocks run to 10^6–10^8 arrangements; there the result is a bound, no lower term among the spins that could be run. Two fillings (elements 102 and 106) didn't finish in the runner's time.

H.3 The singlet's symmetry, and what the calculation establishes. Four-dimensional d orbitals form the representation (1,1) of $SO(4) \cong SU(2) \times SU(2)/\mathbb{Z}_2$. Two electrons in them make spatially symmetric states (2,2) ⊕ (2,0) ⊕ (0,2) ⊕ (1,1) ⊕ (0,0), all spin singlets (25, 10, 9 and 1 states), and antisymmetric ones (2,1) ⊕ (1,2) ⊕ (1,0) ⊕ (0,1), triplets (30 and 6). The singlet that comes lowest is the ten-state (2,0) ⊕ (0,2): two irreducible representations of SO(4), exchanged by a reflection, so a single level once reflections are counted. In ours the triplet ^{3}F is the term that keeps the electrons nearest right angles (0.225, against ^{1}D's 0.292 and random's 0.333), and it is the lowest for every value of F^4/F^2. Titanium's d^2 has ^{3}F lowest and its first singlet 1.25 eV up.

What this establishes is narrower than a whole-atom ground term. The term energies are computed within one subshell on the atom's own maximum-spin orbital, which is neither relaxed term by term nor correlated beyond the subshell, and the correction they give the table (section 5.3) is first order. The inversion holds under both laws tested, and for any orbital whose F^4/F^2 is below 5/3. Hund's first rule has been studied away from three dimensions, in two-dimensional helium-like atoms and quantum dots (Sako et al. 2012; Sako, Paldus, and Diercksen 2010) and by dimensional scaling (González 1991),

and two-electron matrix elements for hydrogenic d^2 and f^2 pairs have been derived for any D (Summerfield and Loeser 1999). We have not been able to read the last two in full, and none of the sources we could read states this inversion.

H.4 The corrected table. Every d- and f-block element (92 of them) was searched again, with the neighbours' configurations as extra seeds (49–65 settled candidates each). Every candidate got its open subshells' gaps added, and the candidates were ranked again, with and without the correction, so that a change from the wider search is told apart from one from the terms. **The terms change no ground configuration.** They lower the eight Hund-breaking atoms' energies, and they move first ionizations in pairs. An atom whose own ground term gains the gap holds its electron harder: 54 (6.41 → 6.71), 66 (18.82 → 19.40), 114 (6.92 → 7.27), 126 (24.92 → 25.55) eV. The next element, whose ion lands on the broken filling, gives one up more easily: 55 (8.28 → 7.91), 67 (20.56 → 19.93), 115 (9.28 → 8.83), 127 (27.23 → 26.56) eV. That is 0.30–0.67 eV either way. The wider search found one configuration the bake had missed. Element 82 is $4f^1\ 5d^1\ 6s^2$, 0.36 eV below the bake's $4f^2\ 6s^2$, which is our cerium's kind; its ionization goes 7.03 → 7.39 eV. The table of section 4 is the corrected one (`data/table4.csv`). The Hund-spin table it replaces is supplement S1 (`data/table4-hundspin.csv`). How far to trust each configuration follows from appendix E.3. Take a bias of about a volt toward d and f: a ground configuration with a competitor holding fewer d and f electrons within a volt of it could be the other way round. 11 of the 92 d- and f-block elements have one: 30 ($3d^6\ 4s^2$, 0.55 eV up), 37 ($3d^{13}\ 4s^2$, 0.55 eV up), 53 ($5s^2\ 5p^1$, 0.01 eV up), 58 ($4d^6\ 5s^2$, 0.20 eV up), 61 ($4d^9\ 5s^2$, 0.25 eV up), 64 ($4d^{13}\ 5s^1$, 0.95 eV up), 107 ($5p^8\ 6s^2$, 0.20 eV up), 108 ($5p^8\ 6s^2$, 0.79 eV up), 109 ($5p^8\ 6s^2$, 0.57 eV up), 122 ($5d^{10}\ 6s^2$, 0.54 eV up), 124 ($5d^{13}\ 6s^1$, 0.55 eV up).

H.5 The d^2 singlet in a ligand field. A spin singlet is not orbitally inert: the ten states of (2,0) ⊕ (0,2) still carry orbital structure, and in a compound a ligand field splits them, and might hand the ground to a triplet. The ion here is element 26's 2+, $3d^2$ over closed shells, with point charges −1 at 4 bohr in three arrangements of four dimensions: along the axes (eight, the 16-cell, four dimensions' octahedron), at a tesseract's corners (sixteen), and at the 24-cell's vertices (twenty-four). The field is projected exactly onto the degrees a d × d product holds, scaled from zero, and every determinant of the two electrons is diagonalized, as in section 5.1. A four-dimensional d subshell splits 6 + 3 on the axes and 3 + 6 at the corners (ours' octahedron 3 + 2, cube 2 + 3). On the axes the two electrons go into the six-fold set, and the singlet stays the ground at every field strength: its ten states split, the lowest of them 2-fold, an orbital doublet that a distortion could act on, and the lowest triplet stays 0.44 eV up, the free ion's gap. At the corners the three-fold set is lowest, and two electrons in it favour a triplet: the singlet holds until the one-electron spread Δ reaches 2.4–2.5 eV, and a triplet is the ground beyond, but only just, 0.07 eV below the singlet even at Δ = 7.6 eV. The 24-cell splits the d subshell not at all (none): its vertices average every harmonic of degree 2 and 4 to zero, as the icosahedron does for ours, so a d ion in that site keeps its free-ion terms. Ours' Ti^{2+} ($3d^2$) is the check, with its triplet the ground at every field in both shapes, as Tanabe and Sugano (1954) found for d^2. These are point charges, a model of the field's symmetry rather than of its size (`atom/tools/ligandd2.mjs`, `data/ligandd2.json`).

I · Starting data for chemistry

I.1 Valence states. Chemistry can pay to move an electron if the bonds it then makes pay more, so the reference records the cost. For each of 11 representative atoms, every configuration one electron away from the ground (any electron near the top moved to any subshell near the top) was solved by local exchange and by Hartree–Fock (table 1; all 11 with the ground's spin ladder in `data/valence.json`). Ours is the calibration, 33 states against NIST's levels (Sansonetti and Martin 2005). Moving an electron into a higher shell of the kind that makes a Rydberg series is good to a median 0.4 and 0.3 eV (18 states). The s → p promotion, the price of hybrid bonding, is not. For atoms whose p shell is less than half full (Be, B, C, Mg, Al, Si) both methods make it too cheap, by −2.3 to −0.6 (local exchange) and −1.7 to −0.9 (Hartree–Fock) eV, the near-degeneracy of an s^2 pair with p^2 that neither method holds; past half full (N, P, S) they make it too dear (−0.2 to 1.0 and 0.3 to 2.7 eV), where the measured state mixes with others. In four dimensions the p subshell holds eight, so every second-row atom up to element 8 is in the first group: element 4's 2s → 2p costs 1.3 and 1.5 eV by the two methods, darbon's (element 7) 3.4 and 4.0 eV, element 14's 3s → 3p 3.1 and 2.9 eV, and on our evidence each is probably a volt or two short of the true cost. Element 26's three lowest moves (into 4p, from 4s into 3d, from 3d into 4p) lie within 0.2 eV of each other by Hartree–Fock and a volt apart by local exchange, so its chemistry has several states to draw on and the methods disagree about which is cheapest. For element 53 Hartree–Fock finds a state below the table's ground, the open configuration of appendix E.6.

Table 1. Valence states of representative four-dimensional atoms: the ground configuration (outer subshells) and the three lowest configurations one electron away by Hartree–Fock, each labelled by the move and given by local exchange and Hartree–Fock (eV above each method's own ground; a negative value is a state below the table's ground).

element	ground	lowest states by Hartree–Fock: move, local exchange / Hartree–Fock (eV)
4	$1s^2\,2s^2$	2s → 2p 1.3 / 1.5; 2s → 3s 3.4 / 3.4
7	$1s^2\,2s^2\,2p^3$	2s → 2p 3.4 / 4.0; 2p → 3s 6.1 / 5.5; 2p → 3p 7.2 / 6.7
10	$1s^2\,2s^2\,2p^6$	2p → 3s 8.9 / 8.2; 2p → 3p 10.8 / 10.0; 2s → 2p 14.0 / 17.4
12	$1s^2\,2s^2\,2p^8$	2p → 3s 13.7 / 12.9
14	$2s^2\,2p^8\,3s^2$	3s → 3p 3.1 / 2.9; 3s → 4s 4.3 / 4.2
26	$3p^8\,3d^2\,4s^2$	4s → 4p 3.3 / 3.1; 4s → 3d 2.3 / 3.1; 3d → 4p 4.3 / 3.3
39	$3p^8\,3d^{16}\,4s^1$	3d → 4s 2.5 / 0.4; 4s → 3d 3.2 / 5.0; 4s → 4p 7.8 / 7.1
42	$3p^8\,3d^{18}\,4s^2$	4s → 4p 9.3 / 8.3; 4s → 5s 11.1 / 9.9; 4s → 4d 13.0 / 11.8
53	$4p^8\,4d^1\,5s^2$	4d → 5p 0.0 / −0.8; 4d → 6s 1.8 / 1.1; 4d → 4f 3.8 / 2.9
82	$5p^8\,5d^1\,6s^2$	4f → 5d 4.7 / 1.5; 5d → 4f 0.4 / 1.8; 5d → 6p 2.5 / 1.9
128	$5s^2\,5p^8\,5d^{18}$	5d → 6s 8.0 / 8.6

I.2 Darbon's hydride holds five hydrogens. The table implies a valence rule: an atom with v electrons outside its core makes v bonds, up to $D + 1$ of them, and $2(D + 1) - v$ past that. That is a claim about molecules, and one hydride tests it directly. Darbon (element 7, $2s^2 2p^3$) should make five bonds, so DH_5 should exist, its hydrogens at the corners of the 5-cell, the simplest regular figure in four dimensions. Making five bonds takes the s → p promotion of appendix I.1. At that geometry (D–H 3.23 bohr, by Hartree–Fock) we computed the whole curvature matrix of the energy by central differences, in all 24 coordinates of six atoms in four dimensions. With the 10 translations and rotations set apart, the 14 curvatures fall in groups of 5 + 4 + 1 + 4, as the 5-cell's symmetry requires. None is negative: the lowest is 0.020 hartree/bohr², and the gradient is 5×10^{-6}. So DH_5 is a true minimum, not a saddle that a symmetric search could settle on. It also lies below both balanced ways of coming apart. Losing one hydrogen, $DH_5 \rightarrow DH_4 + H$, costs 1.55 eV by Hartree–Fock and 2.32 eV with MP2. Losing two as a molecule, $DH_5 \rightarrow DH_3 + H_2$, costs 2.13 and 3.43 eV. Each fragment is in its lowest spin: DH_4 is a doublet and DH_3 a triplet, with DH_3's singlet 1.01 (Hartree–Fock) and 0.85 eV (MP2) higher. Energies are by unrestricted Hartree–Fock and MP2 in the molecular basis. The fragments' geometries were optimized only within symmetric shapes, with their hydrogens equivalent; DH_3 and DH_4 settle flat, their hydrogens in a plane or a 3-space through darbon. A lower, less symmetric fragment could lower both costs, and the curvature matrix, which describes DH_5 only, says nothing about that. The systematic test, every hydride and its fragments searched without symmetry, belongs to the chemistry this table starts.

J · The atoms' response, and joining and separating them

J.1 Polarizability, dispersion and the permanent pull. For the closed-shell atoms and the one-s-electron alkalis of both worlds (table 2) the static polarizability $\alpha_p(0)$, the polarizability at imaginary frequency and from it the dispersion coefficient C_6 by Casimir and Polder's integral are computed by the coupled response of the same solver (Hartree plus local exchange, each electron's own share removed; exact for hydrogen in both worlds). The set is the atoms for which that response's angular bookkeeping is exact: full subshells and a single s electron. Doubling the frequency quadrature moves C_6 by at most 2×10^{-6}, and the other radial grids move $\alpha_p(0)$ and C_6 by at most 0.5 %. The sum rule $\omega^2 \alpha_p(i\omega) \rightarrow Z$, read at ω = 300 hartree, holds to 0.90 or better; the shortfall in the heaviest atoms is core response at higher frequencies still. In ours, against the reference values of Derevianko, Porsev, and Babb (2010), the noble gases' $\alpha_p(0)$ is off by −8 % to +4 %, and the alkalis and alkaline earths are too easily polarized, by +7 % to +25 % in $\alpha_p(0)$ and +11 % to +35 % in C_6 (median errors 9 % and 13 %), as local exchange without correlation is known to make them.

Table 2. Response of closed-shell and one-s-electron atoms (atomic units): static polarizability, C_6, $Q = \sum \langle r^2 \rangle$, in four dimensions $C_5 = 3Q^2/64$ and $R^* = C_6/C_5$; for ours, beside each, the reference value (ref.) of Derevianko, Porsev, and Babb (2010). Data: `data/response.json`, with $\alpha_p(i\omega)$ on the quadrature nodes and each atom's convergence.

world	element	outer subshells	$\alpha_p(0)$	ref.	C_6	ref.	Q	C_5	R^*
ours	1	$1s^1$	4.5	4.5	6.5	6.5	3	0	
ours	2	$1s^2$	1.32	1.38	1.38	1.46	2.37	0	
ours	3	$1s^2\,2s^1$	184	164	1 600	1 389	18.8	0	
ours	4	$1s^2\,2s^2$	47.1	37.8	288	214	17.5	0	
ours	10	$2s^2\,2p^6$	2.45	2.67	5.68	6.38	9.51	0	
ours	11	$2p^6\,3s^1$	174	163	1 722	1 556	26.3	0	
ours	12	$2p^6\,3s^2$	78.6	71.3	721	627	29	0	
ours	18	$3s^2\,3p^6$	11.4	11.1	66.4	64.3	26.5	0	
ours	19	$3p^6\,4s^1$	357	290	5 215	3 897	49.1	0	
ours	20	$3p^6\,4s^2$	171	157	2 442	2 121	54.7	0	
ours	30	$3d^{10}\,4s^2$	45.5		338		33.7	0	
ours	36	$4s^2\,4p^6$	17.5	16.8	136	130	40.1	0	
4D	1	$1s^1$	34.8		166		11.3	5.93	28
4D	2	$1s^2$	9.92		33.8		8.82	3.65	9.26
4D	3	$1s^2\,2s^1$	471		7 518		41.3	79.9	94.1
4D	4	$1s^2\,2s^2$	132		1 548		37.1	64.5	24
4D	12	$2s^2\,2p^8$	3.54		13.3		16.3	12.4	1.07
4D	13	$2p^8\,3s^1$	122		1 184		34.2	54.7	21.7
4D	14	$2p^8\,3s^2$	74.9		785		37.6	66.4	11.8
4D	22	$3s^2\,3p^8$	9.9		71.7		34.2	54.9	1.3
4D	23	$3p^8\,4s^1$	129		1 393		51.4	124	11.2
4D	24	$3p^8\,4s^2$	97.5		1 257		57.2	153	8.2
4D	42	$3d^{18}\,4s^2$	11.7		58.7		28.6	38.4	1.53
4D	50	$4s^2\,4p^8$	7.7		53.2		37	64.3	0.827
4D	51	$4p^8\,5s^1$	77.2		688		51.1	123	5.61
4D	52	$4p^8\,5s^2$	67.1		758		57.4	154	4.92
4D	78	$5s^2\,5p^8$	8.39		67.3		49.3	114	0.591

J.2 Joining and separating atoms. A molecule's binding energy is measured from its atoms, so the atoms the molecular codes reach when a bond is pulled apart have to be the atoms of this paper. The light atoms of both worlds (22 of them) were computed by every method the later papers use: numerical Hartree–Fock, the local exchange of the table, and the Gaussian code (`gauss.js`, unrestricted Hartree–Fock and MP2) in the basis the molecular work uses. For the closed shells that basis lies 0.00 to 6.60 mHa above the numerical Hartree–Fock limit. For the open shells the unrestricted energy lies −5.50 to 2.30 mHa from the restricted numerical one; below it mostly, because an unrestricted determinant can polarize its spins, which is not a better basis. Then pairs of like atoms (ours H_2, N_2, F_2, Ne_2; in four dimensions elements 1, 2, 7, 10, 11, 12) were stretched from 6 to 16 bohr, each started three ways (from bare nuclei and from the two atoms with parallel and with antiparallel spins) and the lowest kept. Every pair separates into neutral atoms (Mulliken charges below 10^{-5}) with each atom's own spin, but not from every start: the bare-nuclei start lands up to 16 hartree too high, on ion pairs and wrong spin states. A molecular code that starts one way only can report a binding energy measured from the wrong atoms. In ours the separated pairs meet the atoms to 0.001 mHa at 16 bohr. In four dimensions they don't meet them at all, as section 4 says they shouldn't. Two closed-shell atoms keep the permanent pull, and Hartree–Fock reproduces its far form: element 12's pair at 16 bohr is bound by −0.0119 against −0.0116 mHa, and the closed shells sit at 1.02–1.06 times the far form there. Hydrogen, whose four-dimensional orbital is diffuse, is at 1.15, still approaching it; its excess over the far form falls by five between 10 and 16 bohr, as shorter-ranged terms do. The open p shells are at 1.4–1.9: an unrestricted atom with a partly filled p shell is not round, and the ordinary quadrupole that ours has too adds a pull of the same order, one that depends on how the two atoms are turned. The atomic energies, the curves and each atom's charge and spin are in `data/separate.json`.

K · Reproducing every number

Every number in the text is read by `papers/p1/build.mjs` from one of the files below, and the selftest (block 67) fails if any claim in `ledger.json` stops holding or any generated file is stale. What made each file:

file	made by	where
`atom/js/table.js` (the bake; supplement S1)	`atom/tools/bake.mjs`, then `--repair` for 88, 101–103, 105–107	GitHub's runners, six minutes a shard
`data/termtable.json` (the main table)	`atom/tools/termtable.mjs`, collected by `termtable-collect.mjs`	runner run `hund-f-and-terms`
`data/hund.json`	`atom/tools/hund.mjs survey`, collected by `hund-collect.mjs`	sandbox (p, d, first f) and runner run `hund-f`
`data/hund-law.json`	`papers/p1/hundlaw.mjs`	a minute
`data/promotion3.json`	`termtable.mjs` with `TERM_MODE=full` on ours, `promotion-collect.mjs`	sandbox, minutes
`data/shelltable.json`	`atom/tools/lawtable.mjs shell`, `repair`, `bake`	runner runs `lawtable-shell`, `lawtable-repair`
`data/audit-scf.json`, `data/convergence.json`	`audit-scf.mjs`, `convergence.mjs`	runner
`data/profiles.json`	`profiles.mjs`	sandbox
`data/ions.json`	`atom/tools/termtable.mjs` with a charge, collected by `ions-collect.mjs`	runner runs `ions-1`, `ions-23`
`data/hftable.json`	`atom/tools/hftable.mjs rank`, collected by `hf-collect.mjs`	runner runs `hf-table-1`, `hf-lanth-1`, `hf-wide-1`, `hf-climb-1`
`data/valence.json`	`atom/tools/hftable.mjs map`, collected by `valence-collect.mjs`	sandbox
`data/valencetest.json`	`atom/tools/valencetest.mjs`, collected by `valencetest-collect.mjs`	runner run `valence-1`
`data/response.json`	`response-table.mjs`	sandbox, six minutes
`data/ligandd2.json`	`atom/tools/ligandd2.mjs`	sandbox, seconds
`data/separate.json`	`atom/tools/separate.mjs`, collected by `separate-collect.mjs`	sandbox
`data/nist-*.json`, `data/response-ref.json` (measured and reference values)	fetched from NIST (the levels by `nist-levels.py`); the reference by hand from its source	by hand
`data/table3.*`, `data/table4.*`, `figdata/`, `figures/`, `ledger.json`, `draft/`	`build.mjs`	seconds
`data/crossref.json` (the references' metadata)	`refs-crossref.mjs`, from each DOI's Crossref record	by hand, when a DOI is added
`draft/*.html`, `draft/paper.pdf`	`page.mjs` (pandoc, with citeproc), `pdf.cjs`	seconds

`papers/p1/reproduce.mjs` re-solves a sample of atoms from the solver and checks them against the paper's data; `--all` re-solves every element. The runner's logs, with each command's exit status, are kept on the repository's `atom4-data` branch under `runs/`. The solvers, the known-answer test assembled from the selftest's method blocks, `reproduce.mjs` and the paper's tables are mirrored under the MIT licence at tangled.org/morphyxmino.bsky.social/four-atom (`npm test` runs the known-answer test; `node papers/p1/reproduce.mjs` re-solves the sample), refreshed on every change to them.

Supplement and data

Supplement S1 is the table with every open subshell given Hund spins (`data/table4-hundspin.csv`). The 40 claims this text rests on, and the data behind them, are in `ledger.json` and `data/`.

References

Al-Jaber, Sami M. 1998. "Hydrogen Atom in N Dimensions." *International Journal of Theoretical Physics* 37 (4): 1289–98. https://doi.org/10.1023/A:1026679921970.

Alliluev, S. P. 1958. "On the Relation Between 'Accidental' Degeneracy and 'Hidden' Symmetry of a System." *Soviet Physics JETP* 6: 156–58. https://www.jetp.ras.ru/cgi-bin/dn/e_006_01_0156.pdf.

Bander, M., and C. Itzykson. 1966. "Group Theory and the Hydrogen Atom (I)." *Reviews of Modern Physics* 38 (2): 330–45. https://doi.org/10.1103/RevModPhys.38.330.

Barbatis, Gerassimos. 2024. "The Hardy Constant: A Review." In *Modern Problems in PDEs and Applications*, 3–11. Trends in Mathematics. Birkhäuser. https://doi.org/10.1007/978-3-031-56732-2_1.

Boys, S. F. 1950. "Electronic Wave Functions. I. A General Method of Calculation for the Stationary States of Any Molecular System." *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* 200 (1063): 542–54. https://doi.org/10.1098/rspa.1950.0036.

Callender, Craig. 2005. "Answers in Search of a Question: 'Proofs' of the Tri-Dimensionality of Space." *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics* 36 (1): 113–36. https://doi.org/10.1016/j.shpsb.2004.09.002.

Case, K. M. 1950. "Singular Potentials." *Physical Review* 80 (5): 797–806. https://doi.org/10.1103/PhysRev.80.797.

Derevianko, Andrei, Sergey G. Porsev, and James F. Babb. 2010. "Electric Dipole Polarizabilities at Imaginary Frequencies for Hydrogen, the Alkali-Metal, Alkaline-Earth, and Noble Gas Atoms." *Atomic Data and Nuclear Data Tables* 96 (3): 323–31. https://doi.org/10.1016/j.adt.2009.12.002.

Duc Pham, L V, Pascal Sattler, Miguel A L Marques, and Carlos L Benavides-Riveros. 2023. "Homogeneous Electron Liquid in Arbitrary Dimensions Beyond the Random Phase Approximation." *New Journal of Physics* 25 (8): 083040. https://doi.org/10.1088/1367-2630/acef4c.

Dyson, Freeman J., and A. Lenard. 1967. "Stability of Matter. I." *Journal of Mathematical Physics* 8 (3): 423–34. https://doi.org/10.1063/1.1705209.

Ehrenfest, P. 1918. “In What Way Does It Become Manifest in the Fundamental Laws of Physics That Space Has Three Dimensions?” *Proceedings of the Koninklijke Akademie van Wetenschappen Te Amsterdam* 20: 200–209. https://dwc.knaw.nl/DL/publications/PU00012213.pdf.

———. 1920. “Welche Rolle spielt die Dreidimensionalität des Raumes in den Grundgesetzen der Physik?” *Annalen der Physik* 366 (5): 440–46. https://doi.org/10.1002/andp.19203660503.

Escobar-Ruiz, A M, H Olivares-Pilón, N Aquino, and S A Cruz. 2021. “Helium-Like Ions in d-Dimensions: Analyticity and Generalized Ground State Majorana Solutions.” *Journal of Physics B: Atomic, Molecular and Optical Physics* 54 (23): 235002. https://doi.org/10.1088/1361-6455/ac3fbf.

FIORANI, EMANUELE. 2009. “Momentum Maps, Independent First Integrals and Integrability for Central Potentials.” *International Journal of Geometric Methods in Modern Physics* 06 (08): 1323–41. https://doi.org/10.1142/s0219887809004247.

Fock, V. 1935. “Zur Theorie des Wasserstoffatoms.” *Zeitschrift für Physik* 98 (3-4): 145–54. https://doi.org/10.1007/BF01336904.

Ghosh, Kumar J. B., Sabre Kais, and Dudley R. Herschbach. 2020. “Unorthodox Dimensional Interpolations for He, Li, Be Atoms and Hydrogen Molecule.” *Frontiers in Physics* 8 (August): 331. https://doi.org/10.3389/fphy.2020.00331.

Glasser, M. L., and J. Boersma. 1983. “Exchange Energy of an Electron Gas of Arbitrary Dimensionality.” *SIAM Journal on Applied Mathematics* 43 (3): 535–45. https://doi.org/10.1137/0143034.

González, Augusto. 1991. “The Quantum Few-Body Problem and the 1/D Method.” *Few-Body Systems* 10 (2): 43–57. https://doi.org/10.1007/bf01352401.

GU, XIAO-YAN, ZHONG-QI MA, and SHI-HAI DONG. 2002. “Exact Solutions to the Dirac Equation for a Coulomb Potential in D + 1 Dimensions.” *International Journal of Modern Physics E* 11 (04): 335–46. https://doi.org/10.1142/s0218301302000879.

Hainzl, Christian, and Robert Seiringer. 2002. “General Decomposition of Radial Functions on Rn and Applications to N-Body Quantum Systems.” *Letters in Mathematical Physics* 61 (1): 75–84. https://doi.org/10.1023/a:1020204818938.

Herschbach, Dudley R., John Avery, and Osvaldo Goscinski, eds. 1993. *Dimensional Scaling in Chemical Physics*. Dordrecht: Kluwer. https://doi.org/10.1007/978-94-011-1836-1.

Heydeman, Matthew, Christian B. Jepsen, Ziming Ji, and Amos Yarom. 2020. “Renormalization and Conformal Invariance of Non-Local Quantum Electrodynamics.” *Journal of High Energy Physics* 2020 (8): 7. https://doi.org/10.1007/jhep08(2020)007.

Katsura, Hosho, and Hideo Aoki. 2006. “Exact Supersymmetry in the Relativistic Hydrogen Atom in General Dimensions—Supercharge and the Generalized Johnson-Lippmann Operator.” *Journal of Mathematical Physics* 47 (3): 032301. https://doi.org/10.1063/1.2173173.

Kramida, A., Yu. Ralchenko, J. Reader, and NIST ASD Team. 2026. “NIST Atomic Spectra Database.” National Institute of Standards and Technology. 2026. https://physics.nist.gov/asd.

Kventsel, G. F., and J. Katriel. 1981. “Thomas–Fermi Atom in n Dimensions.” *Physical Review A* 24 (5): 2299–2301. https://doi.org/10.1103/physreva.24.2299.

La Nave, Gabriele, Kridsanaphong Limtragool, and Philip W. Phillips. 2019. “Colloquium : Fractional Electromagnetism in Quantum Matter and High-Energy Physics.” *Reviews of Modern Physics* 91 (2): 021003. https://doi.org/10.1103/revmodphys.91.021003.

Landau, L. D., and E. M. Lifshitz. 1977. *Quantum Mechanics: Non-Relativistic Theory*. 3rd ed. Course of Theoretical Physics 3. Oxford: Pergamon. https://doi.org/10.1016/C2013-0-02793-4.

Lewin, Mathieu. 2022. “Coulomb and Riesz Gases: The Known and the Unknown.” *Journal of Mathematical Physics* 63 (6): 061101. https://doi.org/10.1063/5.0086835.

Lieb, Elliott H., and Walter E. Thirring. 1975. “Bound for the Kinetic Energy of Fermions Which Proves the Stability of Matter.” *Physical Review Letters* 35 (11): 687–89. https://doi.org/10.1103/PhysRevLett.35.687.

Loeser, J. G. 1987. “Atomic Energies from the Large-Dimension Limit.” *The Journal of Chemical Physics* 86 (10): 5635–46. https://doi.org/10.1063/1.452540.

Loeser, J. G., and D. R. Herschbach. 1986a. “Hartree–Fock Approximation for D-Dimensional Two-Electron Atoms.” *The Journal of Chemical Physics* 84 (7): 3893–3900. https://doi.org/10.1063/1.450100.

———. 1986b. “Hylleraas–Pekeris Treatment of D-Dimensional Two-Electron Atoms.” *The Journal of Chemical Physics* 84 (7): 3882–92. https://doi.org/10.1063/1.450099.

Loeser, J. G., J. H. Summerfield, A. L. Tan, and Z. Zheng. 1994. “Correlated Electronic Structure Models Suggested by the Large-Dimension Limit.” *The Journal of Chemical Physics* 100 (7): 5036–53. https://doi.org/10.1063/1.467221.

Loos, Pierre-François, and Peter M. W. Gill. 2016. “The Uniform Electron Gas.” *WIREs Computational Molecular Science* 6 (4): 410–29. https://doi.org/10.1002/wcms.1257.

Madelung, Erwin. 1936. *Die mathematischen Hilfsmittel des Physikers*. 3rd ed. Grundlehren der mathematischen Wissenschaften 4. Berlin: Springer. https://doi.org/10.1007/978-3-662-02177-4.

Marino, E. C., Leandro O. Nascimento, Van Sérgio Alves, and C. Morais Smith. 2014. “Unitarity of Theories Containing Fractional Powers of the d’Alembertian Operator.” *Physical Review D* 90 (10): 105003. https://doi.org/10.1103/physrevd.90.105003.

McMurchie, Larry E, and Ernest R Davidson. 1978. “One- and Two-Electron Integrals over Cartesian Gaussian Functions.” *Journal of Computational Physics* 26 (2): 218–31. https://doi.org/10.1016/0021-9991(78)90092-X.

Négadi, T., and M. Kibler. 1996. “The Periodic Table in Flatland.” *International Journal of Quantum Chemistry* 57 (1): 53–61. https://doi.org/10.1002/(SICI)1097-461X(1996)57:1<53::AID-QUA6>3.0.CO;2-Y.

Nieto, Michael Martin. 1979. “Hydrogen Atom and Relativistic Pi-Mesic Atom in N-Space Dimensions.” *American Journal of Physics* 47 (12): 1067–72. https://doi.org/10.1119/1.11976.

Perdew, J. P., and Alex Zunger. 1981. “Self-Interaction Correction to Density-Functional Approximations for Many-Electron Systems.” *Physical Review B* 23 (10): 5048–79. https://doi.org/10.1103/PhysRevB.23.5048.

Peter Lindqvist. 2017. *Notes on the p-Laplace Equation, Second Edition*. 2nd ed. Vol. 161. Report, Department of Mathematics and Statistics. University of Jyväskylä. https://jyx.jyu.fi/bitstreams/2a8cbc94-58e9-4e75-94d6-0f1f5bc22ac1/download.

Pyykkö, Pekka, and Yongfang Zhao. 1991. “The Elements of Flatland: Hartree–Fock Atomic Ground States in Two Dimensions for Z = 1–24.” *International Journal of Quantum Chemistry* 40 (4): 527–44. https://doi.org/10.1002/qua.560400408.

Rienstra-Kiracofe, Jonathan C., Gregory S. Tschumper, Henry F. Schaefer, Sreela Nandi, and G. Barney Ellison. 2002. “Atomic and Molecular Electron Affinities: Photoelectron Experiments and Theoretical Computations.” *Chemical Reviews* 102 (1): 231–82. https://doi.org/10.1021/cr990044u.

Sako, Tokuei, Josef Paldus, and Geerd H. F. Diercksen. 2010. “Origin of Hund’s Multiplicity Rule in Quasi-Two-Dimensional Two-Electron Quantum Dots.” *Physical Review A* 81 (2): 022501. https://doi.org/10.1103/physreva.81.022501.

Sako, Tokuei, Josef Paldus, Atsushi Ichimura, and Geerd H F Diercksen. 2012. “Origin of the First Hund Rule and the Structure of Fermi Holes in Two-Dimensional He-Like Atoms and Two-Electron Quantum Dots.” *Journal of Physics B: Atomic, Molecular and Optical Physics* 45 (23): 235001. https://doi.org/10.1088/0953-4075/45/23/235001.

Sansonetti, J. E., and W. C. Martin. 2005. “Handbook of Basic Atomic Spectroscopic Data.” *Journal of Physical and Chemical Reference Data* 34 (4): 1559–2259. https://doi.org/10.1063/1.1800011.

Santos, F. C., V. Soares, and A. C. Tort. 2007. “An English Translation of Bertrand’s Theorem.” arXiv. 2007. https://arxiv.org/abs/0704.2396.

Schlesier, Robert, Carlos L. Benavides-Riveros, and Miguel A. L. Marques. 2020. “Homogeneous Electron Gas in Arbitrary Dimensions.” *Physical Review B* 102 (3): 035123. https://doi.org/10.1103/physrevb.102.035123.

Summerfield, J.H., and J.G. Loeser. 1999. “Dimension-dependent Two-electron Hamiltonian Matrix Elements.” *Journal of Mathematical Chemistry* 25 (2-3): 309–15. https://doi.org/10.1023/a:1019196719751.

Sun, Qiming, Timothy C. Berkelbach, Nick S. Blunt, George H. Booth, Sheng Guo, Zhendong Li, Junzi Liu, et al. 2018. “PySCF: The Python-Based Simulations of Chemistry Framework.” *WIREs Computational Molecular Science* 8 (1): e1340. https://doi.org/10.1002/wcms.1340.

Sun, Qiming, Xing Zhang, Samragni Banerjee, Peng Bao, Marc Barbry, Nick S. Blunt, Nikolay A. Bogdanov, et al. 2020. “Recent Developments in the PySCF Program Package.” *The Journal of Chemical Physics* 153 (2): 024109. https://doi.org/10.1063/5.0006074.

Tanabe, Yukito, and Satoru Sugano. 1954. “On the Absorption Spectra of Complex Ions I.” *Journal of the Physical Society of Japan* 9 (5): 753–66. https://doi.org/10.1143/JPSJ.9.753.

Tangherlini, F. R. 1963. “Schwarzschild Field in n Dimensions and the Dimensionality of Space Problem.” *Il Nuovo Cimento* 27 (3): 636–51. https://doi.org/10.1007/BF02784569.

Tarasov, Vasily E. 2016. “Electric Field in Media with Power-Law Spatial Dispersion.” *Modern Physics Letters B* 30 (10): 1650132. https://doi.org/10.1142/s0217984916501323.

Tarasov, Vasily E., and Juan J. Trujillo. 2013. “Fractional Power-Law Spatial Dispersion in Electrodynamics.” *Annals of Physics* 334 (July): 1–23. https://doi.org/10.1016/j.aop.2013.03.014.