

The Answer of Assignment 2

WEI SHUANG

31/8/2025

Problem 1 Solution

- (1) 根据线性空间的同构理论，任何维数相同的线性空间都是同构的。因此可知 $\sum_{i,j=0}^1 c_{ij} |i,j\rangle$ 与 $\{(x_1, x_2, x_3, x_4)\}, x \in C$ 同构。即 $v_1 = (c_{00}, c_{01}, c_{02}, c_{03})^*, v_2 = (d_{00}, d_{01}, d_{10}, d_{11})^*$
- (2) $\langle \psi_1 | \psi_2 \rangle = v_1^\dagger v_2$ 成立，这是因为 $\langle \psi_1 | \psi_2 \rangle = (\sum_{i,j=0}^1 c_{i,j}^* \langle ij|) (\sum_{i,j=0}^1 c_{i,j} |ij\rangle) =$
- (3) $v_1^\dagger v_2$
- (4)

Problem 2 Solution

- (1)
- (2)
- (3)
- (4)

Problem 3 Solution

- (1)
- (2)
- (3)

(4) how should the Pauli operators σ_i^+ and σ_i^z can be written in terms of the f operators? we have the definition that:

$$f_i^\dagger = \left(\prod_{j<i} \sigma_j^z \right) \sigma_i^+$$

so from $\langle \overline{n_1 n_2 \dots n_i} | f_i^\dagger | \overline{n_1 n_2 n_3 \dots n_i} \rangle = \langle \overline{n_1 n_2 \dots n_i} | f_i | \overline{n_1 n_2 \dots n_i} \rangle^*$ we can get the relation between f_i and σ_i^+, σ_i^z as follows:

$$f_i = \left(\prod_{j<i} \sigma_j^z \right) \sigma_i^-$$

then try to represent σ_i^z and σ_i^+ in terms of f_i : we use the matrix representation of the Pauli operators:

$$\sigma_i^z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \sigma_i^+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \sigma_i^- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

σ_i^z can be expressed as: $2\sigma_i^+ \sigma_i^- - I = 2f_i^\dagger f_i - I$

like wise:

$$\begin{aligned} \sigma_i^+ &= f_i^\dagger \left(\prod_{j<i} \sigma_j^z \right)^{-1} \\ &= f_i^\dagger \left(\prod_{j<i} (2f_j^\dagger f_j - I) \right) \end{aligned}$$