

# Economics 103 – Statistics for Economists

Minsu Chang

University of Pennsylvania

Lecture # 3

## Recall From Last Lecture:

### Range

Maximum Observation - Minimum Observation

### Interquartile Range (IQR)

$$\text{IQR} = Q_3 - Q_1$$

### Variance

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$$

### Standard Deviation

$$s = \sqrt{s^2}$$

## Recall From Last Lecture:

### Variance

Essentially the average squared distance from the mean. Sensitive to both skewness and outliers.

### Standard Deviation

$\sqrt{\text{Variance}}$ , but more convenient since **same units as data**

### Range

Difference between largest and smallest observations. *Very* sensitive to outliers. Displayed in boxplot.

### Interquartile Range

Range of middle 50% of the data. Insensitive to outliers, skewness. Displayed in boxplot.

## Why Squares?

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$$

What's Wrong With This?

$$\frac{1}{n-1} \sum_{i=1}^N (x_i - \bar{x}) =$$

## Why Squares?

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$$

What's Wrong With This?

$$\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x}) \quad = \quad \frac{1}{n-1} \left[ \sum_{i=1}^n x_i - \sum_{i=1}^n \bar{x} \right] =$$

## Why Squares?

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$$

What's Wrong With This?

$$\begin{aligned} \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x}) &= \frac{1}{n-1} \left[ \sum_{i=1}^n x_i - \sum_{i=1}^n \bar{x} \right] = \frac{1}{n-1} \left[ \sum_{i=1}^n x_i - n\bar{x} \right] \\ &= \end{aligned}$$

## Why Squares?

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$$

What's Wrong With This?

$$\begin{aligned} \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x}) &= \frac{1}{n-1} \left[ \sum_{i=1}^n x_i - \sum_{i=1}^n \bar{x} \right] = \frac{1}{n-1} \left[ \sum_{i=1}^n x_i - n\bar{x} \right] \\ &= \frac{1}{n-1} \left[ \sum_{i=1}^n x_i - n \cdot \frac{1}{n} \sum_{i=1}^n x_i \right] \\ &= \end{aligned}$$

## Why Squares?

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$$

What's Wrong With This?

$$\begin{aligned} \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x}) &= \frac{1}{n-1} \left[ \sum_{i=1}^n x_i - \sum_{i=1}^n \bar{x} \right] = \frac{1}{n-1} \left[ \sum_{i=1}^n x_i - n\bar{x} \right] \\ &= \frac{1}{n-1} \left[ \sum_{i=1}^n x_i - n \cdot \frac{1}{n} \sum_{i=1}^n x_i \right] \\ &= \frac{1}{n-1} \left[ \sum_{i=1}^n x_i - \sum_{i=1}^n x_i \right] = \end{aligned}$$



## Why Squares?

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$$

What's Wrong With This?

$$\begin{aligned} \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x}) &= \frac{1}{n-1} \left[ \sum_{i=1}^n x_i - \sum_{i=1}^n \bar{x} \right] = \frac{1}{n-1} \left[ \sum_{i=1}^n x_i - n\bar{x} \right] \\ &= \frac{1}{n-1} \left[ \sum_{i=1}^n x_i - n \cdot \frac{1}{n} \sum_{i=1}^n x_i \right] \\ &= \frac{1}{n-1} \left[ \sum_{i=1}^n x_i - \sum_{i=1}^n x_i \right] = 0 \end{aligned}$$

# Variance is Sensitive to Skewness and Outliers

And so is Standard Deviation!

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$$

## Outliers

Differentiate with respect to  $(x_i - \bar{x}) \Rightarrow$  the farther an observation is from the mean, the *larger* its effect on the variance.

## Skewness

Variance measures average squared distance from center, taking **mean** as the center, but the mean is sensitive to skewness!

## Skewness – A Measure of Symmetry

$$\text{Skewness} = \frac{1}{n} \frac{\sum_{i=1}^n (x_i - \bar{x})^3}{s^3}$$

## Skewness – A Measure of Symmetry

$$\text{Skewness} = \frac{1}{n} \frac{\sum_{i=1}^n (x_i - \bar{x})^3}{s^3}$$

What do the values indicate?

Zero  $\Rightarrow$  symmetry, positive right-skewed, negative left-skewed.

## Skewness – A Measure of Symmetry

$$\text{Skewness} = \frac{1}{n} \frac{\sum_{i=1}^n (x_i - \bar{x})^3}{s^3}$$

What do the values indicate?

Zero  $\Rightarrow$  symmetry, positive right-skewed, negative left-skewed.

Why cubed?

To get the desired sign.

## Skewness – A Measure of Symmetry

$$\text{Skewness} = \frac{1}{n} \frac{\sum_{i=1}^n (x_i - \bar{x})^3}{s^3}$$

What do the values indicate?

Zero  $\Rightarrow$  symmetry, positive right-skewed, negative left-skewed.

Why cubed?

To get the desired sign.

Why divide by  $s^3$ ?

So that skewness is unitless

## Skewness – A Measure of Symmetry

$$\text{Skewness} = \frac{1}{n} \frac{\sum_{i=1}^n (x_i - \bar{x})^3}{s^3}$$

What do the values indicate?

Zero  $\Rightarrow$  symmetry, positive right-skewed, negative left-skewed.

Why cubed?

To get the desired sign.

Why divide by  $s^3$ ?


So that skewness is unitless

Rule of Thumb

Typically (but not always), right-skewed  $\Rightarrow$  mean  $>$  median


left-skewed  $\Rightarrow$  mean  $<$  median

# Histogram of Handspan





# Histogram of Handedness



## Essential Distinction: Sample vs. Population

For now, you can think of the population as a list of  $N$  objects:

Population:  $x_1, x_2, \dots, x_N$

from which we draw a sample of size  $n < N$  objects:

Sample:  $x_1, x_2, \dots, x_n$

# Essential Distinction: Sample vs. Population

For now, you can think of the population as a list of  $N$  objects:

Population:  $x_1, x_2, \dots, x_N$

from which we draw a sample of size  $n < N$  objects:

Sample:  $x_1, x_2, \dots, x_n$

## Important Point:

Later in the course we'll be more formal by considering **probability models** that represent the *act of sampling* from a population rather than thinking of a population as a list of objects. Once we do this we will no longer use the notation  $N$  as the population will be *conceptually infinite*.

# Essential Distinction: Parameter vs. Statistic

$N$  individuals in the Population,  $n$  individuals in the Sample:

	Parameter (Population)	Statistic (Sample)
Mean	$\mu = \frac{1}{N} \sum_{i=1}^N x_i$	$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
Var.	$\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$	$s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$
S.D.	$\sigma = \sqrt{\sigma^2}$	$s = \sqrt{s^2}$

## Key Point

We use a **sample**  $x_1, \dots, x_n$  to calculate **statistics** (e.g.  $\bar{x}$ ,  $s^2$ ,  $s$ ) that serve as **estimates** of the corresponding population **parameters** (e.g.  $\mu$ ,  $\sigma^2$ ,  $\sigma$ ).

## Why Do Sample Variance and Std. Dev. Divide by $n - 1$ ?

Pop. Var. $\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$	Sample Var. $s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$
Pop. S.D. $\sigma = \sqrt{\sigma^2}$	Sample S.D. $s = \sqrt{s^2}$

There is an important reason for this, but explaining it requires some concepts we haven't learned yet.

# Why Mean and Variance (and Std. Dev. )?

## Empirical Rule

For large populations that are approximately bell-shaped, std. dev. tells where most observations will be relative to the mean:

- ▶  $\approx 68\%$  of observations are in the interval  $\mu \pm \sigma$
- ▶  $\approx 95\%$  of observations are in the interval  $\mu \pm 2\sigma$
- ▶ Almost all of observations are in the interval  $\mu \pm 3\sigma$

# Why Mean and Variance (and Std. Dev. )?

## Empirical Rule

For large populations that are approximately bell-shaped, std. dev. tells where most observations will be relative to the mean:

- ▶  $\approx 68\%$  of observations are in the interval  $\mu \pm \sigma$
- ▶  $\approx 95\%$  of observations are in the interval  $\mu \pm 2\sigma$
- ▶ Almost all of observations are in the interval  $\mu \pm 3\sigma$

## Therefore

We will be interested in  $\bar{x}$  as an estimate of  $\mu$  and  $s$  as an estimate of  $\sigma$  since these population parameters are so informative.



Which is more “extreme?”

(a) Handspan of 27cm

(b) Height of 78in



## Centering: Subtract the Mean

Handspan	Height
$27\text{cm} - 20.6\text{cm} = 6.4\text{cm}$	$78\text{in} - 67.6\text{in} = 10.4\text{in}$

## Standardizing: Divide by S.D.

Handspan	Height
$27\text{cm} - 20.6\text{cm} = 6.4\text{cm}$	$78\text{in} - 67.6\text{in} = 10.4\text{in}$
$6.4\text{cm}/2.2\text{cm} \approx 2.9$	$10.4\text{in}/4.5\text{in} \approx 2.3$

## Standardizing: Divide by S.D.

Handspan	Height
$27\text{cm} - 20.6\text{cm} = 6.4\text{cm}$	$78\text{in} - 67.6\text{in} = 10.4\text{in}$
$6.4\text{cm}/2.2\text{cm} \approx 2.9$	$10.4\text{in}/4.5\text{in} \approx 2.3$

The units have disappeared!

# Z-scores: How many standard deviations from the mean?

Best for Symmetric Distribution, No Outliers (Why?)

$$z_i = \frac{x_i - \bar{x}}{s}$$

# Z-scores: How many standard deviations from the mean?

Best for Symmetric Distribution, No Outliers (Why?)

$$z_i = \frac{x_i - \bar{x}}{s}$$

Unitless

Allows comparison of variables with different units.

# Z-scores: How many standard deviations from the mean?

Best for Symmetric Distribution, No Outliers (Why?)

$$z_i = \frac{x_i - \bar{x}}{s}$$

## Unitless

Allows comparison of variables with different units.

## Detecting Outliers

Measures how “extreme” one observation is relative to the others.

# Z-scores: How many standard deviations from the mean?

Best for Symmetric Distribution, No Outliers (Why?)

$$z_i = \frac{x_i - \bar{x}}{s}$$

## Unitless

Allows comparison of variables with different units.

## Detecting Outliers

Measures how “extreme” one observation is relative to the others.

## Linear Transformation

What is the sample mean of the z-scores?

$$\bar{z} = \frac{1}{n} \sum_{i=1}^n z_i$$



What is the sample mean of the z-scores?

$$\bar{z} = \frac{1}{n} \sum_{i=1}^n z_i = \frac{1}{n} \sum_{i=1}^n \frac{x_i - \bar{x}}{s} =$$

What is the sample mean of the z-scores?

$$\begin{aligned}\bar{z} &= \frac{1}{n} \sum_{i=1}^n z_i = \frac{1}{n} \sum_{i=1}^n \frac{x_i - \bar{x}}{s} = \frac{1}{n \cdot s} \left[ \sum_{i=1}^n x_i - \sum_{i=1}^n \bar{x} \right] \\ &= \end{aligned}$$

What is the sample mean of the z-scores?

$$\begin{aligned}\bar{z} &= \frac{1}{n} \sum_{i=1}^n z_i = \frac{1}{n} \sum_{i=1}^n \frac{x_i - \bar{x}}{s} = \frac{1}{n \cdot s} \left[ \sum_{i=1}^n x_i - \sum_{i=1}^n \bar{x} \right] \\ &= \frac{1}{n \cdot s} \left[ \sum_{i=1}^n x_i - n\bar{x} \right] =\end{aligned}$$

What is the sample mean of the z-scores?

$$\begin{aligned}\bar{z} &= \frac{1}{n} \sum_{i=1}^n z_i = \frac{1}{n} \sum_{i=1}^n \frac{x_i - \bar{x}}{s} = \frac{1}{n \cdot s} \left[ \sum_{i=1}^n x_i - \sum_{i=1}^n \bar{x} \right] \\ &= \frac{1}{n \cdot s} \left[ \sum_{i=1}^n x_i - n\bar{x} \right] = \frac{1}{n \cdot s} \left[ \sum_{i=1}^n x_i - n \cdot \frac{1}{n} \sum_{i=1}^n x_i \right] \\ &= \end{aligned}$$

What is the sample mean of the z-scores?

$$\begin{aligned}\bar{z} &= \frac{1}{n} \sum_{i=1}^n z_i = \frac{1}{n} \sum_{i=1}^n \frac{x_i - \bar{x}}{s} = \frac{1}{n \cdot s} \left[ \sum_{i=1}^n x_i - \sum_{i=1}^n \bar{x} \right] \\&= \frac{1}{n \cdot s} \left[ \sum_{i=1}^n x_i - n\bar{x} \right] = \frac{1}{n \cdot s} \left[ \sum_{i=1}^n x_i - n \cdot \frac{1}{n} \sum_{i=1}^n x_i \right] \\&= \frac{1}{n \cdot s} \left[ \sum_{i=1}^n x_i - \sum_{i=1}^n x_i \right] =\end{aligned}$$

What is the sample mean of the z-scores?

$$\begin{aligned}\bar{z} &= \frac{1}{n} \sum_{i=1}^n z_i = \frac{1}{n} \sum_{i=1}^n \frac{x_i - \bar{x}}{s} = \frac{1}{n \cdot s} \left[ \sum_{i=1}^n x_i - \sum_{i=1}^n \bar{x} \right] \\&= \frac{1}{n \cdot s} \left[ \sum_{i=1}^n x_i - n\bar{x} \right] = \frac{1}{n \cdot s} \left[ \sum_{i=1}^n x_i - n \cdot \frac{1}{n} \sum_{i=1}^n x_i \right] \\&= \frac{1}{n \cdot s} \left[ \sum_{i=1}^n x_i - \sum_{i=1}^n x_i \right] = 0\end{aligned}$$

What is the variance of the z-scores?

$$s_z^2 = \frac{1}{n-1} \sum_{i=1}^n (z_i - \bar{z})^2 =$$

What is the variance of the z-scores?

$$s_z^2 = \frac{1}{n-1} \sum_{i=1}^n (z_i - \bar{z})^2 = \frac{1}{n-1} \sum_{i=1}^n z_i^2 =$$



What is the variance of the z-scores?

$$s_z^2 = \frac{1}{n-1} \sum_{i=1}^n (z_i - \bar{z})^2 = \frac{1}{n-1} \sum_{i=1}^n z_i^2 = \frac{1}{n-1} \sum_{i=1}^n \left( \frac{x_i - \bar{x}}{s_x} \right)^2$$

=

What is the variance of the z-scores?

$$\begin{aligned}s_z^2 &= \frac{1}{n-1} \sum_{i=1}^n (z_i - \bar{z})^2 = \frac{1}{n-1} \sum_{i=1}^n z_i^2 = \frac{1}{n-1} \sum_{i=1}^n \left( \frac{x_i - \bar{x}}{s_x} \right)^2 \\&= \frac{1}{s_x^2} \left[ \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2 \right] =\end{aligned}$$

What is the variance of the z-scores?

$$\begin{aligned}s_z^2 &= \frac{1}{n-1} \sum_{i=1}^n (z_i - \bar{z})^2 = \frac{1}{n-1} \sum_{i=1}^n z_i^2 = \frac{1}{n-1} \sum_{i=1}^n \left( \frac{x_i - \bar{x}}{s_x} \right)^2 \\&= \frac{1}{s_x^2} \left[ \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2 \right] = \frac{s_x^2}{s_x^2} =\end{aligned}$$

What is the variance of the z-scores?

$$\begin{aligned}s_z^2 &= \frac{1}{n-1} \sum_{i=1}^n (z_i - \bar{z})^2 = \frac{1}{n-1} \sum_{i=1}^n z_i^2 = \frac{1}{n-1} \sum_{i=1}^n \left( \frac{x_i - \bar{x}}{s_x} \right)^2 \\&= \frac{1}{s_x^2} \left[ \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2 \right] = \frac{s_x^2}{s_x^2} = 1\end{aligned}$$

What is the variance of the z-scores?

$$\begin{aligned}s_z^2 &= \frac{1}{n-1} \sum_{i=1}^n (z_i - \bar{z})^2 = \frac{1}{n-1} \sum_{i=1}^n z_i^2 = \frac{1}{n-1} \sum_{i=1}^n \left( \frac{x_i - \bar{x}}{s_x} \right)^2 \\&= \frac{1}{s_x^2} \left[ \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2 \right] = \frac{s_x^2}{s_x^2} = 1\end{aligned}$$

So what is the *standard deviation* of the z-scores?



## Population Z-scores and the Empirical Rule: $\mu \pm 2\sigma$

If we knew the population mean  $\mu$  and standard deviation  $\sigma$  we could create a *population version* of a z-score. This leads to an important way of rewriting the Empirical Rule:

## Population Z-scores and the Empirical Rule: $\mu \pm 2\sigma$

If we knew the population mean  $\mu$  and standard deviation  $\sigma$  we could create a *population version* of a z-score. This leads to an important way of rewriting the Empirical Rule:

Bell-shaped population  $\Rightarrow$  approx. 95% of observations  $x_i$  satisfy

$$\mu - 2\sigma \leq x_i \leq \mu + 2\sigma$$

## Population Z-scores and the Empirical Rule: $\mu \pm 2\sigma$

If we knew the population mean  $\mu$  and standard deviation  $\sigma$  we could create a *population version* of a z-score. This leads to an important way of rewriting the Empirical Rule:

Bell-shaped population  $\Rightarrow$  approx. 95% of observations  $x_i$  satisfy

$$\mu - 2\sigma \leq x_i \leq \mu + 2\sigma$$

$$-2\sigma \leq x_i - \mu \leq 2\sigma$$



## Population Z-scores and the Empirical Rule: $\mu \pm 2\sigma$

If we knew the population mean  $\mu$  and standard deviation  $\sigma$  we could create a *population version* of a z-score. This leads to an important way of rewriting the Empirical Rule:

Bell-shaped population  $\Rightarrow$  approx. 95% of observations  $x_i$  satisfy

$$\mu - 2\sigma \leq x_i \leq \mu + 2\sigma$$

$$-2\sigma \leq x_i - \mu \leq 2\sigma$$

$$-2 \leq \frac{x_i - \mu}{\sigma} \leq 2$$

# Relationships Between Variables

## Crosstabs – Show Relationship between Categorical Vars.

(aka Contingency Tables)

<i>Eye Color</i>	<i>Sex</i>		Total
	Male	Female	
Black	5	2	7
Blue	6	4	10
Brown	26	31	57
Copper	1	0	1
Dark Brown	0	1	1
Green	4	1	5
Hazel	2	2	4
Maroon	1	0	1
Total	45	41	86

## Example with Crosstab in *Percents*

# Who Supported the Vietnam War?

In January 1971 the Gallup poll asked: “A proposal has been made in Congress to require the U.S. government to bring home all U.S. troops before the end of this year. Would you like to have your congressman vote for or against this proposal?”

Guess the results, for respondents in each education category, and fill out this table (the two numbers in each column should add up to 100%):

	Adults with:			Total adults
	Grade school education	High school education	College education	
% for withdrawal of U.S. troops (doves)				73%
% against withdrawal of U.S. troops (hawks)				27%
Total	100%	100%	100%	100%

## Who Were the Doves?

Which group do you think was most strongly in favor of the withdrawal of US troops from Vietnam?

- (a) Adults with only a Grade School Education
- (b) Adults with a High School Education
- (c) Adults with a College Education

## Who Were the Hawks?

Which group do you think was most strongly **opposed to** the withdrawal of US troops from Vietnam?

- (a) Adults with only a Grade School Education
- (b) Adults with a High School Education
- (c) Adults with a College Education

## From The Economist – “Lexington,” October 4th, 2001

*“Back in the Vietnam days, the anti-war movement spread from the intelligentsia into the rest of the population, eventually paralyzing the country’s will to fight.”*



# Who *Really* Supported the Vietnam War

Gallup Poll, January 1971

	Adults with:			
	Grade school education	High school education	College education	Total adults
% for withdrawal of U.S. troops (doves)	80%	75%	60%	73%
% against withdrawal of U.S. troops (hawks)	20%	25%	40%	27%
Total	100%	100%	100%	100%

What about numeric data?

# Covariance and Correlation: Linear Dependence Measures

## Two Samples of Numeric Data

$x_1, \dots, x_n$  and  $y_1, \dots, y_n$

## Dependence

Do  $x$  and  $y$  both tend to be large (or small) at the same time?

## Key Point

Use the idea of centering and standardizing to decide what “big” or “small” means in this context.

# Notation

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$

$$\bar{y} = \frac{1}{n} \sum_{i=1}^n y_i$$

$$s_x = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2}$$

$$s_y = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (y_i - \bar{y})^2}$$

# Covariance

$$s_{xy} = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})$$

- ▶ Centers each observation around its mean and multiplies.
- ▶ Zero  $\Rightarrow$  no linear dependence
- ▶ Positive  $\Rightarrow$  positive linear dependence
- ▶ Negative  $\Rightarrow$  negative linear dependence
- ▶ Population parameter:  $\sigma_{xy}$
- ▶ Units?

# Correlation

$$r_{xy} = \frac{1}{n-1} \sum_{i=1}^n \left( \frac{x_i - \bar{x}}{s_x} \right) \left( \frac{y_i - \bar{y}}{s_y} \right) = \frac{s_{xy}}{s_x s_y}$$

- ▶ Centers *and* standardizes each observation
- ▶ Bounded between -1 and 1
- ▶ Zero  $\Rightarrow$  no linear dependence
- ▶ Positive  $\Rightarrow$  positive linear dependence
- ▶ Negative  $\Rightarrow$  negative linear dependence
- ▶ Population parameter:  $\rho_{xy}$
- ▶ Unitless

We'll have more to say about correlation and covariance when we discuss linear regression.

# Essential Distinction: Parameter vs. Statistic

## And Population vs. Sample

$N$  individuals in the Population,  $n$  individuals in the Sample:

	Parameter (Population)	Statistic (Sample)
Mean	$\mu_x = \frac{1}{N} \sum_{i=1}^N x_i$	$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
Var.	$\sigma_x^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$	$s_x^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$
S.D.	$\sigma_x = \sqrt{\sigma_x^2}$	$s_x = \sqrt{s^2}$
Cov.	$\sigma_{xy} = \frac{\sum_{i=1}^N (x_i - \mu_x)(y_i - \mu_y)}{N}$	$s_{xy} = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{n-1}$
Corr.	$\rho = \frac{\sigma_{xy}}{\sigma_x \sigma_y}$	$r = \frac{s_{xy}}{s_x s_y}$