

LIMIT THEOREMS FOR HIGHER RANK ACTIONS ON HEISENBERG NILMANIFOLDS

MINSUNG KIM*

ABSTRACT. The main result of this paper is a construction of finitely additive measures for higher rank abelian actions on Heisenberg nilmanifolds. Under a full measure set of Diophantine condition for the generators of the action, we construct *Bufetov functional* on rectangles on $2g + 1$ dimensional Heisenberg manifold. We prove that the deviation of ergodic integral of higher rank actions is described by the asymptotics of Bufetov functionals for a sufficiently smooth function. As a corollary, the distribution of normalized ergodic integrals to have variance 1, converges along certain subsequences to a non-degenerate compactly supported measure on the real line.

CONTENTS

1.	Introduction	2
1.1.	Introduction	2
1.2.	Definitions and statement of results	3
	Main results	5
2.	Analysis on Heisenberg manifolds	8
2.1.	Sobolev space	8
2.2.	Invariant currents	9
2.3.	Diophantine condition	10
3.	Constructions of the functionals	12
3.1.	Remainder estimates	12
3.2.	Constructions of the functionals	16
3.3.	Proof of Theorem 1.3	18
3.4.	Proof of Theorem 1.6	20
4.	Limit distributions	21
4.1.	Proof of Theorem 1.7	21
5.	L^2 -lower bounds	24
5.1.	Structure of return map	24
5.2.	Higher cohomology for \mathbb{Z}^d -action of skew-shifts	26
5.3.	Changes of coordinates	27
5.4.	L^2 -lower bound of functional	28
6.	Analyticity of functionals	31
6.1.	Analyticity	31
6.2.	Extensions of domain	34
7.	Measure estimation for bounded-type	35

Date: December 10, 2021.

2010 Mathematics Subject Classification. 37A17, 37A20, 37A44, 60F05.

Key words and phrases. Higher rank abelian actions, Limit theorems, Heisenberg groups.

1	Acknowledgement	37
2	References	37

3 1. INTRODUCTION

4 **1.1. Introduction.** The asymptotic behavior and limiting distribution of ergodic
5 averages of translation flow were studied by Alexander Bufetov in the series of his
6 works [Buf14a, Buf10, Buf14b]. He constructed finitely-additive Hölder measures
7 and cocycles over translation flows that are known as *Bufetov functionals*. Such
8 functionals are constructed to derive the deviation of ergodic integrals of translation
9 flows by providing a new proof of the celebrated work of G. Forni [For02]. Bufetov
10 noticed a duality between his finitely-additive Hölder measures and Forni's invariant
11 distributions, which played a crucial role in his work of cohomological equations.
12 Following these observations, the duality between such functionals and the space
13 of invariant distributions was constructed analogously for other parabolic flows
14 and higher rank actions (e.g. [Buf13, BS13, BF14, FK20b, For20a]). Bufetov also
15 proved the probabilistic results for normalized ergodic integrals as a part of his
16 main theorem. Furthermore, it turned out that his construction of functional is also
17 closely related to limit shapes of ergodic sums for interval exchange transformation
18 [MMY10, MUY20].

19 In this paper, our main results are on effective equidistributions for higher rank
20 abelian actions and limit theorems on Heisenberg nilmanifolds. We firstly introduce
21 the construction of the Bufetov functional for higher rank abelian actions. The
22 main argument is based on the renormalizations and induction argument from the
23 work of Cosentino and Flaminio [CF15]. However, in our case, we extended their
24 constructions to the rectangular domain and allowed general products of intervals.
25 We assume that the functional exists for a full measure set of frame α under a
26 sufficient Diophantine condition obtained from recurrent conditions (see §2.3.2).
27 Likewise, we also constructed the duality between functionals and invariant currents
28 in [CF15], enabling the deviation formula. As a corollary, we prove existence of
29 the limit distributions of (normalized) ergodic integrals for higher rank actions.
30 More specifically, as a random variable, normalized ergodic integrals converge in
31 distribution along certain subsequences to a non-degenerate, compactly supported
32 measure on the real line.

33 As a minor application, it is not surprising that the related result for exponential
34 sums appeared from connections between the Heisenberg group and theta series in
35 several contexts. For Heisenberg flows, as a first return map on a transverse torus
36 is obtained as a skew-translation, its limit theorem of theta sums was studied by J.
37 Griffin J. Marklof and Cellarosi-Marklof [GM14, CM16]. It is recently generalized by
38 Forni and Kanigowski [FK20b]. Likewise, we obtain limit theorems for theta series
39 on Siegel half spaces which generalize results [GG04, Mar99] (see [Tol78, MM07,
40 MNN07] for a general introduction).

41 Furthermore, L^2 -lower bound of Bufetov functional on a transverse torus and
42 its analyticity on the higher dimensional rectangular domain follows. This part
43 contributes that our functional does not vanish and extends to the complex domain
44 in the weighted space of fast-decaying coefficient of functionals. From these results,

polynomial type of lower bounds for the analytic function is obtained on certain sub-level sets. Such polynomial estimates on the measure are derived by real analyticity of a functional along the leaves of a foliation transverse to the actions based on results of [Bru99]. This tool was already devised by Forni and Kanigowski [FK20b] to study the bound of correlations for time-changes of nilflows. Unfortunately, it turned out that we can not obtain similar results because the time-changes of higher rank abelian actions on Heisenberg nilmanifolds are all trivial. It follows from the triviality of the first cohomology group (see [CF15, Theorem 3.16]) thus, they are conjugated to the linear action and never mixing.

As a rank 1-action, mixing properties for time changes of Heisenberg nilflows were firstly studied in [AFU11]. Then, it was extended in the dense set of non-trivial time changes for any uniquely ergodic nilflows on general nilmanifolds [Rav18, AFRU21]. As a special case, on time-changes of Heisenberg flows, multiple mixing [FK20a] was proved by Ratner property. In \mathbb{Z}^k -actions, mixing of shapes for automorphisms on nilmanifold is proved in [GS14, GS15].

Comprehensive studies of spatial and temporal limit theorems for horocycle flows have been carried out in the work of D. Dolgopyat and O. Sarig [DS17]. Recent work of Ravotti [Rav21] proved spatial and temporal limit theorems for horocycle flows in alternative ways by using Ratner's argument. His work did not rely on the methods from invariant distributions of solving the cohomological equations. However, it is still unknown if temporal (or weaker) limit theorems for Heisenberg flows can be proved, and it will be a possible subject of further works.

1.2. Definitions and statement of results. We review definitions about Heisenberg manifold and its moduli space.

1.2.1. Heisenberg manifold. Let $g \geq 1$ and H^g be the standard $2g + 1$ dimensional Heisenberg group and set $\Gamma := \mathbb{Z}^g \times \mathbb{Z}^g \times \frac{1}{2}\mathbb{Z}$ a discrete and co-compact subgroup of H^g . We shall call it standard lattice of H^g and the quotient $M := H^g/\Gamma$ will be called *Heisenberg manifold*. Lie algebra $\mathfrak{h}^g = Lie(H^g)$ is equipped with a basis $(X_1, \dots, X_g, Y_1, \dots, Y_g, Z)$ satisfying canonical commutation relations

$$(1) \quad [X_i, Y_j] = \delta_{ij}Z, \quad \text{where } \delta_{ij} = 1 \text{ if } i = j, \text{ and } \delta_{ij} = 0 \text{ otherwise.}$$

For $1 \leq d \leq g$, let $P := P^d$ be a subgroup of H^g where its Lie algebra $\mathfrak{p} := Lie(P^d)$ is generated by $\{X_1, \dots, X_d\}$. For any $\alpha \in Sp_{2g}(\mathbb{R})$, set $(X_i^\alpha, Y_i^\alpha, Z) = \alpha^{-1}(X_i, Y_i, Z)$ for $1 \leq i \leq d$. We define a parametrization of the subgroup $P^{d,\alpha} = \alpha^{-1}(P^d)$ according to

$$P_x^{d,\alpha} := \exp(x_1 X_1^\alpha + \dots + x_d X_d^\alpha), \quad x = (x_1, \dots, x_d) \in \mathbb{R}^d.$$

By central extension of \mathbb{R}^{2g} by \mathbb{R} , we have an exact sequence

$$0 \rightarrow Z(H^g) \rightarrow H^g \rightarrow \mathbb{R}^{2g} \rightarrow 0.$$

The natural projection $pr : M \rightarrow H^g/(\Gamma Z(H^g))$ maps M onto a $2g$ -dimensional torus $\mathbb{T}^{2g} := \mathbb{R}^{2g}/\mathbb{Z}^{2g}$.

1.2.2. Moduli space. The group of automorphisms of H^g that are trivial on the center is denoted by $Aut_0(H^g) = Sp_{2g}(\mathbb{R}) \ltimes \mathbb{R}^{2g}$. Since dynamical properties of actions are invariant under inner automorphism, we restrict our interest to $Sp_{2g}(\mathbb{R})$. We regard $Sp_{2g}(\mathbb{R})$ as the *deformation space* of the standard Heisenberg manifold M and call the quotient $\mathfrak{M}_g = Sp_{2g}(\mathbb{R})/Sp_{2g}(\mathbb{Z})$ the *moduli space* of (standard) Heisenberg manifold.

1 *Siegel modular variety* is a double coset space $\Sigma_g = K_g \backslash Sp_{2g}(\mathbb{R}) / Sp_{2g}(\mathbb{Z})$ where
 2 K_g is a maximal compact subgroup $Sp_{2g}(\mathbb{R}) \cap SO_{2g}(\mathbb{R})$ of $Sp_{2g}(\mathbb{R})$. For $\alpha \in$
 3 $Sp_{2g}(\mathbb{R})$, we denote $[\alpha] := \alpha Sp_{2g}(\mathbb{Z})$ by its projection on the moduli space \mathfrak{M}_g
 4 and write $[[\alpha]] := K_g \alpha Sp_{2g}(\mathbb{Z})$ the projection of α to the Siegel modular variety
 5 Σ_g .

6 Double coset $K_g \backslash Sp_{2g}(\mathbb{R}) / 1_{2g}$ is identified to the Siegel upper half space $\mathfrak{H}_g :=$
 7 $\{Z \in Sym_g(\mathbb{C}) \mid \Im(Z) > 0\}$. *Siegel upper half space* of genus g is complex manifold
 8 of symmetric complex $g \times g$ matrices $Z = X + iY$ with positive-definite symmetric
 9 imaginary part $\Im(Z) = Y$ and arbitrary (symmetric) real part X . Denote by
 10 $\Sigma_g \approx Sp_{2g}(\mathbb{Z}) \backslash \mathfrak{H}_g$.

11 1.2.3. *Representation.* We write the Hilbert sum decomposition of

$$(2) \quad L^2(M) = \bigoplus_{n \in \mathbb{Z}} H_n$$

into closed H^g -invariant subspaces. Set $f = \sum_{n \in \mathbb{Z}} f_n \in L^2(M)$ and $f_n \in H_n$ where

$$H_n = \{f \in L^2(M) \mid \exp(tZ)f = \exp(2\pi i n K t)f\}$$

12 for some fixed $K > 0$. The center $Z(H^g)$ has spectrum $2\pi\mathbb{Z} \setminus \{0\}$ and the space
 13 $L^2(M)$ splits as Hilbert sum of H^g -module H_n , which is equivalent to irreducible
 14 representation π .

By Stone-Von Neumann theorem, the unitary irreducible representation π of the
 Heisenberg group of non-zero central parameter $K > 0$ is unitarily equivalent to the
 Schrödinger representation π . By differentiating the Schrödinger representation, we
 obtain a representation of the Lie algebra \mathfrak{h}^g on Schwartz space $\mathcal{S}(\mathbb{R}^g) \subset L^2(\mathbb{R}^g)$
 (as a C^∞ -vector). This is called *infinitesimal derived representation* $d\pi_*$ with pa-
 rameter $n \in \mathbb{Z}$, and for each $k = 1, 2, \dots, g$,

$$d\pi_*(X_k) = \frac{\partial}{\partial x_k}, \quad d\pi_*(Y_k) = 2\pi i n K x_k, \quad d\pi_*(Z) = 2\pi i n K$$

15 acts on $L^2(\mathbb{R}^g) \simeq L^2(H_n)$.

16 Given a basis (V_i) of the Lie algebra, we set a Laplacian $\Delta = -\sum_i V_i^2$ and define
 17 L^2 -Sobolev norm $\|f\|_s^2 = \langle f, (1 + \Delta)^s f \rangle$ where $\langle \cdot, \cdot \rangle$ is an ordinary inner product.
 18 For $s > 0$, the *Sobolev space* $W^s(M)$ is defined by a completion of $C^\infty(M)$ equipped
 19 with the norm $\|\cdot\|_s$. The Sobolev space $W^s(M) = \bigoplus_{n \in \mathbb{Z}} W^s(H_n)$ decomposes to
 20 closed H^g -invariant subspaces $W^s(H_n) = W^s(M) \cap H_n$.

21 1.2.4. *Renormalization flow.* Denote diagonal matrix $\delta_i = \text{diag}(d_1, \dots, d_g)$ with
 22 $d_i = 1$, $d_k = 0$ if $k \neq i$. Then, for each $1 \leq i \leq g$, we denote $\hat{\delta}_i = \begin{bmatrix} \delta_i & 0 \\ 0 & -\delta_i \end{bmatrix} \in$
 23 $\mathfrak{sp}_{2g} = \text{Lie}(Sp_{2g}(\mathbb{R}))$.

24 Any such $\hat{\delta}_i$ generate a one-parameter subgroup of automorphism (renormaliza-
 25 tion flow) $r_i^t := e^{t\hat{\delta}_i}$. We denote (rank d) renormalization actions $r_t := r_{i_1}^{t_1} \cdots r_{i_d}^{t_d}$
 26 for $t = (t_1, \dots, t_d)$ and $1 \leq i_1, \dots, i_d \leq g$. We also write the corresponding auto-
 27 morphism

$$(3) \quad \exp(t\hat{\delta}(d)) : (x, y, z) \mapsto (e^{t\hat{\delta}}x, e^{-t\hat{\delta}}y, z)$$

28 of the Heisenberg group (see §2.2.3 for its use).

1 **Main results.** One of the main objects in this paper is to construct finitely-
 2 additive measures defined on the space of rectangles on Heisenberg manifold M .
 3 We state our results beginning with an overview of Bufetov functional.

4 **Definition 1.1.** For $(m, \mathbf{T}) \in M \times \mathbb{R}_+^d$, denote the *standard rectangle* for action P ,

$$(4) \quad \Gamma_{\mathbf{T}}^X(m) = \{P_{\mathbf{t}}^{d,\alpha}(m) \mid \mathbf{t} \in U(\mathbf{T}) = [0, \mathbf{T}^{(1)}] \times \cdots \times [0, \mathbf{T}^{(d)}]\}.$$

5 Let $Q_y^{d,Y} := \exp(y_1 Y_1 + \cdots + y_d Y_d)$, $y = (y_1, \dots, y_d) \in \mathbb{R}^d$ be an action generated
 6 by elements Y_i of standard basis. Let $\phi_z^Z := \exp(zZ)$, $z \in \mathbb{R}$ be a flow generated
 7 by the central element Z .

8 **Definition 1.2.** Let \mathfrak{R} be the collection of the *generalized rectangles* in M . For
 9 any $1 \leq j \leq d \leq g$ and $\mathbf{t} = (t_1, \dots, t_d)$, we set

$$(5) \quad \mathfrak{R} := \bigcup_{1 \leq i \leq d} \bigcup_{(y,z) \in \mathbb{R}^j \times \mathbb{R}} \bigcup_{(m,\mathbf{T}) \in M \times \mathbb{R}_+^d} \{(\phi_{t_i z}^Z) \circ Q_y^{j,Y} \circ P_{\mathbf{t}}^{d,\alpha}(m) \mid \mathbf{t} \in U(\mathbf{T})\}.$$

10 **Theorem 1.3.** For any irreducible representation H , there exists a finitely-additive
 11 measure $\hat{\beta}_H(\Gamma) \in \mathbb{C}$ defined on every standard d -rectangle Γ such that the following
 12 holds:

(1) (Additive property) For any decomposition of disjoint rectangles $\Gamma = \bigcup_{i=1}^n \Gamma_i$
 or whose intersections have zero measure,

$$\hat{\beta}_H(\alpha, \Gamma) = \sum_{i=1}^n \hat{\beta}_H(\alpha, \Gamma_i).$$

(2) (Scaling property) For $\mathbf{t} \in \mathbb{R}^d$,

$$\hat{\beta}_H(r_{\mathbf{t}}(\alpha), \Gamma) = e^{-(t_1 + \cdots + t_d)/2} \hat{\beta}_H(\alpha, \Gamma).$$

(3) (Invariance property) For any action $Q_{\tau}^{j,Y}$ generated by Y_i 's for $\tau \in \mathbb{R}_+^j$
 and $1 \leq j \leq d$,

$$\hat{\beta}_H(\alpha, (Q_{\tau}^{j,Y})_* \Gamma) = \hat{\beta}_H(\alpha, \Gamma).$$

(4) (Bounded property) For any rectangle $\Gamma \in \mathfrak{R}$, there exists a constant $C(\Gamma) >$
 0 such that for $\hat{X} = \hat{X}_1 \wedge \cdots \wedge \hat{X}_d$,

$$|\hat{\beta}_H(\alpha, \Gamma)| \leq C(\Gamma) \left(\int_{\Gamma} |\hat{X}| \right)^{d/2}.$$

13 **Corollary 1.4.** The functional $\hat{\beta}_H$ defined on standard rectangle $\Gamma_{\mathbf{T}}^X$ extends to the
 14 class \mathfrak{R} .

15 **Remark.** By the additive property of functionals on rectangles, it is plausible to
 16 extend the shape of domain to a general boundary by approximating with rectangles.
 17 However, this method may cause limitations by having a weaker estimate of equidis-
 18 tributions. It may be also interesting to compare the methods of Shah [S09a, S09b]
 19 and to obtain equidistribution results for smooth (or Lipschitz) boundary in our
 20 settings.

21 Let us consider arbitrary two d -standard rectangles $U(\mathbf{T}_1)$ and $U(\mathbf{T}_2)$. For
 22 convenience, these rectangles are translated to intersect at only one vertex as in the
 23 figure. Without loss of generality, assume that d distinct faces of $U(\mathbf{T}_1)$ emanate




FIGURE 1. Illustration of the rectangles $U(\mathbf{T}_1)$ and $U(\mathbf{T}_2)$ on i, j -th coordinate.

- 1 from the origin and $U(\mathbf{T}_1) \cap U(\mathbf{T}_2) = \{(\mathbf{T}_1^{(l)})_{1 \leq l \leq d}\}$. Denote by $\mathbf{P}(\mathbf{T}_1)$ a collection
 2 of 2^d vertices $v = (v^{(1)}, v^{(2)}, \dots, v^{(d)}) \in \mathbb{R}_+^d$ of $U(\mathbf{T}_1)$.

Then we define a vector $\mathbf{T}_v = (\mathbf{T}_v^{(l)}) \in \mathbb{R}_+^d$ associated with v given by

$$\mathbf{T}_v^{(l)} := \begin{cases} \mathbf{T}_2^{(l)} & \text{if } v^{(l)} = \mathbf{T}_1^{(l)} \\ \mathbf{T}_1^{(l)} & \text{if } v^{(l)} = 0. \end{cases}$$

- 3 **Corollary 1.5.** *Let us denote*

$$(6) \quad \beta_H(\alpha, m, \mathbf{T}) := \hat{\beta}_H(\alpha, \Gamma_{\mathbf{T}}^X(m)).$$

- 4 *The function β_H satisfies the following properties:*

- (1) (Cocycle property) *For all $(m, \mathbf{T}_1, \mathbf{T}_2) \in M \times \mathbb{R}_+^d \times \mathbb{R}_+^d$,*

$$\beta_H(\alpha, m, \mathbf{T}_1 + \mathbf{T}_2) = \sum_{v \in \mathbf{P}(\mathbf{T}_1)} \beta_H(\alpha, \mathbf{P}_v^{d, \alpha}(m), \mathbf{T}_v).$$

- (2) (Scaling property) *For all $m \in M$ and $\mathbf{t} = (t_1, \dots, t_d)$,*

$$\beta_H(r_{\mathbf{t}}(\alpha), m, \mathbf{T}) = e^{-(t_1 + \dots + t_d)/2} \beta_H(\alpha, m, \mathbf{T}).$$

- (3) (Bounded property) *Let us denote the largest length of edges of $U(\mathbf{T})$ by $T_{\max} = \max_{1 \leq i \leq d} \mathbf{T}^{(i)}$. Then there exists a constant $C_H > 0$ such that*

$$\beta_H(\alpha, m, \mathbf{T}) \leq C_H T_{\max}^{d/2}.$$

- (4) (Orthogonality) *For all $\mathbf{T} \in \mathbb{R}^d$, $\beta_H(\alpha, \cdot, \mathbf{T})$ belongs to an irreducible component, i.e.,*

$$\beta_H(\alpha, \cdot, \mathbf{T}) \in H \subset L^2(M).$$

- 5 By representation theory introduced in §1.2.3, for any $f = \sum_H f_H \in W^s(M)$,
 6 define a *Bufetov cocycle* associated to f as a sum

$$(7) \quad \beta^f(\alpha, m, \mathbf{T}) = \sum_H D_{\alpha}^H(f) \beta_H(\alpha, m, \mathbf{T}).$$

Notation. Let \hat{X}_i^{α} be a 1-form dual to the vector field X_i^{α} , in the sense that $\hat{X}_i^{\alpha}(X_i^{\alpha}) = 1$ and $\hat{X}_i^{\alpha}(X_j^{\alpha}) = 0$ if $i \neq j$ on M . Given a Jordan region U and a point $m \in M$, set $\mathcal{P}_U^{d, \alpha} m$ the *Birkhoff integrals* (currents) associated to the action $\mathbf{P}_x^{d, \alpha}$ given by

$$\langle \mathcal{P}_U^{d, \alpha} m, \omega_f \rangle := \int_U f(\mathbf{P}_x^{d, \alpha} m) dx_1 \cdots dx_d$$

1 for any degree d \mathfrak{p} -form $\omega_f = f \hat{X}_1^\alpha \wedge \cdots \wedge \hat{X}_d^\alpha$ with a smooth function $f \in C_0^\infty(M)$
 2 with zero averages.

3 **Theorem 1.6.** *For all $s > d(d+11)/4 + g + 1$, there exists a constant $C_s > 0$ such*
 4 *that for almost all frequency α and for all $f \in W^s(M)$ and for all $(m, \mathbf{T}) \in M \times \mathbb{R}^d$,*
 5 *we have*

$$(8) \quad \left| \left\langle \mathcal{P}_{U(\mathbf{T})}^{d,\alpha} m, \omega_f \right\rangle - \beta^f(\alpha, m, \mathbf{T}) \right| \leq C_s \|\omega_f\|_s.$$

6 for $U(\mathbf{T}) = [0, T^{(1)}] \times \cdots \times [0, T^{(d)}]$ and $\omega_f = f \omega^{d,\alpha} \in \Lambda^d \mathfrak{p} \otimes W^s(M)$.

The family of random variable

$$E_{\mathbf{T}_n}(f) := \frac{1}{\text{vol}(U(\mathbf{T}_n))^{1/2}} \left\langle \mathcal{P}_{U(\mathbf{T}_n)}^{d,\alpha}(m), \omega_f \right\rangle$$

7 is defined where $U(\mathbf{T}_n)$ is a sequence of rectangles. The point $m \in M$ is distributed
 8 accordingly to the probability measure vol on M . Our goal is to understand the
 9 asymptotic behavior of the probability distributions of $E_{\mathbf{T}_n}(f)$ as $U(\mathbf{T}_n) \nearrow \mathbb{R}^d$ in
 10 a sense of Følner.

Theorem 1.7. *Let $\{\mathbf{T}_n\}$ be any sequence such that*

$$\lim_{n \rightarrow \infty} r_{\log \mathbf{T}_n}[\alpha] = \alpha_\infty \in \mathfrak{M}_g.$$

11 *For every closed form $\omega_f \in \Lambda^d \mathfrak{p} \otimes W^s(M)$ with $s > d(d+11)/4 + g + 1$, which is not*
 12 *a coboundary, the limit distribution of the family of random variables $E_{\mathbf{T}_n}(f)$ exists*
 13 *along a subsequence of $\{\mathbf{T}_n\}$. In particular, for almost all α , the limit distribution*
 14 *of $E_{\mathbf{T}_n}(f)$ has compact support.*

15 We finish the section by giving some remarks on the new adaptation of transfer
 16 operator techniques from hyperbolic dynamics. The method stems from the analysis
 17 of the transfer operator, firstly treated by P. Gieulietti and Liverani [GL19]. They
 18 set up a non-linear flow on the torus and proved asymptotics of ergodic averages
 19 with expansions of invariant distributions and eigenvalues of transfer operators
 20 called *Ruelle resonances* (see also [AB18, FGL19, For20b, B19] for applications in
 21 parabolic flows).

22 A recent work of L. Simonelli and O. Butterley [BS20] reproved some of the re-
 23 sults of Flaminio and Forni [FF06] by analytical methods of hyperbolic theory, not
 24 relying on representation theory. Although their work is restricted to the periodic
 25 type of α in the moduli space \mathfrak{M} (the flow is only renormalized by partially hyper-
 26 bolic diffeomorphism, or α is periodic) their methods showed indirect similarities
 27 with [FK20b] and our current work.¹ However, it is still not known if it is possible to
 28 extend their approach to full measure set of α , requiring construction of renormal-
 29 ization cocycle, so called *Transfer cocycle*. It is also not well-studied how to replace
 30 the use of Anisotropic norm, not relying on the result of Faure-Tsujii [FT15].²

31 On higher step nilmanifolds, there does not exist a renormalization flow (mod-
 32 uli space \mathfrak{M} is trivial). It is not possible to construct the Bufetov functionals

¹For instance, it is plausible to view their construction of functionals obtained by spectral projection as a functional $\hat{\beta}_H$ on each sub-representation H in our setting (see [BS20, §4]).

²Furthermore, it is unknown how to set Diophantine conditions for the recurrence of frequency α . Instead of relying on the methods in Sobolev constant techniques introduced in [FF06], we do not have proper tools on Anisotropic spaces yet.

1 by the same strategies introduced in here, but other methods in handling non-
 2 renormalizable flows are possibly applied (cf. [FF14, FFT16, Kim21]).

3 *Outline of the paper.* In section 2, we give basic definitions on higher rank actions,
 4 moduli spaces and Sobolev spaces. In section 3, we state main theorem and prove
 5 constructions of Bufetov functionals with main properties. In section 4, we prove
 6 asymptotic formula of ergodic integrals and limit theorems. In section 5, we prove
 7 L^2 -lower bound of the Bufetov functionals. In Section 6, analyticity of functional
 8 and extensions of domain are provided. In section 7, there exist measure estimates
 9 of functionals on the sets where values of functionals are small. This result only
 10 holds when frame α is of bounded type.

11 2. ANALYSIS ON HEISENBERG MANIFOLDS

12 In this section, we review definitions of Sobolev space, currents, representation
 13 and renormalization flows on moduli space.

14 2.1. Sobolev space.

2.1.1. *Sobolev norm.* Given a basis (V_i) of the Lie algebra \mathfrak{h}^g , we write a new basis
 $((\alpha^{-1})_* V_i)$ for $\alpha \in \text{Aut}_0(\mathcal{H}^g)$. Similarly, denote by Laplacian $\Delta_\alpha = -\sum_i (\alpha^{-1})_* V_i^2$
 with respect to the new basis. For any $s \in \mathbb{R}$ and any C^∞ -function $f \in L^2(M)$,
Sobolev norm is defined by

$$\|f\|_{\alpha,s} = \langle f, (1 + \Delta_\alpha)^s f \rangle^{1/2}.$$

15 Let $W_\alpha^s(M)$ be a completion of $C^\infty(M)$ with the norm above. The dual space of
 16 $W_\alpha^s(M)$ is denoted by $W_\alpha^{-s}(M)$ and it is isomorphic to $W_\alpha^s(M)$. Extending it to the
 17 exterior algebra, define the Sobolev spaces of the form $\Lambda^d \mathfrak{p} \otimes W_\alpha^s(M)$ of cochains
 18 of degree d and we use the same notations for the norm.

2.1.2. *Sobolev bundle.* The group $Sp_{2g}(\mathbb{Z})$ acts on the right on the trivial bundles

$$Sp_{2g}(\mathbb{R}) \times W^s(M) \rightarrow Sp_{2g}(\mathbb{R})$$

where

$$(\alpha, \varphi) \mapsto (\alpha, \varphi)\gamma = (\alpha\gamma, \gamma^* \varphi), \quad \gamma \in Sp_{2g}(\mathbb{Z}).$$

We obtain the quotient flat bundle of Sobolev spaces over the moduli space:

$$(Sp_{2g}(\mathbb{R}) \times W^s(M))/Sp_{2g}(\mathbb{Z}) \rightarrow \mathfrak{M}_g = Sp_{2g}(\mathbb{R})/Sp_{2g}(\mathbb{Z})$$

19 and the fiber over $[\alpha] \in \mathfrak{M}_g$ is locally identified with the space $W_\alpha^s(M)$.

By invariance of $Sp_{2g}(\mathbb{Z})$ action, the class of (α, φ) is denoted by $[\alpha, \varphi]$ and
 $Sp_{2g}(\mathbb{Z})$ -invariant Sobolev norm is written by

$$\|f\|_{\alpha,s} := \|[\alpha, f]\|_s.$$

We denote the bundle of \mathfrak{p} -forms of degree j of Sobolev order s by $A^j(\mathfrak{p}, \mathfrak{M}^s)$.
 The space of continuous linear functional on $A^j(\mathfrak{p}, \mathfrak{M}^s)$ will be called the *space*
of currents of dimension j and denoted by $A_j(\mathfrak{p}, \mathfrak{M}^{-s})$. There is a flat bundle of
 (currents) distribution $A_j(\mathfrak{p}, \mathfrak{M}^{-s})$ whose fiber over $[\alpha]$ is locally identified with
 the space $W_\alpha^{-s}(M)$. Likewise, the space of Sobolev currents $A_j(\mathfrak{p}, W^{-s}(M))$ of
 dimension j with order s is identified with $\Lambda^j \mathfrak{p} \otimes W^{-s}(M)$. We write the norm for
 form ω and currents \mathscr{D} by

$$\|\omega\|_{\alpha,s} := \|[\alpha, \omega]\|_s, \quad \|\mathscr{D}\|_{\alpha,s} := \|[\alpha, \mathscr{D}]\|_s.$$

2.1.3. *Best Sobolev constant.* The *Sobolev embedding theorem* implies that for any $\alpha \in Sp_{2g}(\mathbb{R})$ and $s > g + 1/2$, there exists a constant $B_s(\alpha)$ such that for any $f \in W_\alpha^s(M)$,

$$(9) \quad B_s(\alpha) := \sup_{f \in W_\alpha^s(M) \setminus \{0\}} \frac{\|f\|_\infty}{\|f\|_{\alpha,s}}.$$

By Lemma 4.4 in [CF15], the *best Sobolev constant* B_s is a $Sp_{2g}(\mathbb{Z})$ -modular function on \mathfrak{H}_g . Thus, we shall write $B_s([\alpha])$ or $B_s(\alpha)$ for $B_s(\alpha)$. By the Sobolev embedding theorem, we have a bound for the Birkhoff integral current.

Lemma 2.1. [CF15, Lemma 5.5] *For any Jordan region $U \subset \mathbb{R}^d$ with Lebesgue measure $|U|$, for any $s > g + 1/2$ and all $m \in M$,*

$$\left\| [\alpha, \mathcal{P}_U^{d,\alpha} m] \right\|_{-s} \leq B_s([\alpha])|U|.$$

2.2. Invariant currents.

2.2.1. *Identification.* The *boundary operators*

$$\partial : A_j(\mathfrak{p}, W^{-s}(M)) \rightarrow A_{j-1}(\mathfrak{p}, W^{-s}(M))$$

are adjoint of the differentials d such that $\langle \partial \mathcal{D}, \omega \rangle = \langle \mathcal{D}, d\omega \rangle$ for any $\omega \in \Lambda^{j-1} \mathfrak{p} \otimes W^s(M)$. A current \mathcal{D} is called *closed* if $\partial \mathcal{D} = 0$.

For $s > 0$, we denote $Z_d(\mathfrak{p}, W^{-s}(M))$ by the space of closed currents of dimension d and Sobolev order s . $I_d(\mathfrak{p}, W^{-s}(M))$ is denoted by the d -dimensional space of \mathbf{P} -invariant currents of Sobolev order s . Now we find the relations between these two currents.

Proposition 2.2. [CF15, Proposition 3.13] *For any $s > d/2 = \dim \mathbf{P}/2$, we have $I_d(\mathfrak{p}, \mathcal{S}(\mathbb{R}^g)) \subset W^{-d/2-\epsilon}(\mathbb{R}^g)$ for all $\epsilon > 0$. Additionally,*

- $I_d(\mathfrak{p}, \mathcal{S}(\mathbb{R}^g))$ is one dimensional space if $\dim \mathbf{P} = g$,
- $I_d(\mathfrak{p}, \mathcal{S}(\mathbb{R}^g))$ is an infinite-dimensional space if $\dim \mathbf{P} < g$,
- $I_d(\mathfrak{p}, \mathcal{S}(\mathbb{R}^g)) = Z_d(\mathfrak{p}, W^{-s}(M))$ for any $1 \leq d \leq g$.

2.2.2. *Basic currents.* The current B_α is called *basic* if for all $j \in \{i_1, \dots, i_d\}$,

$$\iota_{X_j} B_\alpha = L_{X_j} B_\alpha = 0.$$

For an irreducible representation H , there exists a unique *basic current* B_α^H (of degree $2g + 1 - d$ and dimension d) associated to invariant distribution D_α^H . It is defined by $B_\alpha^H = D_\alpha^H \eta_X$ and this formula implies that for every d -form ξ ,

$$(10) \quad B_\alpha^H(\xi) = D_\alpha^H \left(\frac{\eta_X \wedge \xi}{\omega_{vol}} \right)$$

where $\eta_X := \iota_{X_{i_1}} \cdots \iota_{X_{i_d}} \omega_{vol}$ and ω_{vol} is an invariant volume form (with total unit volume).

The basic current B_α^H belongs to the Sobolev space of currents and it is \mathbf{P} -invariant. It follows that for all $s > d/2$, by Sobolev embedding theorem, $B_\alpha^H \in A_j(\mathfrak{p}, W_\alpha^{-s}(M))$ if and only if $D_\alpha^H \in W_\alpha^{-s}(M)$ for all $s > d/2$.

Remark. The formula (10) yields an isomorphism between the space of basic (closed) currents and invariant distributions (see also [For02, §6.1] and [BF14,

§2.1)). For any d -dimensional \mathfrak{p} -form $\omega_f = f \hat{X}_1 \wedge \cdots \wedge \hat{X}_d$ with $f \in C^\infty(M)$, we will identify currents \mathcal{D} with distribution D by writing

$$\langle D, f \rangle = \langle \mathcal{D}, \omega_f \rangle.$$

2.2.3. *Renormalization.* Let $s > d/2$. Recall the definition of renormalization flow (3) in §1.2.4. For $\omega \in \Lambda^d \mathfrak{p} \otimes W^s(\mathbb{R}^g)$, $\mathcal{D} \in Z_d(\mathfrak{p}, W^{-s}(M))$, and $t \in \mathbb{R}$

$$r_i^t[\alpha, \omega] = [r_i^t \alpha, \omega], \quad r_i^t[\alpha, \mathcal{D}] = [r_i^t \alpha, \mathcal{D}].$$

- 1 By Proposition 5.2 in [CF15], the sub-bundle $Z_d(\mathfrak{p}, W^{-s}(M))$ is invariant under
2 the renormalization flows r_i^t . Furthermore, we have

$$\|r_1^{t_1} \cdots r_d^{t_d}[\alpha, \mathcal{D}]\|_{-s} = e^{-(t_1 + \cdots + t_d)/2} \|[\alpha, \mathcal{D}]\|_{-s}.$$

- 3 Let $U_{\mathbf{t}} : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$ be unitary operator for $\mathbf{t} = (t_1, \dots, t_d)$,

$$(11) \quad U_{\mathbf{t}} f(x) = e^{-(t_1 + \cdots + t_d)/2} f(e^{t_1} x_1, \dots, e^{t_d} x_d).$$

Then, for any invariant distribution D_α^H ,

$$D_{r_{\mathbf{t}}(\alpha)}^H = e^{(t_1 + \cdots + t_d)/2} D_\alpha^H.$$

- 4 By reparametrization of \mathbb{R}^d -action $r_{\mathbf{t}} = r_1^{t_1} \cdots r_d^{t_d}$,

$$(12) \quad \mathbf{P}_x^{d, (r_1^{t_1} \cdots r_d^{t_d} \alpha)} = \mathbf{P}_{(e^{-t_1} x_1, \dots, e^{-t_d} x_d)}^{d, \alpha}.$$

- 5 Then the Birkhoff integral current also satisfies the following identity

$$(13) \quad \mathcal{P}_U^{d, (r_1^{t_1} \cdots r_d^{t_d} \alpha)} m = e^{(t_1 + \cdots + t_d)/2} \mathcal{P}_{(e^{-t_1}, \dots, e^{-t_d})U}^{d, \alpha} m.$$

6 2.3. Diophantine condition.

7 2.3.1. Height function.

Definition 2.3 (Height function). The *height* of a point $Z \in \mathfrak{H}_g$ in Siegel upper half space is the positive number

$$hgt(Z) := \det \Im(Z).$$

The *maximal height function* $Hgt : \Sigma_g \rightarrow \mathbb{R}^+$ is the maximal height of a $Sp_{2g}(\mathbb{Z})$ orbit of Z . That is, for the class of $[Z] \in \Sigma_g$,

$$Hgt([Z]) := \max_{\gamma \in Sp_{2g}(\mathbb{Z})} hgt(\gamma(Z)).$$

- 8 By Proposition 4.8 of [CF15], there exists a universal constant $C(s) > 0$ such
9 that the best Sobolev constant satisfies the estimate

$$(14) \quad B_s([\alpha]) \leq C(s) \cdot (Hgt([\alpha]))^{1/4}.$$

- 10 We rephrase Lemma 4.9 in [CF15] regarding the bound of renormalized height.

- 11 **Lemma 2.4.** For any $[\alpha] \in \mathfrak{M}_g$ and any $\mathbf{t} \in \mathbb{R}_+^d$,

$$(15) \quad Hgt([\exp(-\mathbf{t} \hat{\delta}(d)) \alpha]) \leq (\det(e^{\mathbf{t} \hat{\delta}}))^2 Hgt([\alpha]).$$

2.3.2. *Diophantine condition.*

Definition 2.5. An automorphism $\alpha \in Sp_{2g}(\mathbb{R})$ or a point $[\alpha] \in \mathfrak{M}_g$ is $\hat{\delta}(d)$ -*Diophantine* of type σ if there exists a $\sigma > 0$ and a constant $C > 0$ such that

$$(16) \quad Hgt([\exp(-t\hat{\delta}(d))\alpha]) \leq CHgt([\exp(-t\hat{\delta}(d))])^{(1-\sigma)} Hgt([\alpha]), \quad \forall t \in \mathbb{R}_+^d.$$

This states that $\alpha \in Sp_{2g}(\mathbb{R})$ satisfies $\hat{\delta}(d)$ -Diophantine if the height of the projection of $\exp(-t\hat{\delta}(d))\alpha$ in the Siegel modular variety Σ_g is bounded by $e^{2(t_1+\dots+t_d)(1-\sigma)}$. Furthermore,

- $[\alpha] \in \mathfrak{M}_g$ satisfies a $\hat{\delta}(d)$ -*Roth condition* if for any $\epsilon > 0$ there exists a constant $C > 0$ such that

$$(17) \quad Hgt([\exp(-t\hat{\delta}(d))\alpha]) \leq CHgt([\exp(-t\hat{\delta}(d))])^\epsilon Hgt([\alpha]), \quad \forall t \in \mathbb{R}_+^d.$$

That is, $\hat{\delta}(d)$ -Diophantine of type $0 < \sigma < 1$.

- $[\alpha]$ is of *bounded type* if there exists a constant $C > 0$ such that

$$(18) \quad Hgt([\exp(-t\hat{\delta}(d))]) \leq C, \quad \forall t \in \mathbb{R}_+^d.$$

Definition 2.6. Let $X = G/\Lambda$ be a homogeneous space equipped with the probability Haar measure μ . A function $\phi : X \rightarrow \mathbb{R}$ is said k -DL (distance like) for some exponent $k > 0$ if it is uniformly continuous and if there exist constants $C_1, C_2 > 0$ such that

$$C_1 e^{-kz} \leq \mu(\{x \in X \mid \phi(x) \geq z\}) \leq C_2 e^{-kz}, \quad \forall z \in \mathbb{R}.$$

By the work of Kleinblock and Margulis, a multi-parameter generalization of Khinchin-Sullivan logarithm law for geodesic excursion [Sul82] holds.

Theorem 2.7. [KM99, Theorem 1.9] Let G be a connected semisimple Lie group without compact factors, μ its normalized Haar measure, $\Lambda \subset G$ an irreducible lattice, \mathfrak{a} a Cartan subalgebra of the Lie algebra of G . Let \mathfrak{d}_+ be a non-empty open cone in a d -dimensional subalgebra \mathfrak{d} of \mathfrak{a} ($1 \leq d \leq \text{rank}_{\mathbb{R}}(G)$). If $\phi : G/\Lambda \rightarrow \mathbb{R}$ is a k -DL function for some $k > 0$, then for μ -almost all $x \in G/\Lambda$ one has

$$\limsup_{\mathbf{z} \in \mathfrak{d}_+, \mathbf{z} \rightarrow \infty} \frac{\phi(\exp(\mathbf{z})x)}{\log \|\mathbf{z}\|} = \frac{d}{k}$$

By Lemma 4.7 of [CF15], the logarithm of Height function is DL-function with exponent $k = \frac{g+1}{2}$ on the Siegel variety Σ_g (and induces on $\mathfrak{M}_g = Sp_{2g}(\mathbb{R})/Sp_{2g}(\mathbb{Z})$). Hence, we obtain the following proposition.

Proposition 2.8. Under the assumption $X = \mathfrak{M}_g$ of Theorem 2.7, for $s > g+1/2$, there exists a full measure set $\Omega_g(\hat{\delta})$ and for all $[\alpha] \in \Omega_g(\hat{\delta}) \subset \mathfrak{M}_g$

$$(19) \quad \limsup_{t \in \mathbb{R}_+^d, t \rightarrow \infty} \frac{\log Hgt([\exp(-t\hat{\delta}(d))\alpha])}{\log \|t\|} \leq \frac{2d}{g+1}.$$

Any such $[\alpha]$ satisfies a $\hat{\delta}(d)$ -Roth condition (17).

For any $L > 0$ and $1 \leq d \leq g$, let $DC(L)$ denote the set of $[\alpha] \in \mathfrak{M}_g$ such that

$$(20) \quad \int_0^\infty \dots \int_0^\infty e^{-(t_1+\dots+t_d)/2} Hgt([r_{-\mathbf{t}}(\alpha)])^{1/4} dt_1 \dots dt_d \leq L.$$

Let DC denote the union of the sets $DC(L)$ over all $L > 0$. It follows immediately that the set $DC \subset \mathfrak{M}_g$ has full Haar volume.

Remark. In the work of Cosentino-Flaminio [CF15], Diophantine condition is restricted to one-parameter subgroup $\exp(-t\hat{\delta})$ for non-negative (g -dimensional) diagonal directions. This is based on the easy part of theorem in Kleinblock and Margulis [KM99, Theorem 1.7]. Our condition is for d -dimensional renormalizations actions. We will write the notation r_{-t} instead of $\exp(-t\hat{\delta}(d))$ for later sections.

3. CONSTRUCTIONS OF THE FUNCTIONALS

In this section, we construct Bufetov functionals for higher rank actions on the standard rectangular domain. Firstly, we prove the bound of Birkhoff integrals on rectangles. It consists of similar proofs of Theorem 5.10 in [CF15], but this type of estimate will be used to control remainder estimate (Lemma 3.4) to define functionals under Diophantine condition (Lemma 3.5). Furthermore, we will verify the properties of functionals $\hat{\beta}_H$ and cocycles β_H stated in Theorem 1.3.

3.1. Remainder estimates. For any exponent $s > d/2$, Hilbert bundle induces an orthogonal decomposition

$$A_d(\mathfrak{p}, \mathfrak{M}^{-s}) = Z_d(\mathfrak{p}, \mathfrak{M}^{-s}) \oplus R_d(\mathfrak{p}, \mathfrak{M}^{-s})$$

where $R_d(\mathfrak{p}, \mathfrak{M}^{-s}) = Z_d(\mathfrak{p}, \mathfrak{M}^{-s})^\perp$. Denote by \mathcal{I}^{-s} and \mathcal{R}^{-s} the corresponding orthogonal projection operator and by \mathcal{I}_α^{-s} and \mathcal{R}_α^{-s} the restrictions to the fiber over $[\alpha] \in \mathfrak{M}$ for $\alpha \in Sp_{2g}(\mathbb{R})$. In particular, for the current (Birkhoff integrals) $\mathcal{D} = \mathcal{P}_U^{d,\alpha} m$, we call $\mathcal{I}_\alpha^{-s}(\mathcal{D}) = \mathcal{I}^{-s}[\alpha, \mathcal{D}]$ boundary term and $\mathcal{R}_\alpha^{-s}(\mathcal{D}) = \mathcal{R}^{-s}[\alpha, \mathcal{D}]$ remainder term respectively. Consider the orthogonal projection

$$(21) \quad \mathcal{D} = \mathcal{I}_{r_{-t}[\alpha]}^{-s}(\mathcal{D}) + \mathcal{R}_{r_{-t}[\alpha]}^{-s}(\mathcal{D}).$$

We firstly recall the following estimate of boundary term.

Lemma 3.1. [CF15, Lemma 5.7] *Let $s > d/2 + 2$. There exists a constant $C = C(s) > 0$ such that for all $t_i \geq 0$ for $1 \leq i \leq d$, we have*

$$\begin{aligned} \|\mathcal{I}^{-s}[\alpha, \mathcal{D}]\|_{-s} &\leq e^{-(t_1 + \dots + t_d)/2} \|\mathcal{I}^{-s}[r_1^{-t_1} \dots r_d^{-t_d} \alpha, \mathcal{D}]\|_{-s} \\ &+ C_1 |t_1 + \dots + t_d| \int_0^1 e^{-u(t_1 + \dots + t_d)/2} \|\mathcal{R}^{-s}[r_1^{-ut_1} \dots r_d^{-ut_d} \alpha, \mathcal{D}]\|_{-(s-2)} du. \end{aligned}$$

By Stokes' theorem, we have the following remainder estimate.

Lemma 3.2. [CF15, Lemma 5.6] *Let $s > g + d/2 + 1$. For any non-negative $s' < s - (d+1)/2$ and Jordan region $U \subset \mathbb{R}^d$, there exists $C = C(d, g, s, s') > 0$ such that*

$$\|\mathcal{R}^{-s}[\alpha, (\mathcal{P}_U^{d,\alpha} m)]\|_{-s} \leq C \|\alpha, \partial(\mathcal{P}_U^{d,\alpha} m)\|_{-s'}.$$

Quantitative bound of Birkhoff integrals on the square domain was obtained in [CF15], but we need to extend the result to the rectangular shapes for analyticity of functionals in the section §6.

From now on, assume that $s_{d,g} := d(d+11)/4 + g + 1/2$.

Theorem 3.3. *For $s > s_{d,g}$, there exists a constant $C(s, d) > 0$ such that the following holds. For any $t_i > 0$, $m \in M$ and $U_d(t) = [0, e^{t_1}] \times \cdots \times [0, e^{t_d}]$, we have*

(22)

$$\begin{aligned} \left\| [\alpha, (\mathcal{P}_{U_d(t)}^{d,\alpha} m)] \right\|_{-s} &\leq C \sum_{k=0}^d \sum_{1 \leq i_1 < \cdots < i_k \leq d} \int_0^{t_{i_k}} \cdots \int_0^{t_{i_1}} \exp\left(\frac{1}{2} \sum_{l=1}^d t_l - \frac{1}{2} \sum_{l=1}^k u_{i_l}\right) \\ &\quad \times Hgt\left(\left[\prod_{1 \leq j \leq d} r_j^{-t_j} \prod_{l=1}^k r_{i_l}^{u_{i_l}} \alpha\right]\right)^{1/4} du_{i_1} \cdots du_{i_k}. \end{aligned}$$

- 1 *Proof.* We proceed by induction. For $d = 1$, it follows from the Theorem 5.8
 2 in [CF15]. We assume that the result holds for $d - 1$. Decompose the current as a
 3 sum of boundary and remainder term as in (21).

Step 1. We estimate the boundary term. By Lemma 3.1, renormalize terms with $r^u = r_1^u \cdots r_d^u$. Then, we have

$$\begin{aligned} \left\| \mathcal{I}^{-s}[\alpha, (\mathcal{P}_{U_d(t)}^{d,\alpha} m)] \right\|_{-s} &\leq e^{-(t_1 + \cdots + t_d)/2} \left\| \mathcal{I}^{-s}[r_1^{-t_1} \cdots r_d^{-t_d} \alpha, (\mathcal{P}_{U_d(t)}^{d,\alpha} m)] \right\|_{-s} \\ (23) \quad &+ C_1(s) \int_0^{t_1 + \cdots + t_d} e^{-ud/2} \left\| \mathcal{R}^{-s}[r^{-u} \alpha, (\mathcal{P}_{U_d(t)}^{d,\alpha} m)] \right\|_{-(s-2)} du \\ &:= (I) + (II). \end{aligned}$$

By renormalization (13) and Lemma 2.1 for unit volume,

$$\begin{aligned} \left\| \mathcal{I}^{-s}[r_1^{-t_1} \cdots r_d^{-t_d} \alpha, (\mathcal{P}_{U_d(t)}^{d,\alpha} m)] \right\|_{-s} &= e^{t_1 + \cdots + t_d} \left\| \mathcal{I}^{-s}[r_1^{-t_1} \cdots r_d^{-t_d} \alpha, (\mathcal{P}_{U_d(0)}^{d, r_1^{-t_1} \cdots r_d^{-t_d} \alpha} m)] \right\|_{-s} \\ &\leq C_2 e^{t_1 + \cdots + t_d} Hgt([r_1^{-t_1} \cdots r_d^{-t_d} \alpha])^{1/4}. \end{aligned}$$

Hence

$$I \leq C_2 e^{(t_1 + \cdots + t_d)/2} Hgt([r_1^{-t_1} \cdots r_d^{-t_d} \alpha])^{1/4},$$

- 4 where the sum corresponds to the first term ($k = 0$) in the statement.

Step 2. To estimate (II),

$$\begin{aligned} \left\| \mathcal{R}^{-s}[r^{-u} \alpha, (\mathcal{P}_{U_d(t)}^{d,\alpha} m)] \right\|_{-(s-2)} &= \left\| e^{ud} \mathcal{R}^{-s}[r^{-u} \alpha, (\mathcal{P}_{U_d(t-u)}^{d, r^{-u} \alpha} m)] \right\|_{-(s-2)} \\ (24) \quad &\leq C_3(s, s') e^{ud} \left\| [r^{-u} \alpha, \partial(\mathcal{P}_{U_d(t-u)}^{d, r^{-u} \alpha} m)] \right\|_{-s'}. \end{aligned}$$

The boundary $\partial(\mathcal{P}_{U_d}^{d, r^{-t} \alpha})$ is the sum of $2d$ currents of dimension $d - 1$. These currents are Birkhoff sums of d face subgroups obtained from $\mathcal{P}_{U_d}^{d, r^{-t} \alpha}$ by omitting one of the base vector fields X_i . It is reduced to $(d - 1)$ dimensional shape obtained from $U_d(t - u) := [0, e^{t_1 - u}] \times \cdots \times [0, e^{t_d - u}]$. For each $1 \leq j \leq d$, there are Birkhoff sums along $d - 1$ dimensional cubes. By induction hypothesis, we add all the $d - 1$

dimensional cubes by adding all the terms along j :

(25)

$$\begin{aligned} & \left\| [r^{-u} \alpha, (\mathcal{P}_{U_{d-1}(t-u)}^{d-1, r^{-u} \alpha} m)] \right\|_{-s'} \leq C_4(s', d-1) \sum_{j=1}^d \sum_{k=0}^{d-1} \sum_{\substack{1 \leq i_1 < \dots < i_k \leq d \\ i_l \neq j, \forall l}} \int_0^{t_{i_k}-u} \dots \int_0^{t_{i_1}-u} \\ & \exp\left(\frac{1}{2} \sum_{\substack{l=1 \\ l \neq j}}^d (t_l - u) - \frac{1}{2} \sum_{l=1}^k u_{i_l}\right) Hgt\left(\left[\prod_{\substack{1 \leq l \leq d \\ l \neq j}} r_l^{-(t_l-u)} \prod_{l=1}^k r_{i_l}^{u_{i_l}} (r^{-u} \alpha)\right]\right)^{1/4} du_{i_1} \dots du_{i_k}. \end{aligned}$$

1 Combining (23) and (24), we obtain the estimate for (II).

$$\begin{aligned} (II) & \leq C_5(s', d-1) \sum_{j=1}^d \sum_{k=0}^{d-1} \sum_{\substack{1 \leq i_1 < \dots < i_k \leq d \\ i_l \neq j}} \int_0^{t_1+\dots+t_d} \int_0^{t_{i_k}-u} \dots \int_0^{t_{i_1}-u} du_{i_1} \dots du_{i_k} du \\ & \times \exp\left(\frac{1}{2} \sum_{\substack{l=1 \\ l \neq j}}^d t_l - \frac{1}{2} u - \frac{1}{2} \sum_{l=1}^k u_{i_l}\right) Hgt\left(\left[\prod_{1 \leq l \leq d} r_l^{-t_l} \prod_{l=1}^k r_{i_l}^{u_{i_l}} (r_j^{-u+t_j} \alpha)\right]\right)^{1/4}. \end{aligned}$$

Applying the change of variable $u_j = t_j - u$, we obtain

$$\begin{aligned} (II) & \leq C_6(s', d-1) \sum_{j=1}^d \sum_{k=1}^{d-1} \sum_{\substack{1 \leq i_1 < \dots < i_k \leq d \\ i_l \neq j}} \\ & \times \left(\int_{-(t_1+\dots+t_d)+t_j}^{t_j} \int_0^{t_{i_k}-u} \dots \int_0^{t_{i_1}-u} du_{i_1} \dots du_{i_k} du_j \right. \\ & \times \exp\left(\frac{1}{2} (t_1 + \dots + t_d) - \frac{1}{2} u_j - \frac{1}{2} \sum_{l=1}^k u_{i_l}\right) Hgt\left(\left[\prod_{1 \leq l \leq d} r_l^{-t_l} \prod_{l=1}^k r_{i_l}^{u_{i_l}} (r_j^{-u_j} \alpha)\right]\right)^{1/4} \Big). \end{aligned}$$

Simplifying multi-summation above, (with $-(t_1 + \dots + t_d) + t_j \leq 0$)

$$\begin{aligned} (II) & \leq C_7(s', d) \sum_{k=1}^d \sum_{1 \leq i_1 < \dots < i_k \leq d} \int_0^{t_{i_k}} \dots \int_0^{t_{i_1}} du_{i_1} \dots du_{i_k} \\ & \times \exp\left(\frac{1}{2} (t_1 + \dots + t_d) - \frac{1}{2} \sum_{l=1}^k u_{i_l}\right) Hgt\left(\left[\prod_{1 \leq l \leq d} r_l^{-t_l} \prod_{l=1}^k r_{i_l}^{u_{i_l}} \alpha\right]\right)^{1/4}. \end{aligned}$$

2 *Step 3. (Remainder estimate).* The remainder term is obtained from Lemma 3.2
 3 (Stokes' theorem). Following step 2, estimate of remainder reduces to that of $d-1$
 4 form. Combining with the step 1, we have the following
 (26)

$$\left\| \mathcal{R}^{-s}[\alpha, \partial(\mathcal{P}_{U_d}^{d, \alpha} m)] \right\|_{-s} \leq C(s) \sum_{i=1}^{d-1} \left\| \mathcal{I}^{-s}[\alpha, (\mathcal{P}_{U_i}^{i, \alpha} m)] \right\|_{-s} + \left\| \mathcal{R}^{-s}[\alpha, (\mathcal{P}_{U_1}^{1, \alpha} m)] \right\|_{-s}$$

where U_i is i -dimensional rectangle. Sum of the boundary terms are absorbed in the bound of (I) + (II). For 1-dimensional remainder with interval Γ_T , the boundary is a 0-dimensional current. Then,

$$\langle \partial(\mathcal{P}_{[0,T]}^{1, \alpha} m), f \rangle = f(P_T^{1, \alpha} m) - f(m).$$

Hence, by Sobolev embedding theorem and by definition of Sobolev constant (9) and (14),

$$\left\| \mathcal{R}^{-s}[\alpha, \partial(\mathcal{P}_{[0,T]}^{1,\alpha} m)] \right\|_{-s} \leq 2B_s([\alpha]) \leq C(s)Hgt([\alpha])^{1/4}.$$

Then, by inequality (15)

$$\begin{aligned} C(s)Hgt([\alpha])^{1/4} &= C(s)Hgt([r_{\mathbf{t}} r_{-\mathbf{t}} \alpha])^{1/4} \\ &\leq C(s)e^{(t_1 + \dots + t_d)/2} Hgt([r_1^{-t_1} \dots r_d^{-t_d} \alpha])^{1/4}. \end{aligned}$$

- 1 This implies that remainder term produces one more term like the bound of (I).
 2 Therefore, the Theorem follows from combining all the terms (I), (II), and remain-
 3 der. \square

4 **Remark.** It is also reasonable to decompose the rectangle $U_d(t)$ as a union of
 5 several squares and renormalize their faces. However, it may also involve compu-
 6 tational difficulty and such approach of summing up squares may provide a rough
 7 upper bound of ergodic integrals. To obtain a necessary bound for the Lemma 3.4,
 8 we rather simply generalized the methods of Theorem 5.10 in [CF15].

9 Let us set

$$(27) \quad \mathcal{K}_{\alpha, \mathbf{t}, s}(\Gamma) := \left\| \mathcal{R}^{-s}[r_{-\mathbf{t}}(\alpha), (\mathcal{P}_{U_\Gamma}^{d, r_{-\mathbf{t}}(\alpha)} m)] \right\|_{-(s+1)}.$$

- 10 Now we prove the remainder estimate that will be used in (34).

Lemma 3.4. Let $s > s_{d,g}$. There exists a constant $C(s, \Gamma) > 0$ such that for any rectangle $U_\Gamma = [0, e^{\Gamma_1}] \times \dots \times [0, e^{\Gamma_d}]$,

$$\mathcal{K}_{\alpha, \mathbf{t}, s}(\Gamma) \leq C(s, \Gamma)Hgt([r_{-\mathbf{t}}(\alpha)])^{1/4}.$$

Proof. By Lemma 3.2, we begin to find the bound of $d-1$ (renormalized) currents. By Theorem 3.3, we obtain the remainder estimate

$$\begin{aligned} \mathcal{K}_{\alpha, \mathbf{t}}(\Gamma) &\leq C \sum_{k=0}^{d-1} \sum_{1 \leq i_1 < \dots < i_k \leq d-1} \int_0^{\Gamma_{i_k}} \dots \int_0^{\Gamma_{i_1}} \exp\left(\frac{1}{2} \sum_{l=1}^{d-1} \Gamma_l - \frac{1}{2} \sum_{l=1}^k u_{i_l}\right) \\ &\quad \times Hgt\left(\left[\prod_{1 \leq j \leq d-1} r_j^{-\Gamma_j} \prod_{l=1}^k r_{i_l}^{u_{i_l}}(r_{-\mathbf{t}}(\alpha))\right]\right)^{1/4} du_{i_1} \dots du_{i_k}. \end{aligned}$$

It follows from (15) that for $0 \leq k \leq d-1$,

$$Hgt\left(\left[\prod_{1 \leq j \leq d-1} r_j^{-\Gamma_j} \prod_{l=1}^k r_{i_l}^{u_{i_l}}(r_{-\mathbf{t}}(\alpha))\right]\right)^{1/4} \leq e^{\frac{1}{2}(\sum_{l=1}^k u_{i_l} - \sum_{l=1}^{d-1} \Gamma_l)} Hgt([r_{-\mathbf{t}}(\alpha)])^{1/4}.$$

Then, we obtain

$$\begin{aligned} (28) \quad \left\| \mathcal{R}^{-s}[r_{-\mathbf{t}}(\alpha), (\mathcal{P}_{U_\Gamma}^{d, r_{-\mathbf{t}}(\alpha)} m)] \right\|_{-s} &\leq C(s) \left(\sum_{k=0}^{d-1} \sum_{1 \leq i_1 < \dots < i_k \leq d-1} \prod_{l=1}^k \Gamma_{i_l} \right) Hgt([r_{-\mathbf{t}}(\alpha)])^{1/4} \\ &\leq C(s, \Gamma)Hgt([r_{-\mathbf{t}}(\alpha)])^{1/4}. \end{aligned}$$

- 11 Therefore, we obtain the conclusion. \square

3.2. Constructions of the functionals. For fixed $\alpha \in \text{Aut}_0(H^g)$, let $\Pi_H^{-s} : A_d(\mathfrak{p}, W_\alpha^{-s}(M)) \rightarrow A_d(\mathfrak{p}, W_\alpha^{-s}(H))$ denote the orthogonal projection on a single irreducible unitary representation H . We further decompose projection operators with basic current $B_\alpha^{-s,H}$ and its remainder $R_\alpha^{-s,H}$ given by

$$\Pi_H^{-s} = \mathcal{B}_\alpha^{-s}(\Gamma) B_\alpha^{-s,H} + R_\alpha^{-s,H}$$

1 where $\mathcal{B}_{H,\alpha}^{-s} : A_d(\mathfrak{p}, W_\alpha^{-s}(M)) \rightarrow \mathbb{C}$ denote the orthogonal component map in the
2 direction of basic current, supported on a single irreducible unitary representation.

3 The *Bufetov functionals* on a standard rectangle Γ are defined for all $[\alpha] \in DC$
4 as follows.

Lemma 3.5. *Let $[\alpha] \in DC(L)$. For $s > s_{d,g}$, the limit*

$$\hat{\beta}_H(\alpha, \Gamma) = \lim_{t_1, \dots, t_d \rightarrow \infty} e^{-(t_1 + \dots + t_d)/2} \mathcal{B}_{H,r_t(\alpha)}^{-s}(\Gamma)$$

5 *exists and it defines a finitely-additive measure on the set of standard rectangles.*
6 *Moreover, there exists a constant $C(s, \Gamma) > 0$ such that the following estimate holds:*

$$(29) \quad \left\| \Pi_{H,\alpha}^{-s}(\Gamma) - \hat{\beta}_H(\alpha, \Gamma) B_\alpha^H \right\|_{\alpha, -s} \leq C(s, \Gamma)(1 + L).$$

Proof. For simplicity, we omit dependence of H . For every $\mathbf{t} \in \mathbb{R}^d$, we have the following orthogonal splitting:

$$\Pi_{H,\alpha}^{-s}(\Gamma) = \mathcal{B}_{\alpha,\mathbf{t}}^{-s}(\Gamma) B_{\alpha,\mathbf{t}} + R_{\alpha,\mathbf{t}},$$

where

$$\mathcal{B}_{\alpha,\mathbf{t}}^{-s} := \mathcal{B}_{H,r_{-\mathbf{t}}(\alpha)}^{-s}, \quad B_{\alpha,\mathbf{t}} := B_{r_{-\mathbf{t}}(\alpha)}^{-s,H}, \quad R_{\alpha,\mathbf{t}} := R_{r_{-\mathbf{t}}(\alpha)}^{-s,H}.$$

For any $\mathbf{h} \in \mathbb{R}^d$, we have

$$\mathcal{B}_{\alpha,\mathbf{t}+\mathbf{h}}^{-s}(\Gamma) B_{\alpha,\mathbf{t}+\mathbf{h}} + R_{\alpha,\mathbf{t}+\mathbf{h}} = \mathcal{B}_{\alpha,\mathbf{t}}^{-s}(\Gamma) B_{\alpha,\mathbf{t}} + R_{\alpha,\mathbf{t}}.$$

7 By reparametrization (12), we have $B_{\mathbf{t}+\mathbf{h}} = e^{-(h_1 + \dots + h_d)/2} B_{\mathbf{t}}$ and

$$(30) \quad \mathcal{B}_{\alpha,\mathbf{t}+\mathbf{h}}^{-s}(\Gamma) = e^{(h_1 + \dots + h_d)/2} \mathcal{B}_{\alpha,\mathbf{t}}^{-s}(\Gamma) + \mathcal{B}_{\alpha,\mathbf{t}+\mathbf{h}}^{-s}(R_{\alpha,\mathbf{t}})$$

and it follows that

$$\mathcal{B}_{\alpha,\mathbf{t}+\mathbf{h}}^{-s}(\Gamma) = e^{h_1/2} \mathcal{B}_{\alpha,t_1,t_2+h_2,\dots,t_d+h_d}^{-s}(\Gamma) + \mathcal{B}_{\alpha,\mathbf{t}+\mathbf{h}}^{-s}(R_{\alpha,\mathbf{t}}).$$

8 By differentiating at $h_1 = 0$,

$$(31) \quad \frac{d}{dt_1} \mathcal{B}_{\alpha,t_1,t_2+h_2,\dots,t_d+h_d}^{-s}(\Gamma) = \frac{1}{2} \mathcal{B}_{\alpha,t_1,t_2+h_2,\dots,t_d+h_d}^{-s}(\Gamma) + \left[\frac{d}{dh_1} \mathcal{B}_{\alpha,\mathbf{t}+\mathbf{h}}^{-s}(R_{\alpha,\mathbf{t}}) \right]_{h_1=0}.$$

Therefore, we solve the following first order ODE

$$\frac{d}{dt_1} \mathcal{B}_{\alpha,t_1,t_2+h_2,\dots,t_d+h_d}^{-s}(\Gamma) = \frac{1}{2} \mathcal{B}_{\alpha,t_1,t_2+h_2,\dots,t_d+h_d}^{-s}(\Gamma) + \mathcal{K}_{\alpha,\mathbf{t}}^{(1)}(\Gamma)$$

where

$$\mathcal{K}_{\alpha,\mathbf{t},s}^{(1)}(\Gamma) := \left[\frac{d}{dh_1} \mathcal{B}_{\alpha,\mathbf{t}+\mathbf{h}}^{-s}(R_{\alpha,\mathbf{t}}) \right]_{h_1=0}.$$

Then, the solution of the differential equation is

$$\begin{aligned}\mathcal{B}_{\alpha,t_1,t_2+h_2,\dots,t_d+h_d}^{-s}(\Gamma) &= e^{t_1/2} \left(\mathcal{B}_{\alpha,0,t_2+h_2,\dots,t_d+h_d}^{-s}(\Gamma) + \int_0^{t_1} e^{-\tau_1/2} \mathcal{K}_{\alpha,\tau,s}^{(1)}(\Gamma) d\tau_1 \right) \\ &= e^{t_1/2} \mathcal{B}_{\alpha,0,t_2+h_2,\dots,t_d+h_d}^{-s}(\Gamma) + \int_0^{t_1} e^{(t_1-\tau_1)/2} \mathcal{K}_{\alpha,\tau,s}^{(1)}(\Gamma) d\tau_1.\end{aligned}$$

Note by reparametrization

$$e^{t_1/2} \mathcal{B}_{\alpha,0,t_2+h_2,\dots,t_d+h_d}^{-s}(\Gamma) = e^{h_2/2} \mathcal{B}_{\alpha,t_1,t_2,t_3+h_3,\dots,t_d+h_d}^{-s}(\Gamma)$$

- 1 and it is possible to differentiate the previous equation with respect to h_2 again.
2 Then

$$(32) \quad \frac{d}{dt_2} \mathcal{B}_{\alpha,t_1,t_2,\dots,t_d+h_d}^{-s}(\Gamma) = \frac{1}{2} \mathcal{B}_{\alpha,t_1,t_2,t_3+h_3,\dots,t_d+h_d}^{-s} + \int_0^{t_1} e^{-\tau_1/2} \mathcal{K}_{\alpha,\tau,s}^{(2)}(\Gamma) d\tau_1$$

- 3 where $\mathcal{K}_{\alpha,\tau,s}^{(2)}(\Gamma) = \frac{d}{dh_2} \mathcal{K}_{\alpha,\tau,s}^{(1)}(\Gamma)$.

Then, the solution of equation (32) is

$$\begin{aligned}\mathcal{B}_{\alpha,t_1,t_2,\dots,t_d+h_d}^{-s}(\Gamma) &= e^{t_2/2} \left(\mathcal{B}_{\alpha,t_1,0,t_3+h_3,\dots,t_d+h_d}^{-s} + \int_0^{t_2} e^{-\tau_2/2} \int_0^{t_1} e^{(t_1-\tau_1)/2} \mathcal{K}_{\alpha,\tau,s}^{(2)}(\Gamma) d\tau_1 d\tau_2 \right) \\ &= e^{h_3/2} \mathcal{B}_{\alpha,t_1,t_2,t_3,\dots,t_d+h_d}^{-s} + \int_0^{t_2} e^{(t_2-\tau_2)/2} \int_0^{t_1} e^{(t_1-\tau_1)/2} \mathcal{K}_{\alpha,\tau,s}^{(2)}(\Gamma) d\tau_1 d\tau_2.\end{aligned}$$

- 4 Inductively, we solve the first order ODE repeatedly and obtain the following
5 solution

$$(33) \quad \mathcal{B}_{\alpha,t}^{-s}(\Gamma) = e^{(t_1+\dots+t_d)/2} \left(\mathcal{B}_{\alpha,0}^{-s} + \int_0^{t_d} \dots \int_0^{t_1} e^{-(\tau_1+\dots+\tau_d)/2} \mathcal{K}_{\alpha,\tau,s}^{(d)}(\Gamma) d\tau_1 \dots d\tau_d \right)$$

where

$$\mathcal{K}_{\alpha,t,s}^{(d)}(\Gamma) = \left[\frac{d}{dh_d} \dots \frac{d}{dh_1} \mathcal{B}_{\alpha,t+h}^{-s}(R_{\alpha,t}) \right]_{h_d \dots h_1=0}.$$

Let $\langle \cdot, \cdot \rangle_{\alpha,t}$ denote the inner product in the space of Hilbert current $A_d(\mathfrak{p}, W_{r-t(\alpha)}^{-s}(H))$.
By the intertwining formula (11),

$$\begin{aligned}\mathcal{B}_{\alpha,t+h}^{-s}(R_{\alpha,t}) &= \langle R_{\alpha,t}, \frac{B_{\alpha,t+h}}{|B_{\alpha,t+h}|_{t+h}^2} \rangle_{\alpha,t+h} \\ &= \langle R_{\alpha,t} \circ U_{-h}, \frac{B_{\alpha,t+h} \circ U_{-h}}{|B_{\alpha,t+h}|_{t+h}^2} \rangle_{\alpha,t} \\ &= \langle R_{\alpha,t} \circ U_{-h}, \frac{B_{\alpha,t}}{|B_{\alpha,t}|_t^2} \rangle_{\alpha,t} = \mathcal{B}_{\alpha,t}^{-s}(R_{\alpha,t} \circ U_{-h}).\end{aligned}$$

In the sense of distributions,

$$\begin{aligned}\frac{d}{dh_d} \dots \frac{d}{dh_1} (R_{\alpha,t} \circ U_{-h}) &= -R_{\alpha,t} \circ \left(\frac{d}{2} + \sum_{i=1}^d X_i(t) \right) \circ U_{-h} \\ &= \left[\left(\sum_{i=1}^d X_i(t) - \frac{d}{2} \right) R_{\alpha,t} \right] \circ U_{-h}.\end{aligned}$$

Then we compute

$$\left[\frac{d}{dh_d} \cdots \frac{d}{dh_1} (\mathcal{B}_{\alpha, \mathbf{t} + \mathbf{h}}^{-s}(R_{\alpha, \mathbf{t}})) \right]_{\mathbf{h}=0} = -\mathcal{B}_{\alpha, \mathbf{t}}^{-s} \left(\left(\sum_{i=1}^d X_i(t) - \frac{d}{2} \right) R_{\alpha, \mathbf{t}} \right).$$

- 1 Recall by (27) that $\mathcal{K}_{\alpha, \mathbf{t}, s}(\Gamma) = \|\mathcal{R}_{\alpha, \mathbf{t}}^{-s}\|_{r_{-\mathbf{t}}(\alpha), -(s+1)}$. By Proposition 2.8 and
 2 Lemma 3.4, for any $\epsilon > 0$

$$(34) \quad |\mathcal{B}_{\alpha, \mathbf{t}}^{-s} \left(\left(\sum_{i=1}^d X_i(t) - \frac{d}{2} \right) R_{\alpha, \mathbf{t}} \right)| \leq \mathcal{K}_{\alpha, \mathbf{t}, s}(\Gamma) \leq C(s, \Gamma) \text{Hgt}([r_{-\mathbf{t}}(\alpha)])^{1/4}.$$

Therefore, the solution of equation (33) exists under Diophantine condition (20) and the following holds:

$$\lim_{t_1, \dots, t_d \rightarrow \infty} e^{-(t_1 + \dots + t_d)/2} \mathcal{B}_{\alpha, \mathbf{t}}^{-s}(\Gamma) = \hat{\beta}_H(\alpha, \Gamma).$$

Moreover, the complex number

$$\hat{\beta}_H(\alpha, \Gamma) = \mathcal{B}_{\alpha, 0}^{-s} + \int_0^\infty \cdots \int_0^\infty e^{-(\tau_1 + \dots + \tau_d)/2} \mathcal{K}_{\alpha, \tau, s}(\Gamma) d\tau_1 \cdots d\tau_d$$

depends continuously on $\alpha \in DC(L)$. Since we have

$$\Pi_{H, \alpha}^{-s}(\Gamma) - \hat{\beta}(\alpha, \Gamma) B_\alpha^H = R_0 - \left(\int_0^\infty \cdots \int_0^\infty e^{-(\tau_1 + \dots + \tau_d)/2} \mathcal{K}_{\alpha, \tau, s}(\Gamma) d\tau_1 \cdots d\tau_d \right) B_\alpha^H,$$

by Diophantine condition again,

$$\left\| \Pi_{H, \alpha}^{-s}(\Gamma) - \hat{\beta}_H(\alpha, \Gamma) B_\alpha^H \right\|_{\alpha, -s} \leq C(s, \Gamma)(1 + L).$$

3

□

- 4 **3.3. Proof of Theorem 1.3.** The proof of Theorem 1.3 follows immediately from
 5 the refinement to the constructions of Bufetov functionals (see also [BF14, §2.5] for
 6 horocycle flows).

- 7 *Notation.* The action of flow $\{r_t\}_{t \in \mathbb{R}}$ on a current \mathcal{C} is defined by pull-back as
 8 follows:

$$(r_t^* \mathcal{C})(\omega) = \mathcal{C}(r_{-t}^* \omega), \quad \text{for any smooth form } \omega.$$

Lemma 3.6 (Invariance). *Let $1 \leq d \leq g$. The functional $\hat{\beta}_H$ defined on d -standard rectangle $\Gamma = \Gamma_T^X$ is invariant under the action of $(Q_y^{j, Y})$ for any $y \in \mathbb{R}_+^j$ and $1 \leq j \leq d$. That is,*

$$\hat{\beta}_H(\alpha, (Q_y^{j, Y})_* \Gamma) = \hat{\beta}_H(\alpha, \Gamma).$$

- 9 *Proof.* We will prove the functional $\hat{\beta}_H$ exists with respect to rectangle $(Q_y^{j, Y})_* \Gamma$
 10 and prove its invariance under the action $Q_y^{j, Y}$. It suffices to verify the invariance
 11 property under the rank 1 action $Q_\tau^{1, Y}$ for $\tau \in \mathbb{R}$ since we can apply the statement
 12 for d -rank actions repeatedly.

- 13 Given a standard d -dimensional rectangle Γ , set $\Gamma_Q := (Q_\tau^{1, Y})_* \Gamma$. Let $D(\Gamma, \Gamma_Q)$
 14 be the $(d+1)$ dimensional space spanned by trajectories of the action of $Q_\tau^{1, Y}$
 15 projecting Γ onto Γ_Q . Then $D(\Gamma, \Gamma_Q)$ is a union of all orbits I of action $Q_\tau^{1, Y}$ such
 16 that the boundary of I , d -dimensional faces, is contained in $\Gamma \cup \Gamma_Q$, and interior of I
 17 is disjoint from $\Gamma \cup \Gamma_Q$. $D(\Gamma, \Gamma_Q)$ is defined by integration and it is a $(d+1)$ -current.




FIGURE 2. Illustration of the standard d -rectangles Γ , Γ_Q , $d+1$ dimensional current $D(\Gamma, \Gamma_Q)$ and supports of $r_{-t}(\Gamma)$ and $r_{-t}(\Gamma_Q)$.

For convenience, let us denote renormalization flow by $r_t := r_i^t$ for some i -th coordinates. Then, $r_{-t}(\Gamma)$ and $r_{-t}(\Gamma_Q)$ are respectively the support of the currents $r_t^*\Gamma$ and $r_t^*\Gamma_Q$. Thus, we have the following identity

$$r_t^*D(\Gamma, \Gamma_Q) = D(r_{-t}(\Gamma), r_{-t}(\Gamma_Q)).$$

- 1 Since the current $\partial D(\Gamma, \Gamma_Q) - (\Gamma - \Gamma_Q)$ is composed of orbits for the action $Q_\tau^{1,Y}$,
 2 it follows that

$$(35) \quad \partial[r_t^*D(\Gamma, \Gamma_Q)] - (r_t^*\Gamma - r_t^*\Gamma_Q) = r_t^*[\partial D(\Gamma, \Gamma_Q) - (\Gamma - \Gamma_Q)] \rightarrow 0.$$

Now, we turn to prove the volume of $D(r_{-t}(\Gamma), r_{-t}(\Gamma_Q))$ is uniformly bounded for all $t > 0$. For any $p \in \Gamma$, set $\tau(p)$ be length of the arc lying in $D_Q := D(\Gamma, \Gamma_Q)$, and set $\tau_\Gamma := \sup\{\tau(p) \mid p \in \Gamma\} < \infty$. We write

$$vol_{d+1}(D_Q) = \int_\Gamma \tau dvol_d.$$

- 3 Since $vol_d(r_{-t}(\Gamma)) \leq e^t vol_d(\Gamma)$,

$$(36) \quad vol_{d+1}(r_{-t}(D_Q)) = \int_{r_{-t}(\Gamma)} \tau dvol_d \leq \tau_\Gamma e^{-t} vol_d(r_{-t}(\Gamma)) \leq \tau_\Gamma vol_d(\Gamma) < \infty.$$

- 4 Note that d -dimensional current $(Q_\tau^{1,Y})_*\Gamma - \Gamma$ is equal to the boundary of the
 5 $(d+1)$ dimensional current D_Q . By arguments in remainder estimate (or Sobolev
 6 embedding theorem),

$$(37) \quad \|(Q_\tau^{1,Y})_*\Gamma - \Gamma\|_{r_{-t}(\alpha), -s} \leq C_s \tau B_s([r_{-t}\alpha]) \leq C_s \tau \text{Hgt}([r_{-t}\alpha])^{1/4}$$

- 7 is finite for all $t > 0$.

Then, by (35), (36) and existence of Bufetov functional $\hat{\beta}_H(\alpha, \Gamma)$, the last inequality holds:

$$\|\mathcal{B}_{\alpha, t}^{-s}((Q_\tau^{1,Y})_*\Gamma) - \mathcal{B}_{\alpha, t}^{-s}(\Gamma)\|_{\alpha, -s} < \infty.$$

- 8 Therefore, by the definition of Bufetov functional in the Lemma 3.5, $\hat{\beta}_H(\alpha, (Q_\tau^{1,Y})_*\Gamma)$
 9 exists and $\hat{\beta}_H(\alpha, \Gamma)$ is invariant under the action of $Q_\tau^{1,Y}$. \square

- 10 *Proof of Theorem 1.3. Additive property.* It follows from the linearity of projections
 11 and limit.

- 12 *Scaling property.* It is immediate from the definition.

Bounded property. By the scaling property, for $r_t = r_1^t \cdots r_d^t$ with $t > 0$,

$$\hat{\beta}_H(\alpha, \Gamma) = e^{dt/2} \hat{\beta}_H(r_t(\alpha), \Gamma).$$

Choose $t = \log(\int_{\Gamma} |\hat{X}|)$ and $\hat{X} = \hat{X}_1 \wedge \cdots \wedge \hat{X}_d$, then Bufetov functional on the rectangle Γ is bounded:

$$|\hat{\beta}_H(\alpha, \Gamma)| \leq C(\Gamma) \left(\int_{\Gamma} |\hat{X}| \right)^{d/2}.$$

1 *Invariance property.* Now it follows directly from the Lemma 3.6. \square

2 *Notation.* We write for $\mathbf{T} = (T^{(i)}) \in \mathbb{R}^d$ and $\mathbf{t} = (t_1, \dots, t_d) \in \mathbb{R}^d$,

$$(38) \quad \text{vol}(U(\mathbf{T})) := \prod_{i=1}^d T^{(i)}, \quad \text{vol}(U(\mathbf{t})) = \prod_{i=1}^d t_i.$$

3 Now we extend the properties of functional $\hat{\beta}_H$ to cocycle β_H .

4 *Proof of Corollary 1.5.* Cocycle property of β_H follows from the additive property
5 of $\hat{\beta}_H$. Scaling and bounded properties are immediate. Denote $\mathbf{tT} := (t_1 T^{(1)}, \dots, t_d T^{(d)}) \in$
6 \mathbb{R}_+^d , and we obtain

$$(39) \quad \beta_H(\alpha, m, \mathbf{tT}) = \text{vol}(U(\mathbf{T}))^{1/2} \beta_H(r_{\log \mathbf{T}}(\alpha), m, \mathbf{t}).$$

By Lemma 3.5 and scaling property, we have

$$\begin{aligned} D_{\alpha}^H(f) \beta_H(\alpha, m, \mathbf{t}) &= \lim_{|\mathbf{U}(\mathbf{T})| \rightarrow \infty} \frac{1}{\text{vol}(U(\mathbf{T}))^{1/2}} \beta_H(r_{-\log \mathbf{t}}(\alpha), m, \mathbf{tT}) \\ &= \lim_{|\mathbf{U}(\mathbf{T})| \rightarrow \infty} \frac{1}{\text{vol}(U(\mathbf{T}))^{1/2}} \left\langle \mathcal{P}_{U(\mathbf{tT})}^{d, r_{-\log \mathbf{t}}(\alpha)} m, \omega_{f_H} \right\rangle. \end{aligned}$$

7 It follows that $\beta_H(\alpha, \cdot, \mathbf{t}) \in H$ as a point-wise limit of Birkhoff integrals. This
8 implies orthogonal property. \square

9 For $\gamma \in Sp_{2g}(\mathbb{Z})$, we have

$$\beta_H(\gamma\alpha, \gamma(m), \mathbf{T}) = \beta_H(\alpha, m, \mathbf{T}).$$

10 It means that the function $\beta_H(\cdot, m, \mathbf{T})$ is well-defined on the moduli space \mathfrak{M}_g .

3.4. Proof of Theorem 1.6. We define the excursion function

$$\begin{aligned} E_{\mathfrak{M}}(\alpha, \mathbf{T}) &:= \int_0^{\log T^{(d)}} \cdots \int_0^{\log T^{(1)}} e^{-(t_1 + \cdots + t_d)/2} \text{Hgt}([r_{\mathbf{t} - \log \mathbf{T}}(\alpha)])^{1/4} dt_1 \cdots dt_d \\ &= \prod_{i=1}^d (T^{(i)})^{1/2} \int_0^{\log T^{(d)}} \cdots \int_0^{\log T^{(1)}} e^{(t_1 + \cdots + t_d)/2} \text{Hgt}([r_{\mathbf{t}}(\alpha)])^{1/4} dt_1 \cdots dt_d. \end{aligned}$$

Lemma 3.7 (Bound of cocycle). *For any Diophantine $[\alpha] \in DC(L)$ and for any $f \in W^s(M)$ for $s > s_{d,g} + 1/2$, the Bufetov cocycle β^f is defined by a uniformly convergent series:*

$$|\beta^f(\alpha, m, \mathbf{tT})| \leq C_s \left(L + \text{vol}(U(\mathbf{T}))^{1/2} (1 + \text{vol}(U(\mathbf{t})) + E_{\mathfrak{M}}(\alpha, \mathbf{T})) \right) \|f\|_{\alpha, s}.$$

Proof. Recall that by (7),

$$\beta^f(\alpha, m, \mathbf{T}) = \sum_H D_{\alpha}^H(f) \beta_H(\alpha, m, \mathbf{T}).$$

11 It follows from Lemma 3.5 that there exists a constant $C > 0$ such that whenever
12 $[\alpha] \in DC(L)$, then

$$(40) \quad |\beta_H(\alpha, m, \mathbf{t})| \leq C(1 + L + \text{vol}(U(\mathbf{t}))), \quad (m, \mathbf{t}) \in M \times \mathbb{R}_+^d.$$

By Diophantine condition (20), whenever $[\alpha] \in DC(L)$ then $[r_{\log \mathbf{T}}(\alpha)] \in DC(L_{\mathbf{T}})$ with

$$L_{\mathbf{T}} \leq L \text{vol}(U(\mathbf{T}))^{-1/2} + E_{\mathfrak{M}}(\alpha, \mathbf{T}).$$

Thus by (40), we obtain for all $(m, \mathbf{t}) \in M \times \mathbb{R}^d$,

$$|\beta_H(r_{\log \mathbf{T}}(\alpha), m, \mathbf{t})| \leq C(1 + L_{\mathbf{T}} + \text{vol}(U(\mathbf{t}))).$$

By the scaling property (39), it follows that for all $s > s_{d,g}$ and $q > 1/2$

$$\begin{aligned} |\beta^f(\alpha, m, \mathbf{t}\mathbf{T})| &\leq C_s \text{vol}(U(\mathbf{T}))^{1/2} (1 + L + \text{vol}(U(\mathbf{t}))) \sum_{n \in \mathbb{Z}} \|f_n\|_{\alpha, s} \\ &\leq C_s \text{vol}(U(\mathbf{T}))^{1/2} (1 + L_{\mathbf{T}} + \text{vol}(U(\mathbf{t}))) \left(\sum_{n \in \mathbb{Z}} (1 + n^2)^q \right)^{-1/2} \\ &\quad \times \left(\sum_{n \in \mathbb{Z}} \left\| (1 - Z^2)^{q/2} f_n \right\|_{\alpha, s}^2 \right)^{1/2}. \end{aligned}$$

Therefore, for all $s' = s_{d,g} + q$, there exists a constant $C_{s'} > 0$ such that

$$|\beta^f(\alpha, m, \mathbf{t}\mathbf{T})| \leq C_{s'} \text{vol}(U(\mathbf{T}))^{1/2} (1 + L_{\mathbf{T}} + \text{vol}(U(\mathbf{t}))) \|f\|_{\alpha, s'}.$$

1

□

2 By Lemma 3.5, 3.7 and identification of the norm for the form ω_f , asymptotic
3 formula on each irreducible component provides the following corollary.

4 **Corollary 3.8.** *For all $s > s_{d,g} + 1/2$, there exists a constant $C_s > 0$ such that for*
5 *all $[\alpha] \in DC(L)$, for all $f \in W_{\alpha}^s(M)$ and for all $(m, \mathbf{T}) \in M \times \mathbb{R}^d$, we have*

$$(41) \quad \left| \left\langle \mathcal{P}_{U(\mathbf{T})}^{d, \alpha} m, \omega_f \right\rangle - \beta^f(\alpha, m, \mathbf{T}) \right| \leq C_s (1 + L) \|\omega_f\|_{\alpha, s}$$

6 for $U(\mathbf{T}) = [0, T^{(1)}] \times \cdots \times [0, T^{(d)}]$ and $\omega_f = f \omega^{d, \alpha} \in \Lambda^d \mathfrak{p} \otimes W_{\alpha}^s(M)$.

7 *Proof of Theorem 1.6.* The theorem follows from Corollary 3.8 for $\alpha \in \bigcup_{L>0} DC(L)$.

8

□

9

4. LIMIT DISTRIBUTIONS

10 In this section, we prove Theorem 1.7, limit distribution of normalized ergodic
11 integrals of higher rank actions on the standard rectangles.

12 4.1. Proof of Theorem 1.7.

13 **Lemma 4.1.** *There exists a continuous modular function $\theta_H : \text{Aut}_0(\mathbf{H}^g) \rightarrow H \subset$*
14 *$L^2(M)$ such that for any $\omega_f = f \omega^{d, \alpha} \in \Lambda^d \mathfrak{p} \otimes W_{\alpha}^s(H)$ with $s > d/2$,*

$$(42) \quad \lim_{|U(\mathbf{T})| \rightarrow \infty} \left\| \frac{1}{\text{vol}(U(\mathbf{T}))^{1/2}} \left\langle \mathcal{P}_{U(\mathbf{T})}^{d, \alpha}(\cdot), \omega_f \right\rangle - \theta_H(r_{\log \mathbf{T}}(\alpha)) D_{\alpha}^H(f) \right\|_{L^2(M)} = 0.$$

15 *The family $\{\theta_H(\alpha) \mid \alpha \in \text{Aut}_0(\mathbf{H}^g)\}$ has a constant norm in $L^2(M)$.*

Proof. By the Fourier transform, the space of smooth vectors and Sobolev space $W^s(H)$ is represented as the Schwartz space $\mathcal{S}(\mathbb{R}^d) \subset L^2(\mathbb{R}^d)$ such that

$$\int_{\mathbb{R}^d} \left| \left(1 + \sum_{i=1}^d \left(\frac{\partial^2}{\partial u_i^2} + u_i^2 \right) \right)^{s/2} \hat{f}(u) \right|^2 du < \infty.$$

Let $\mathbf{t}, \mathbf{u} \in \mathbb{R}^d$. Then we claim for any $f \in \mathcal{S}^s(\mathbb{R}^d)$, there exists a function $\theta(\alpha)(\cdot) \in L^2(\mathbb{R}^d)$ such that

$$\lim_{|U(\mathbf{T})| \rightarrow \infty} \left\| \frac{1}{\text{vol}(U(\mathbf{T}))^{1/2}} \int_{U(\mathbf{T})} f(\mathbf{u} + \mathbf{t}) d\mathbf{t} - \theta_H(r_{\log \mathbf{T}}(\alpha))(\mathbf{u}) \text{Leb}(f) \right\|_{L^2(\mathbb{R}^d, d\mathbf{u})} = 0.$$

This is equivalent to the statement (42). By the standard Fourier transform on \mathbb{R}^d , equivalently

$$\lim_{|U(\mathbf{T})| \rightarrow \infty} \left\| \frac{1}{\text{vol}(U(\mathbf{T}))^{1/2}} \int_{U(\mathbf{T})} e^{i\mathbf{t} \cdot \hat{\mathbf{u}}} \hat{f}(\hat{\mathbf{u}}) d\mathbf{t} - \hat{\theta}_H(r_{\log \mathbf{T}}(\alpha))(\hat{\mathbf{u}}) \hat{f}(0) \right\|_{L^2(\mathbb{R}^d, d\hat{\mathbf{u}})} = 0.$$

For $\chi \in L^2(\mathbb{R}^d, d\hat{\mathbf{u}})$ and $\hat{\mathbf{u}} = (\hat{u}_j)_{1 \leq j \leq d}$, we denote

$$\chi_j(\hat{\mathbf{u}}) = \frac{e^{i\hat{u}_j} - 1}{i\hat{u}_j}, \quad \chi(\hat{\mathbf{u}}) = \prod_{j=1}^d \chi_j(\hat{\mathbf{u}}).$$

Let $\hat{\theta}(\alpha)(\hat{\mathbf{u}}) := \chi(\hat{\mathbf{u}})$ for all $\hat{\mathbf{u}} \in \mathbb{R}^d$. Now we will compute $\theta(r_{\log \mathbf{T}}(\alpha))$. By intertwining formula (11) for $\mathbf{T} \in \mathbb{R}^d$,

$$U_{\mathbf{T}}(f)(\hat{\mathbf{u}}) = \prod_{i=1}^d (T^{(i)})^{1/2} f(\mathbf{T}\hat{\mathbf{u}}), \text{ for } \mathbf{T}\hat{\mathbf{u}} = (T^{(1)}\hat{u}_1, \dots, T^{(d)}\hat{u}_d).$$

Then, for all $\alpha \in \text{Aut}_0(\mathbf{H}^g)$,

$$\hat{\theta}(r_{\log \mathbf{T}}(\alpha))(\hat{\mathbf{u}}) = U_{\mathbf{T}}(\chi)(\hat{\mathbf{u}}) = \text{vol}(U(\mathbf{T}))^{1/2} \chi(\mathbf{T}\hat{\mathbf{u}}).$$

The function $\theta(\alpha)$ is defined by inverse Fourier transform of $\hat{\theta}(\alpha)$ and

$$\|\theta_H(\alpha)\|_H = \|\theta(\alpha)\|_{L^2(\mathbb{R}^d)} = \|\hat{\theta}(\alpha)\|_{L^2(\mathbb{R}^d)} = \|\chi(\hat{\mathbf{u}})\|_{L^2(\mathbb{R}^d, d\hat{\mathbf{u}})} = C > 0.$$

By integration,

$$\begin{aligned} (43) \quad & \int_0^{T^{(d)}} \dots \int_0^{T^{(1)}} e^{i\mathbf{t} \cdot \hat{\mathbf{u}}} \hat{f}(\hat{\mathbf{u}}) d\mathbf{t} = \text{vol}(U(\mathbf{T})) \chi(\mathbf{T}\hat{\mathbf{u}}) \hat{f}(\hat{\mathbf{u}}) \\ & = \text{vol}(U(\mathbf{T})) \chi(\mathbf{T}\hat{\mathbf{u}}) (\hat{f}(\hat{\mathbf{u}}) - \hat{f}(0)) + \text{vol}(U(\mathbf{T}))^{1/2} \hat{\theta}(r_{\log \mathbf{T}}(\alpha))(\hat{\mathbf{u}}) \hat{f}(0). \end{aligned}$$

Then the claim reduces to the following:

$$\limsup_{|U(\mathbf{T})| \rightarrow \infty} \left\| \text{vol}(U(\mathbf{T}))^{1/2} \chi(\mathbf{T}\hat{\mathbf{u}}) (\hat{f}(\hat{\mathbf{u}}) - \hat{f}(0)) \right\|_{L^2(\mathbb{R}^d)} = 0.$$

If $f \in \mathcal{S}^s(\mathbb{R}^g)$ with $s > d/2$, function $\hat{f} \in C^0(\mathbb{R}^d)$ and bounded. Thus, by Dominated convergence theorem and change of variables,

$$\left\| \text{vol}(U(\mathbf{T}))^{1/2} \chi(\mathbf{T}\hat{\mathbf{u}}) (\hat{f}(\hat{\mathbf{u}}) - \hat{f}(0)) \right\|_{L^2(\mathbb{R}^d, d\hat{\mathbf{u}})} = \left\| \chi(\nu) \left(\hat{f}\left(\frac{\nu}{\mathbf{T}}\right) - \hat{f}(0) \right) \right\|_{L^2(\mathbb{R}^d, d\nu)} \rightarrow 0.$$

1

□

Corollary 4.2. *For any $s > d/2$, $\alpha \in \text{Aut}_0(\mathbf{H}^g)$ and $f \in W_{\alpha}^s(H)$, there exists a constant $C > 0$ such that*

$$\lim_{|U(\mathbf{T})| \rightarrow \infty} \frac{1}{\text{vol}(U(\mathbf{T}))^{1/2}} \left\| \left\langle \mathcal{P}_{U(\mathbf{T})}^{d, \alpha} m, \omega_f \right\rangle \right\|_{L^2(M)} = C |D_{\alpha}^H(f)|.$$

1 From Corollary 4.2, we derive the following limit result for the L^2 -norm of Bufe-
 2 tov functionals.

Corollary 4.3. *For every irreducible component H and $[\alpha] \in DC$, there exists $C > 0$ such that*

$$\lim_{|U(\mathbf{T})| \rightarrow \infty} \frac{1}{\text{vol}(U(\mathbf{T}))^{1/2}} \|\beta_H(\alpha, \cdot, \mathbf{T})\|_{L^2(M)} = C.$$

Proof. By the normalization of invariant distribution in Sobolev space, for any $\alpha \in \text{Aut}_0(\mathbf{H}^g)$, there exists a function $f_\alpha^H \in W_\alpha^s(H)$ such that $D_\alpha(f_\alpha^H) = \|f_\alpha^H\|_s = 1$. For all $[\alpha] \in DC(L)$, by asymptotic formula (41) for $f = f_H$,

$$|\langle \mathcal{P}_{U(\mathbf{T})}^{d,\alpha} m, \omega_f \rangle - \beta^f(\alpha, m, \mathbf{T})| \leq C_s(1 + L).$$

3 Therefore, L^2 -estimate follows from Corollary 4.2. \square

4 A relation between the Bufetov functional and the modular function θ_H is es-
 5 tablished below.

Corollary 4.4. *For every irreducible component $H \subset L^2(M)$, the following holds. For any $L > 0$ and any $r_{\mathbf{t}}$ -invariant probability measure μ supported on $DC(L) \subset \mathfrak{M}_g$,*

$$\beta_H(\alpha, \cdot, 1) = \theta_H(\alpha)(\cdot), \quad \text{for } \mu\text{-almost all } [\alpha] \in \mathfrak{M}_g.$$

6 *Proof.* By Theorem 1.6 and Lemma 4.1, there exists a constant $C > 0$ such that
 7 for all $[\alpha] \in \text{supp}(\mu) \subset DC(L)$ and $\mathbf{T} \in \mathbb{R}_+^d$, we have

$$(44) \quad \lim_{|U(\mathbf{T})| \rightarrow \infty} \|\beta_H(r_{\log \mathbf{T}}(\alpha), \cdot, 1) - \theta_H(r_{\log \mathbf{T}}(\alpha))\|_{L^2(M)} \leq \frac{C_\mu}{\text{vol}(U(\mathbf{T}))^{1/2}}.$$

8 By Luzin's theorem, for any $\delta > 0$ there exists a compact subset $E(\delta) \subset \mathfrak{M}$ such that
 9 we have the measure bound $\mu(\mathfrak{M} \setminus E(\delta)) < \delta$ and the function $\beta_H(\alpha, \cdot, 1) \in L^2(M)$
 10 depends continuously on $[\alpha] \in E(\delta)$. By Poincaré recurrence, there is a full measure
 11 set $F \subset \mathfrak{M}$ for \mathbb{R}^d -action. Denote by a full measure set $E'(\delta) = E(\delta) \cap F \subset E(\delta)$.

For every $\alpha_0 \in E'(\delta)$, there is a divergent sequence (\mathbf{t}_n) such that $\{r_{\mathbf{t}_n}(\alpha_0)\} \subset E(\delta)$ and $\lim_{n \rightarrow \infty} r_{\mathbf{t}_n}(\alpha_0) = \alpha_0$. By continuity of θ_H and β_H at α_0 , we have

$$(45) \quad \begin{aligned} \|\beta_H(\alpha_0, \cdot, 1) - \theta_H(\alpha_0)\|_{L^2(M)} \\ = \lim_{n \rightarrow \infty} \|\beta_H(r_{\mathbf{t}_n}(\alpha_0), \cdot, 1) - \theta_H(r_{\mathbf{t}_n}(\alpha_0))\|_{L^2(M)} = 0. \end{aligned}$$

12 Then $\beta_H(\alpha, \cdot, 1) = \theta_H(\alpha) \in L^2(M)$ for all $\alpha \in E'(\delta)$. It follows that the set where
 13 the equality (45) fails has a measure less than any $\delta > 0$, thus the identity holds
 14 for μ -almost all $\alpha \in \text{Aut}_0(\mathbf{H}^g)$. \square

15 For all $\alpha \in \text{Aut}_0(\mathbf{H}^g)$, smooth function $f \in W^s(M)$ for $s > s_{d,g} + 1/2$ decompose
 16 as an infinite sum, and the functional θ^f is defined by a convergent series

$$(46) \quad \theta^f(\alpha) := \sum_H D_\alpha^H(f) \theta_H(\alpha).$$

17 Hence, the modular function $\theta^f : \text{Aut}_0(\mathbf{H}^g) \rightarrow L^2(M)$ is continuous.

18 The following result is an extension of Lemma 4.1 to an asymptotic formula.

Lemma 4.5. *For all $\alpha \in \text{Aut}_0(\mathbf{H}^g)$, $f \in W^s(M)$ and $s > s_{d,g} + 1/2$,*

$$\lim_{n \rightarrow \infty} \left\| \frac{1}{\text{vol}(U(\mathbf{T}_n))^{1/2}} \left\langle \mathcal{P}_{U(\mathbf{T}_n)}^{d,\alpha} m, \omega_f \right\rangle - \theta^f(r_{\log \mathbf{T}_n}(\alpha)) \right\|_{L^2(M)} = 0.$$

1 We summarize our results on limit distributions for higher rank actions.

Theorem 4.6 (Theorem 1.7). *Let (\mathbf{T}_n) be any sequence such that*

$$\lim_{n \rightarrow \infty} r_{\log \mathbf{T}_n}[\alpha] = \alpha_\infty \in \mathfrak{M}_g.$$

2 *For every closed form $\omega_f \in \Lambda^d \mathfrak{p} \otimes W^s(M)$ with $s > s_{d,g} + 1/2$, which is not a*
 3 *coboundary, the limit distribution of the family of random variables $E_{\mathbf{T}_n}(f)$ ex-*
 4 *ists along a subsequence of $\{\mathbf{T}_n\}$ and is equal to the distribution of the function*
 5 $\theta^f(\alpha_\infty) = \beta(\alpha_\infty, \cdot, 1) \in L^2(M)$.

6 *If $\alpha_\infty \in DC$, then $\theta^f(\alpha_\infty)$ is a bounded function on M , and the limit distribution*
 7 *has compact support.*

8 *Proof of Theorem 1.7.* Since $\alpha_\infty \in \mathfrak{M}_g$, the existence of limit follows from the
 9 Lemma 4.5 and Theorem 4.6. \square

10 A relation with Birkhoff integrals and theta sum was introduced in [CF15, §5.3],
 11 and as an application, we derive the limit theorem of theta sums.

Corollary 4.7. *Let $\mathcal{Q}[x] = x^\top \mathcal{Q} x$ be the quadratic forms defined by $g \times g$ real matrix \mathcal{Q} , where $\alpha = \begin{pmatrix} I & 0 \\ \mathcal{Q} & I \end{pmatrix} \in Sp_{2g}(\mathbb{R})$ and $\ell(x) = \ell^\top x$ is the linear form defined by $\ell \in \mathbb{R}^g$. Then the theta sum*

$$\Theta(\mathcal{Q}, \ell; N) = N^{-g/2} \sum_{n \in \mathbb{Z}^g \cap [0, N]} \exp(2\pi i(\mathcal{Q}[n] + \ell(n)))$$

12 *has a limit distribution and it has compact support.*

13 5. L^2 -LOWER BOUNDS

14 In this section we prove L^2 - lower bounds of ergodic integrals on transverse torus.

5.1. Structure of return map. The polarized Heisenberg group $\mathbf{H}_{pol}^g \approx \mathbb{R}^g \times \mathbb{R}^g \times \mathbb{R}$ is equipped with the group law $(x, y, z) \cdot (x', y', z') = (x + x', y + y', z + z' + yx')$. Reduced standard Heisenberg group is defined by quotient

$$\mathbf{H}_{red}^g := \mathbf{H}_{pol}^g / (\{0\} \times \{0\} \times \frac{1}{2}\mathbb{Z}) \approx \mathbb{R}^g \times \mathbb{R}^g \times \mathbb{R} / \frac{1}{2}\mathbb{Z}.$$

15 Then the Reduced standard lattice is $\Gamma_{red}^g = \mathbb{Z}^g \times \mathbb{Z}^g \times \{0\} \subset \mathbf{H}_{red}^g$ and the quotient
 16 $\mathbf{H}_{red}^g / \Gamma_{red}^g$ is isomorphic to the standard Heisenberg manifold $M = \mathbf{H}^g / \Gamma$.

Given the standard frame (X_i, Y_i, Z) , $(g + 1)$ -dimensional (transverse) torus is denoted by

$$\mathbb{T}_\Gamma^{g+1} := \{\Gamma \exp(\sum_{i=1}^g y_i Y_i + z Z) \mid (y_i, z) \in \mathbb{R} \times \mathbb{R}\}.$$

Now, we consider a return map of $\mathbf{P}^{d,\alpha}$ on \mathbb{T}_Γ^{g+1} on the coordinates in reduced Heisenberg group. For $x = (x_1, \dots, x_g) \in \mathbb{R}^g$, we write in the coordinate of \mathbf{H}_{red}^g for convenience

$$\exp(x_1 X_1^\alpha + \dots + x_g X_g^\alpha) = (x_\alpha, x_\beta, w \cdot x), \text{ for some } x_\alpha, x_\beta, \text{ and } w \in \mathbb{R}^d.$$

Then by group law,

$$\exp(x_1 X_1^\alpha + \cdots + x_g X_g^\alpha) \cdot (0, y, z) = (x_\alpha, y + x_\beta, z + w \cdot x)$$

1 and given $(n, m, 0) \in \Gamma_{red}^g$,

$$(47) \quad \exp(x_1 X_1^\alpha + \cdots + x_g X_g^\alpha) \cdot (0, y, z) \cdot (n, m, 0) = \exp(x'_1 X_1^\alpha + \cdots + x'_g X_g^\alpha) \cdot (0, y', z')$$

2 if and only if

$$x'_\alpha = x_\alpha + n, y' = y + (x_\beta - x'_\beta) + m \text{ and } z' = z + (w - w') \cdot x + n^\top (y + x_\beta).$$

Assume $\langle X_i^\alpha, X_j \rangle \neq 0$ for all i, j , and we write the first return time for $\mathbf{P}^{d, \alpha}$ action

$$t_{Ret} = (t_{Ret,1}, \dots, t_{Ret,d}) \in \mathbb{R}^d$$

3 on transverse torus \mathbb{T}_Γ^{g+1} . We denote the domain for return time $U(t_{Ret}) =$
4 $[0, t_{Ret,1}] \times \cdots \times [0, t_{Ret,d}]$. Return map of action $\mathbf{P}^{d, \alpha}$ on \mathbb{T}_Γ^{g+1} has a form of
5 skew-shift

$$(48) \quad A_{\rho, \tau}(y, z) = (y + \rho, z + v \cdot y + \tau) \text{ on } \mathbb{R}^g / \mathbb{Z}^g \times \mathbb{R} / K^{-1} \mathbb{Z}$$

6 for some non-zero vectors $\rho, v \in \mathbb{R}^g$ and $\tau \in \mathbb{R}$.

7 Furthermore, $A_{\rho, \tau} = A_{d, \rho, \tau} \circ \cdots \circ A_{1, \rho, \tau}$ decomposes with commuting linear
8 skew-shifts

$$(49) \quad A_{i, \rho, \tau}(y, z) = (y + \rho_i, z + v_i \cdot y + \tau_i) \text{ on } \mathbb{R}^g / \mathbb{Z}^g \times \mathbb{R} / K^{-1} \mathbb{Z}$$

for some $\rho_i, v_i \in \mathbb{R}^g$ and $\tau_i \in \mathbb{R}$. For each $j \neq k$, it is easily verified that

$$A_{j, \rho, \tau} \circ A_{k, \rho, \tau} = A_{k, \rho, \tau} \circ A_{j, \rho, \tau}.$$

Given pair $(\mathbf{m}, n) \in \mathbb{Z}_{K|n|}^g \times \mathbb{Z}$, let $H_{(\mathbf{m}, n)}$ denote the corresponding factor and $C^\infty(H_{(\mathbf{m}, n)})$ be a subspace of smooth functions on $H_{(\mathbf{m}, n)}$. Denote $\{e_{\mathbf{m}, n} \mid (\mathbf{m}, n) \in \mathbb{Z}_{|n|}^g \times \mathbb{Z}\}$ the basis of characters on \mathbb{T}_Γ^{g+1} and for all $(y, z) \in \mathbb{T}^g \times \mathbb{T}$,

$$e_{\mathbf{m}, n}(y, z) := \exp[2\pi i(\mathbf{m} \cdot y + nKz)].$$

For each $A_{i, \rho, \sigma}$, we set $\mathbf{v}_i = (v_{i1}, \dots, v_{ig}) \in \mathbb{Z}_{K|n|}^g$. Then the orbit can be identified with the following dual orbit

$$\begin{aligned} \mathcal{O}_{A_i}(\mathbf{m}, n) &= \{(\mathbf{m} + (nj_i)\mathbf{v}_i, n), j_i \in \mathbb{Z}\} \\ &= \{(m_1 + (nv_{i1})j_i, \dots, m_g + (nv_{ig})j_i, n), j_i \in \mathbb{Z}\}. \end{aligned}$$

If $n = 0$, the orbit $[(\mathbf{m}, 0)] \subset \mathbb{Z}^g \times \mathbb{Z}$ of $(\mathbf{m}, 0)$ is reduced to a single element. If $n \neq 0$, then the dual orbit $[(\mathbf{m}, n)] \subset \mathbb{Z}^{g+1}$ of (\mathbf{m}, n) for higher rank actions is described as follows:

$$\mathcal{O}_A(\mathbf{m}, n) = \{(m_k + n \sum_{i=1}^d (v_{ik} j_i), n)_{1 \leq k \leq d} : j = (j_1, \dots, j_d) \in \mathbb{Z}^d\}.$$

It follows that every A -orbit for rank \mathbb{R}^d -action (or A_i -orbit) can be labeled uniquely by a pair $(\mathbf{m}, n) \in \mathbb{Z}_{|n|}^g \times \mathbb{Z} \setminus \{0\}$ with $\mathbf{m} = (m_1, \dots, m_g)$. Thus, the subspace of functions with non-zero central characters splits as a direct sum of components $H_{(\mathbf{m}, n)}$

$$L^2(\mathbb{T}_\Gamma^{g+1}) = \bigoplus_{\omega \in \mathcal{O}_A} H_\omega, \quad \text{where } H_\omega = \bigoplus_{(\mathbf{m}, n) \in \omega} \mathbb{C} e_{(\mathbf{m}, n)}.$$

5.2. **Higher cohomology for \mathbb{Z}^d -action of skew-shifts.** In this subsection, we will find a relations with the return map for \mathbb{Z}^d action of $P^{d,\alpha}$ on the torus \mathbb{T}_Γ^{g+1} and obstructions to cohomological equation $\omega = d\Omega$.

We will restrict our interest to the following cocycle equation

$$(50) \quad \varphi(x, t) = \mathfrak{D}\Phi(x, t), \quad x \in \mathbb{T}^g, \quad t \in \mathbb{Z}^d$$

where d -cocycle $\varphi : \mathbb{T}_\Gamma^{g+1} \times \mathbb{Z}^d \rightarrow \mathbb{R}$, $(d-1)$ cochain $\Phi : \mathbb{T}_\Gamma^{g+1} \rightarrow \mathbb{R}^d$, $\Phi = (\Phi_1, \dots, \Phi_d)$, and \mathfrak{D} is coboundary operator

$$\mathfrak{D}\Phi = \sum_{i=1}^d (-1)^{i+1} \Delta_i \Phi_i$$

where $\Delta_i \Phi_i = \Phi_i \circ A_{i,\rho,\tau} - \Phi_i$.

In the work of Katok [KK95, §2], he proved the existence of solutions for cocycle equations by studying the dual equations $\hat{\varphi} = \mathfrak{D}\hat{\Phi}$ in the space of Fourier coefficients (dual orbit). We apply this result to our A -orbit and verify the invariant distributions (or currents) explicitly.

Proposition 5.1. [KK95, Proposition 2.2] *A dual cocycle $\hat{\varphi}$ satisfies cocycle equation (50) if and only if $\sum_{j \in \mathbb{Z}^d} \hat{\varphi}(\mathbf{m}, n) \circ A^j = 0$.*

For fixed $(\mathbf{m}, n) \in \mathbb{Z}^g \times \mathbb{Z}$, we denote an obstruction of cohomological equation restricted to the A -orbit of (\mathbf{m}, n) by $\mathcal{D}_{\mathbf{m},n}(\varphi) = \sum_{j \in \mathbb{Z}^d} \hat{\varphi}(\mathbf{m}, n) \circ A^j$. Since A^j is composition of commuting toral automorphisms, we obtain the following generalized formula (see [AFU11, §5] and [Ka03, §11] for rank 1 map on \mathbb{T}^2).

Lemma 5.2. *There exists a distributional obstruction to the existence of a smooth solution $\varphi \in C^\infty(H_{(\mathbf{m},n)})$ of the cohomological equation (50). A generator of the space of invariant distribution $\mathcal{D}_{\mathbf{m},n}$ is given by*

$$\mathcal{D}_{\mathbf{m},n}(e_{a,b}) := e^{-2\pi i \sum_{i=1}^d [(\mathbf{m} \cdot \rho_i + nK\tau_i)j_i + nK\tau_i \binom{j_i}{2}]}$$

if $(a, b) = (m_k + K \sum_{i=1}^d (v_{ik}j_i), n)_{1 \leq k \leq g}$ and 0 otherwise.

Proof. From previous observation, there exists an obstruction

$$(51) \quad \mathcal{D}_{\mathbf{m},n}(\varphi) = \sum_{j \in \mathbb{Z}^d} \int_{\mathbb{T}_\Gamma^{g+1}} \varphi(x, y) \overline{e_{\mathbf{m},n} \circ A_{\rho,\tau}^j} dx dy.$$

By direct computation, for fixed $j = (j_1, \dots, j_d)$,

$$e_{\mathbf{m},n} \circ A_{\rho,\tau}^j(y, z) = \prod_{i=1}^d \left(e^{2\pi i [(\mathbf{m} \cdot \rho_i + nK\tau_i)j_i + nK\tau_i \binom{j_i}{2}]} \right) \left(e^{2\pi i (\mathbf{m} \cdot y + K(z + n \sum_{k=1}^d (v_{ik}j_i)y_k))} \right).$$

Then, we choose $\hat{\varphi} = e_{a,b}$ for $(a, b) = (m_k + K \sum_{i=1}^d (v_{ik}j_i), n)_{1 \leq k \leq g}$ in the non-trivial orbit ($n \neq 0$),

$$(52) \quad \mathcal{D}_{\mathbf{m},n}(e_{a,b}) = e^{-2\pi i \sum_{i=1}^d [(\mathbf{m} \cdot \rho_i + nK\tau_i)j_i + nK\tau_i \binom{j_i}{2}]}.$$

5.3. Changes of coordinates. For any frame $(X_i^\alpha, Y_i^\alpha, Z)_{i=1}^g$ and any $m \in M$, denote a transverse cylinder

$$\mathcal{C}_{\alpha,m} := \{m \exp(\sum_{i=1}^g y'_i Y_i^\alpha + z'Z) \mid (y', z') \in U(t_{Ret}^{-1}) \times \mathbb{T}\} \subset M.$$

For any $\xi \in \mathbb{T}_\Gamma^{g+1}$, let $\xi' \in \mathcal{C}_{\alpha,m}$ denote first intersection of the orbit $\{P_t^{d,\alpha}(\xi) \mid t \in \mathbb{R}_+^d\}$ with transverse cylinder $\mathcal{C}_{\alpha,m}$. Then, there exists a first return time to the cylinder $t(\xi) = (t_1(\xi), \dots, t_d(\xi)) \in \mathbb{R}_+^d$ such that the map $\Phi_{\alpha,m} : \mathbb{T}_\Gamma^{g+1} \rightarrow \mathcal{C}_{\alpha,m}$ is defined by

$$\xi' = \Phi_{\alpha,m}(\xi) = P_{t(\xi)}^{d,\alpha}(\xi), \quad \forall \xi \in \mathbb{T}_\Gamma^{g+1}.$$

Let (y, z) and (y', z') denote the coordinates on \mathbb{T}_Γ^{g+1} and $\mathcal{C}_{\alpha,m}$ respectively, given by the exponential map

$$(y, z) \rightarrow \xi_{y,z} := \Gamma \exp(\sum_{i=1}^g y_i Y_i + zZ), \quad (y', z') \rightarrow m \exp(\sum_{i=1}^g y'_i Y_i^\alpha + z'Z).$$

- 1 Recall that if $\alpha \in Sp_{2g}(\mathbb{R})$, then for $1 \leq i, j \leq g$ there exist matrices $A =$
 2 $(a_{ij}), B = (b_{ij}), C = (c_{ij}), D = (d_{ij})$ such that

$$\alpha := \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in Sp_{2g}(\mathbb{R}),$$

satisfying $A^t D - C^t B = I_{2g}$, $C^t A = A^t C$, $D^t B = B^t D$, and $\det(A) \neq 0$. Set

$$X_i^\alpha = \sum_{j=1}^g (a_{ij} X_j + b_{ij} Y_j) + w_i Z \quad \text{and} \quad Y_i^\alpha = \sum_{j=1}^g (c_{ij} X_j + d_{ij} Y_j) + v_i Z.$$

- 3 Let $x = \Gamma \exp(\sum_{i=1}^d y_{x,i} Y_i + z_x Z) \exp(\sum_{i=1}^d t_{x,i} X_i)$, for some $(y_x, z_x) \in \mathbb{T}^d \times$
 4 $\mathbb{R}/K\mathbb{Z}$ and $t_x = (t_{x,i}) \in [0, 1]^d$. Then, the map $\Phi_{\alpha,x} : \mathbb{T}_\Gamma^{g+1} \rightarrow \mathcal{C}_{\alpha,x}$ is defined by
 5 $\Phi_{\alpha,x}(y, z) = (y', z')$ where

$$(53) \quad \begin{bmatrix} y'_1 \\ y'_2 \\ \vdots \\ y'_g \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1g} \\ a_{21} & a_{22} & \cdots & a_{2g} \\ \vdots & \vdots & \ddots & \vdots \\ a_{g1} & a_{g2} & \cdots & a_{gg} \end{bmatrix} \begin{bmatrix} y_1 - y_{x,1} \\ y_2 - y_{x,2} \\ \vdots \\ y_g - y_{x,g} \end{bmatrix} + \begin{bmatrix} b_{11} & \cdots & b_{1g} \\ b_{21} & \cdots & b_{2g} \\ \vdots & \vdots & \vdots \\ b_{g1} & \cdots & b_{gg} \end{bmatrix} \begin{bmatrix} t_{x,1} \\ t_{x,2} \\ \vdots \\ t_{x,g} \end{bmatrix},$$

- 6 and $z' = z + P(\alpha, x, y)$ for some degree 4 polynomial P .
 Therefore, the map $\Phi_{\alpha,x}$ is invertible with

$$\Phi_{\alpha,x}^*(dy'_1 \wedge \cdots \wedge dy'_g \wedge dz') = \frac{1}{\det(A)} dy_1 \wedge \cdots \wedge dy_g \wedge dz.$$

- 7 Since $A^t D - C^t B = I_{2g}$, by direct computation we obtain return time, we have

$$(54) \quad \begin{bmatrix} t_1(\xi) \\ t_2(\xi) \\ \vdots \\ t_g(\xi) \end{bmatrix} = \begin{bmatrix} d_{11} & \cdots & d_{1g} \\ d_{21} & \cdots & d_{2g} \\ \vdots & \vdots & \vdots \\ d_{d1} & \cdots & d_{dg} \end{bmatrix} \begin{bmatrix} t_{x,1} \\ t_{x,2} \\ \vdots \\ t_{x,d} \end{bmatrix} + \begin{bmatrix} c_{11} & c_{12} & \cdots & c_{1g} \\ c_{21} & c_{22} & \cdots & c_{2g} \\ \vdots & \vdots & \vdots & \vdots \\ c_{g1} & c_{g2} & \cdots & c_{gg} \end{bmatrix} \begin{bmatrix} y_1 - y_{x,1} \\ y_2 - y_{x,2} \\ \vdots \\ y_g - y_{x,g} \end{bmatrix}.$$

Then,

$$\|t(\xi)\| \leq \max_i |t_i(\xi)|^g \leq \max_i \left| \sum_{j=1}^g d_{ij} t_{x,i} + c_{ij} (y_i - y_{x,i}) \right|^g \leq \max_i \|Y_i^\alpha\|^g.$$

- 1 5.4. **L^2 -lower bound of functional.** We will prove the bounds for square mean
 2 of integrals along leaves of foliations of the torus \mathbb{T}_Γ^{g+1} .

Lemma 5.3. *There exists a constant $C > 0$ such that for all $\alpha = (X_i^\alpha, Y_i^\alpha, Z)$, and for every irreducible component $H := H_n$ of central parameter $n \neq 0$, there exists a function f_H such that*

$$\|f_H\|_{L^\infty(H)} \leq C \text{vol}(U(t_{Ret}))^{-1} |\mathcal{D}_\alpha^H(f_H)|,$$

$$\|f_H\|_{\alpha,s} \leq C \text{vol}(U(t_{Ret}))^{-1} |\mathcal{D}_\alpha^H(f_H)| \left(1 + \frac{\Sigma(t_{Ret})}{\text{vol}(U(t_{Ret}))} \|Y\| \right)^s (1+n^2)^{s/2}$$

- 3 where $\|Y\| := \max_{1 \leq i \leq g} \|Y_i^\alpha\|$ and $\Sigma(t_{Ret}) = \sum_{i=1}^g t_{Ret,i}$.

- 4
 5 On rectangular domain $U(\mathbf{T})$, for all $m \in \mathbb{T}_\Gamma^{g+1}$ and $T^{(i)} \in \mathbb{Z}_{t_{Ret,i}}$

$$(55) \quad \left\| \left\langle \mathcal{P}_{U(\mathbf{T})}^{d,\alpha}(\mathbf{Q}_y^{g,Y} m), \omega_H \right\rangle \right\|_{L^2(\mathbb{T}^g, dy)} = |\mathcal{D}_\alpha^H(f_H)| \left(\frac{\text{vol}(U(\mathbf{T}))}{\text{vol}(U(t_{Ret}))} \right)^{1/2}.$$

In addition, whenever $H \perp H' \subset L^2(M)$ the functions

$$\left\langle \mathcal{P}_{U(\mathbf{T})}^{d,\alpha}(\mathbf{Q}_y^{g,Y} m), \omega_H \right\rangle \quad \text{and} \quad \left\langle \mathcal{P}_{U(\mathbf{T})}^{d,\alpha}(\mathbf{Q}_y^{g,Y} m), \omega_{H'} \right\rangle$$

- 6 are orthogonal in $L^2(\mathbb{T}^g, dy)$.

- 7 *Proof.* The operator $I_\alpha : L^2(M) \rightarrow L^2(\mathbb{T}_\Gamma^{g+1})$ is defined by

$$(56) \quad f \rightarrow I_\alpha(f) := \int_{U(t_{Ret})} f \circ \mathbf{P}_s^{d,\alpha}(\cdot) ds.$$

Then operator I_α is surjective linear map of $L^2(M)$ onto $L^2(\mathbb{T}_\Gamma^{g+1})$ with a right inverse R_α^χ defined as follows. Let $\chi \in C_0^\infty(0,1)^g$ be any function of jointly integrable with integral 1. For any $F \in L^2(\mathbb{T}_\Gamma^{g+1})$, let $R_\alpha^\chi(F) \in L^2(M)$ be a function defined by

$$R_\alpha^\chi(F)(\mathbf{P}_v^{d,\alpha}(x)) = \frac{1}{\text{vol}(U(t_{Ret}))} \chi\left(\frac{v}{t_{Ret}}\right) F(x), \quad (x, v) \in \mathbb{T}_\Gamma^{g+1} \times U(t_{Ret}).$$

Then, it follows that there exists a constant $C_\chi > 0$ such that

$$(57) \quad \begin{aligned} \|R_\alpha^\chi(F)\|_{\alpha,s} &\leq C_\chi \text{vol}(U(t_{Ret}))^{-1} \left(1 + \sum_{i=1}^g t_{Ret,i}^{-1} \|Y_i^\alpha\| \right)^s \|F\|_{W^s(\mathbb{T}_\Gamma^{g+1})} \\ &\leq C_\chi \text{vol}(U(t_{Ret}))^{-1} \left(1 + \frac{\Sigma(t_{Ret})}{\text{vol}(U(t_{Ret}))} \|Y\| \right)^s \|F\|_{W^s(\mathbb{T}_\Gamma^{g+1})}. \end{aligned}$$

As explained in §5.1, the space $L^2(\mathbb{T}_\Gamma^{g+1})$ decompose as a direct sum of irreducible subspaces invariant under the action of each $A_{j,\rho,\sigma}$. It follows that the subspace of functions with non-zero central character can be split as direct sum of components

$H_{(\mathbf{m},n)}$ with $(\mathbf{m},n) \in \mathbb{Z}_{|n|}^g \times \mathbb{Z} \setminus \{0\}$ with $\mathbf{m} = (m_1, \dots, m_g)$. For a function $F \in H_{(\mathbf{m},n)}$, it is characterized by Fourier expansion

$$F = \sum_{j \in \mathbb{Z}^d} F_j e^{A^j(\mathbf{m},n)} = \sum_{j \in \mathbb{Z}^d} F_j e^{(m_k + K \sum_{i=1}^d (v_{ik} j_i), n)}.$$

Then, we can choose $f_H := R_\alpha^\chi(e_{\mathbf{m},n}) \in C^\infty(H)$ such that

$$(58) \quad |D_\alpha(f_\alpha^H)| = |D_{\mathbf{m},n}(e_{\mathbf{m},n})| = 1,$$

$$(59) \quad \int_{U(t_{Ret})} f_H \circ \mathbf{P}_t^{d,\alpha}(y, z) dt = e_{\mathbf{m},n}(y, z), \text{ for } (y, z) \in \mathbb{T}_\Gamma^{g+1}.$$

Therefore, it follows from (57) that

$$|f_H|_{L^\infty(H)} \leq C_\chi \text{vol}(U(t_{Ret}))^{-1},$$

$$\|f_H\|_{\alpha,s} \leq C \text{vol}(U(t_{Ret}))^{-1} |\mathcal{D}_\alpha^H(f_H)| \left(1 + \frac{\Sigma(t_{Ret})}{\text{vol}(U(t_{Ret}))} \|Y\|\right)^s (1+n^2)^{s/2}.$$

Moreover, since $\{e_{\mathbf{m},n} \circ A_{\rho,\tau}^j\}_{j \in \mathbb{Z}^d} \subset L^2(\mathbb{T}_\Gamma^g, dy)$ is orthonormal, we verify

$$\begin{aligned} \left\| \left\langle \mathcal{P}_{U(\mathbf{T})}^{d,\alpha}(\mathbf{Q}_y^{g,Y} x), \omega_H \right\rangle \right\|_{L^2(\mathbb{T}^g, dy)} &= \left\| \sum_{j_d=0}^{\lfloor \frac{T(d)}{t_{Ret,d}} \rfloor - 1} \cdots \sum_{j_1=0}^{\lfloor \frac{T(1)}{t_{Ret,1}} \rfloor - 1} e_{\mathbf{m},n} \circ A_{\rho,\tau}^j \right\|_{L^2(\mathbb{T}^g, dy)} \\ &= \left(\frac{\text{vol}(U(\mathbf{T}))}{\text{vol}(U(t_{Ret}))} \right)^{1/2}. \end{aligned}$$

1

□

2 Recall that

$$(60) \quad L^2(M) = \bigoplus_{n \in \mathbb{Z}} H_n := \bigoplus_{n \in \mathbb{Z}} \bigoplus_{i=1}^{\mu(n)} H_{i,n}$$

3 where $H_n = \bigoplus_{i=1}^{\mu(n)} H_{i,n}$ is irreducible representation with a central parameter n
 4 and $\mu(n)$ is countable by Howe-Richardson multiplicity formula.

For any infinite dimensional vector $\mathbf{c} := (c_{i,n}) \in \ell^2$, let $\beta_{\mathbf{c}}$ denote Bufetov functional

$$\beta_{\mathbf{c}} = \sum_{n \in \mathbb{Z}} \sum_{i=1}^{\mu(n)} c_{i,n} \beta_{i,n}^{i,n}.$$

By orthogonal property and from Corollary 4.3, the function $\beta_{\mathbf{c}}(\alpha, \cdot, \mathbf{T}) \in L^2(M)$ for all $(\alpha, \mathbf{T}) \in \text{Aut}_0(\mathbf{H}^g) \times \mathbb{R}_+^d$. Furthermore,

$$\|\beta_{\mathbf{c}}(\alpha, \cdot, \mathbf{T})\|_{L^2(M)}^2 = \sum_{n \in \mathbb{Z}} \sum_{i=1}^{\mu(n)} |c_{i,n}|^2 \|\beta_{i,n}^{i,n}(\alpha, \cdot, \mathbf{T})\|_{L^2(H_{i,n})}^2 \leq C^2 |\mathbf{c}|_{\ell^2}^2 \text{vol}(U(\mathbf{T})).$$

5 For any $\mathbf{c} := (c_{i,n})$, let $|\mathbf{c}|_s$ denote the norm defined by

$$(61) \quad |\mathbf{c}|_s^2 = \sum_{n \in \mathbb{Z} \setminus \{0\}} \sum_{i=1}^{\mu(n)} (1 + K^2 n^2)^s |c_{i,n}|^2.$$

Lemma 5.4. *For any $s > s_{d,g} + 1/2$, there exists a constant $C_s > 0$ such that for all $\alpha \in DC(L)$, for all $\mathbf{c} \in \ell^2$, for all $z \in \mathbb{T}$ and all $T > 0$,*

$$\begin{aligned} & \left| \left\| \beta_{\mathbf{c}}(\alpha, \Phi_{\alpha,x}(\xi_{y,z}), \mathbf{T}) \right\|_{L^2(\mathbb{T}^g, dy)} - \left(\frac{\text{vol}(U(\mathbf{T}))}{\text{vol}(U(t_{Ret}))} \right)^{1/2} |\mathbf{c}|_0 \right| \\ & \leq C_s (\text{vol}(U(t_{Ret})) + \text{vol}(U(t_{Ret}))^{-1}) (1 + L) \left(1 + \frac{\Sigma(t_{Ret})}{\text{vol}(U(t_{Ret}))} \|Y\| \right)^s |\mathbf{c}|_s. \end{aligned}$$

Proof. By Lemma 5.2, there exists a function $f_{i,n} \in C^\infty(H_{i,n})$ with $|\mathcal{D}^{i,n}(f_{i,n})| = 1$. Let $f_{\mathbf{c}} = \sum_{n \in \mathbb{Z}} \sum_{i=1}^{\mu(n)} c_{i,n} f_{i,n} \in C^\infty(M)$, summing up all the functions on irreducibles. Then by the estimates in the Lemma 5.3 and (61),

$$(62) \quad \|f_{\mathbf{c}}\|_{L^\infty(M)} \leq C |\mathbf{c}|_{\ell^1}.$$

$$(63) \quad \|f_{\mathbf{c}}\|_{\alpha,s} \leq C \text{vol}(U(t_{Ret}))^{-1} \left(1 + \frac{\Sigma(t_{Ret})}{\text{vol}(U(t_{Ret}))} \|Y\| \right)^s |\mathbf{c}|_s.$$

By orthogonality,

$$\left\| \left\langle \mathcal{P}_{U(\mathbf{T})}^{d,\alpha} \circ \mathbf{Q}_y^{g,Y}, \omega_{\mathbf{c}} \right\rangle \right\|_{L^2(\mathbb{T}^g, dy)} = \left(\frac{\text{vol}(U(\mathbf{T}))}{\text{vol}(U(t_{Ret}))} \right)^{1/2} |\mathbf{c}|_0.$$

By the estimate in Lemma 5.3 for each $f_{i,n}$, for every $z \in \mathbb{T}$ and all $\mathbf{T} \in \mathbb{R}_+^d$, we have

$$\left\| \left\langle \mathcal{P}_{U(\mathbf{T})}^{d,\alpha}(\Phi_{\alpha,x}(\xi_{y,z})), \omega_{\mathbf{c}} \right\rangle - \left\langle \mathcal{P}_{U(\mathbf{T})}^{d,\alpha}(\xi_{y,z}), \omega_{\mathbf{c}} \right\rangle \right\|_{L^2(\mathbb{T}^g, dy)} \leq 2 \|f_{\mathbf{c}}\|_{L^\infty(M)} \|Y\|.$$

Let $U(\mathbf{T}_{Ret}) = [0, \mathbf{T}_{Ret,1}] \times \cdots \times [0, \mathbf{T}_{Ret,g}]$ where $\mathbf{T}_{Ret,i} := t_{Ret,i}([T^{(i)}/t_{Ret,i}] + 1)$. Then,

$$\left\| \left\langle \mathcal{P}_{U(\mathbf{T})}^{d,\alpha}(\xi_{y,z}), \omega_{\mathbf{c}} \right\rangle - \left\langle \mathcal{P}_{U(\mathbf{T}_{Ret})}^{d,\alpha}(\xi_{y,z}), \omega_{\mathbf{c}} \right\rangle \right\|_{L^2(\mathbb{T}^g, dy)} \leq \text{vol}(U(t_{Ret})) \|f_{\mathbf{c}}\|_{L^\infty(M)}.$$

Therefore, there exists a constant $C' > 0$ such that

$$\begin{aligned} & \left| \left\| \left\langle \mathcal{P}_{U(\mathbf{T})}^{d,\alpha}(\Phi_{\alpha,x}(\xi_{y,z})), \omega_{\mathbf{c}} \right\rangle \right\|_{L^2(\mathbb{T}^g, dy)} - \left(\frac{\text{vol}(U(\mathbf{T}))}{\text{vol}(U(t_{Ret}))} \right)^{1/2} |\mathbf{c}|_0 \right| \\ & \leq C' \text{vol}(U(t_{Ret})) |\mathbf{c}|_{\ell^1}. \end{aligned}$$

For all $s > s_{d,g} + 1/2$, by asymptotic property of Theorem 1.6, for some constant $C_s > 0$,

$$\left| \left\langle \mathcal{P}_{U(\mathbf{T})}^{d,\alpha} m, \omega \right\rangle - \beta_H(\alpha, m, \mathbf{T}) \mathcal{D}_\alpha^H(f_H) \right| \leq C_s (1 + L) \|f\|_{\alpha,s}.$$

Applying $\beta_{\mathbf{c}} = \beta^{f_{\mathbf{c}}}$ and combining bounds on the function $f_{\mathbf{c}}$ with (62),

$$\begin{aligned} & \left| \left\| \beta_{\mathbf{c}}(\alpha, \Phi_{\alpha,x}(\xi_{y,z}), \mathbf{T}) \right\|_{L^2(\mathbb{T}^g, dy)} - \left(\frac{\text{vol}(U(\mathbf{T}))}{\text{vol}(U(t_{Ret}))} \right)^{1/2} |\mathbf{c}|_0 \right| \\ & \leq C' \text{vol}(U(t_{Ret})) |\mathbf{c}|_{\ell^1} + C_s \text{vol}(U(t_{Ret}))^{-1} (1 + L) \|f_{\mathbf{c}}\|_{\alpha,s} \\ & \leq C'_s (\text{vol}(U(t_{Ret})) + \text{vol}(U(t_{Ret}))^{-1}) (1 + L) \left(1 + \frac{\Sigma(t_{Ret})}{\text{vol}(U(t_{Ret}))} \|Y\| \right)^s |\mathbf{c}|_s. \end{aligned}$$

1 Therefore, we derive the estimates in the statement. \square

6. ANALYTICITY OF FUNCTIONALS

In this section we prove that for all $\alpha \in DC$, the Bufetov functionals on any square are real analytic.

6.1. Analyticity. By the orthogonal property of Bufetov cocycle β_H on an irreducible component $H = H_n$ with central parameter $n \in \mathbb{Z} \setminus \{0\}$, the following statement is immediate. For any $(m, \mathbf{T}) \in M \times \mathbb{R}_+^d$ and $t \in \mathbb{R}$,

$$(64) \quad \beta_H(\alpha, \phi_t^Z(m), \mathbf{T}) = e^{2\pi i K n t} \beta_H(\alpha, m, \mathbf{T}).$$

Definition 6.1. For every $t \in \mathbb{R}$, $1 \leq i \leq d$, and $m \in M$, the *stretched (in direction of Z) rectangle* is denoted by

$$(65) \quad [\Gamma_{\mathbf{T}}]_{i,t}^Z(m) := \{(\phi_{ts_i}^Z) \circ \mathbf{P}_{\mathbf{s}}^{d,\alpha}(m) \mid \mathbf{s} \in U(\mathbf{T})\}.$$

For $\mathbf{s} = (s_1, \dots, s_d) \in \mathbb{R}^d$, let us denote by the standard rectangle $\Gamma_{\mathbf{T}}(\mathbf{s}) := (\gamma_1(s_1), \dots, \gamma_d(s_d))$ for $\gamma_i(s_i) = \exp(s_i X_i)$. Similarly, we also write the stretched rectangle

$$(66) \quad [\Gamma_{\mathbf{T}}]_{i,t}^Z(\mathbf{s}) := (\gamma_1(s_1), \dots, \gamma_{i,t}^Z(s_i), \dots, \gamma_d(s_d))$$

where $\gamma_{i,t}^Z(s_i) := \phi_{ts_i}^Z(\gamma_i(s_i))$ is a stretched curve.

Definition 6.2. The *restricted rectangle* $\Gamma_{T,i,s}$ of the standard rectangle $\Gamma_{\mathbf{T}}$ is defined as a restriction on i -th coordinate given by

$$\Gamma_{T,i,s}(\mathbf{s}) := \Gamma_{\mathbf{T}}|_{U_{T,i,s}}(\mathbf{s}),$$

where $U_{T,i,s} = [0, T^{(1)}] \times \dots \times \underbrace{[0, s]}_{i\text{-th}} \times \dots \times [0, T^{(d)}]$ for some $0 < s \leq T^{(i)}$.

Lemma 6.3. For fixed elements (X_i, Y_i, Z) satisfying commutation relation (1), the following formula for rank 1 action holds:

$$\hat{\beta}_H(\alpha, [\Gamma_{\mathbf{T}}]_{i,t}^Z) = e^{2\pi i n K T^{(i)}} \hat{\beta}_H(\alpha, \Gamma_{\mathbf{T}}) - 2\pi i n K t \int_0^{T^{(i)}} e^{2\pi i n K t s_i} \hat{\beta}_H(\alpha, \Gamma_{T,i,s}) ds_i.$$

Proof. Let $\alpha = (X_i, Y_i, Z)$ and ω be d -form supported on a single irreducible representation H . We obtain following the formula for stretches of curve $\gamma_{i,t}^Z$ (see [FK20b, §4 and Lemma 9.1]),

$$\frac{d\gamma_{i,t}^Z}{ds_i} = D\phi_{ts_i}^Z\left(\frac{d\gamma_i}{ds_i}\right) + tZ \circ \gamma_{i,t}^Z.$$

It follows that pairing is given by

$$\begin{aligned} \langle [\Gamma_{\mathbf{T}}]_{i,t}^Z, \omega \rangle &= \int_{U(\mathbf{T})} \omega\left(\frac{d\gamma_1}{ds_1}(s_1), \dots, \frac{d\gamma_{i,t}^Z}{ds_i}(s_i), \dots, \frac{d\gamma_d}{ds_d}(s_d)\right) d\mathbf{s} \\ &= \int_{U(\mathbf{T})} e^{2\pi i n K t s_i} [\omega\left(\frac{d\gamma_1}{ds_1}(s_1), \dots, \frac{d\gamma_d}{ds_d}(s_d)\right)] + \iota_Z \omega \circ [\Gamma_{\mathbf{T}}]_{i,t}^Z(\mathbf{s}) d\mathbf{s} \end{aligned}$$

Denote $d - 1$ dimensional triangle $U_{d-1}(\mathbf{T})$ with $U(\mathbf{T}) = U_{d-1}(\mathbf{T}) \times [0, T^{(i)}]$. Integration by parts for a fixed i -th integral gives

$$\begin{aligned} & \int_{U(\mathbf{T})} e^{2\pi i n K t s_i} [\omega(\frac{d\gamma_1}{ds_1}(s_1), \dots, \frac{d\gamma_d}{ds_d}(s_d))] ds \\ &= e^{2\pi i n K t T^{(i)}} \int_{U(\mathbf{T})} [\omega(\frac{d\gamma_1}{ds_1}(s_1), \dots, \frac{d\gamma_d}{ds_d}(s_d))] ds \\ &- 2\pi i n K t \int_0^{T^{(i)}} e^{2\pi i n K t s_i} \int_{U_{d-1}(\mathbf{T})} \left(\int_0^{s_i} [\omega(\frac{d\gamma_1}{ds_1}(s_1), \dots, \frac{d\gamma_i}{ds_i}(r) \dots, \frac{d\gamma_d}{ds_d}(s_d))] dr \right) ds. \end{aligned}$$

Then, we have the following formula

$$\begin{aligned} \langle [\Gamma_{\mathbf{T}}]_{i,t}^Z, \omega \rangle &= e^{2\pi i n K t T^{(i)}} \langle [\Gamma_{\mathbf{T}}], \omega \rangle - 2\pi i n K t \int_0^{T^{(i)}} e^{2\pi i n K t s_i} \langle \Gamma_{T,i,s}, \omega \rangle ds_i \\ &+ \int_{U(\mathbf{T})} (\iota_Z \omega \circ [\Gamma_{\mathbf{T}}]_{i,t}^Z)(\mathbf{s}) ds. \end{aligned}$$

Since the action of $\mathbf{P}_{\mathbf{t}}^{d,X}$ for $\mathbf{t} \in \mathbb{R}^d$ is identity on the center Z ,

$$\lim_{t_1, \dots, t_d \rightarrow \infty} e^{-(t_1 + \dots + t_d)/2} \int_{U(\mathbf{T})} (\iota_Z (\mathbf{P}_{\mathbf{t}}^{d,X})^* \omega \circ [\Gamma_{\mathbf{T}}]_{i,t}^Z)(\mathbf{s}) ds = 0.$$

- 1 Thus, in particular concerning $d = 1$, it follows by the definition of Bufetov func-
2 tional (Lemma 3.5), the statement holds. \square

Here we define a restricted vector $\mathbf{T}_{i,s}$ of $\mathbf{T} = (T^{(1)}, \dots, T^{(d)}) \in \mathbb{R}^d$. For fixed $i \in [1, d]$, pick $s_i \in [0, T^{(i)}]$ such that $\mathbf{T}_{i,s} \in \mathbb{R}^d$ is a vector with its coordinates

$$T_{i,s}^{(j)} = \begin{cases} T^{(j)} & \text{if } j \neq i \\ s_i & \text{if } j = i. \end{cases}$$

- 3 Similarly, $\mathbf{T}_{i_1, \dots, i_k, s}$ is a vector whose i_1, \dots, i_k -th coordinates are replaced by
4 s_{i_1}, \dots, s_{i_k} .

Lemma 6.4. *For $m \in M$ and $y_i \in \mathbb{R}$, the following property holds:*

$$\begin{aligned} & \beta_H(\alpha, \phi_{y_i}^{Y_i}(m), \mathbf{T}) = \\ & e^{-2\pi i y_i n K T^{(i)}} \beta_H(\alpha, m, \mathbf{T}) + 2\pi i n K y_i \int_0^{T^{(i)}} e^{-2\pi i y_i n K s_i} \beta_H(\alpha, m, \mathbf{T}_{i,s}) ds_i. \end{aligned}$$

Proof. By definition (4), (66) and commutation relation (1), it follows that

$$\phi_{y_i}^{Y_i}(\Gamma_{\mathbf{T}}^X(m)) = [\Gamma_{\mathbf{T}}^X(\phi_{y_i}^{Y_i}(m))]_{i,t}^Z.$$

By the invariance property of Bufetov functional and Lemma 6.3,

$$\begin{aligned} & \beta_H(\alpha, m, \mathbf{T}) = \hat{\beta}_H(\alpha, \phi_{y_i}^{Y_i}(\Gamma_{\mathbf{T}}^X(m))) \\ &= e^{2\pi i y_i n K T^{(i)}} \hat{\beta}_H(\alpha, \Gamma_{\mathbf{T}}^X(\phi_{y_i}^{Y_i}(m))) - 2\pi i n K y_i \int_0^{T^{(i)}} e^{2\pi i n K y_i s_i} \hat{\beta}_H(\alpha, \Gamma_{T,i,s}^X(\phi_{s_i}^{Y_i}(m))) ds_i \\ &= e^{2\pi i y_i n K T^{(i)}} \beta_H(\alpha, \phi_{y_i}^{Y_i}(m), \mathbf{T}) - 2\pi i n K y_i \int_0^{T^{(i)}} e^{2\pi i n K y_i s_i} \beta_H(\alpha, \phi_{s_i}^{Y_i}(m), \mathbf{T}_{i,s}) ds_i. \end{aligned}$$

- 5 Then the statement follows immediately. \square

1 We extend previous Lemma 6.4 to higher rank actions by induction argument.

Lemma 6.5 (Rank- d action). *For $m \in M$ and $y = (y_1, \dots, y_d) \in \mathbb{R}^d$, the following identity for the cocycle β_H holds:*

$$(67) \quad \begin{aligned} \beta_H(\alpha, Q_y^{d,Y}(m), \mathbf{T}) &= e^{-2\pi i \sum_{j=1}^d y_j n K T^{(j)}} \beta_H(\alpha, m, \mathbf{T}) \\ &+ \sum_{k=1}^d \sum_{1 \leq i_1 < \dots < i_k \leq d} \prod_{j=1}^k (2\pi i n K y_{i_j}) e^{-2\pi i n K (\sum_{l \notin \{i_1, \dots, i_k\}} y_l T^{(l)})} \\ &\times \int_0^{T^{(i_1)}} \dots \int_0^{T^{(i_k)}} e^{-2\pi i n K (y_{i_1} s_{i_1} + \dots + y_{i_k} s_{i_k})} \beta_H(\alpha, m, \mathbf{T}_{i_1, \dots, i_k, s}) ds_{i_k} \dots ds_{i_1}. \end{aligned}$$

Proof. We verified that the statement works for $d = 1$ in Lemma 6.4. Assume that (67) holds for rank $d - 1$ action $Q_{y'}^{d-1,Y}$ by induction hypothesis. For convenience, we write

$$Q_y^{d,Y}(m) = \phi_{y_d}^{Y_d} \circ Q_{y'}^{d-1,Y}(m) \text{ for } y' \in \mathbb{R}^{d-1} \text{ and } y = (y', y_d) \in \mathbb{R}^d.$$

By applying Lemma 6.4,

$$(68) \quad \begin{aligned} \beta_H(\alpha, Q_y^{d,Y}(m), \mathbf{T}) &= e^{-2\pi i y_d n K T^{(d)}} \beta_H(\alpha, Q_{y'}^{d-1,Y}(m), \mathbf{T}) \\ &+ 2\pi i n K y_d \int_0^{T^{(d)}} e^{-2\pi i y_d n K s_d} \beta_H(\alpha, Q_{y'}^{d-1,Y}(m), \mathbf{T}_{d,s}) ds_d \\ &:= I + II. \end{aligned}$$

Firstly, by induction hypothesis,

$$\begin{aligned} I &= e^{-2\pi i y_d n K T^{(d)}} \left(e^{-2\pi i \sum_{j=1}^{d-1} y_j n K T^{(j)}} \beta_H(\alpha, m, \mathbf{T}) \right. \\ &+ \sum_{k=1}^{d-1} \sum_{1 \leq i_1 < \dots < i_k \leq d-1} \prod_{j=1}^k (2\pi i n K y_{i_j}) e^{-2\pi i n K (\sum_{l \notin \{i_1, \dots, i_k\}} y_l T^{(l)} + y_d T^{(d)})} \\ &\times \int_0^{T^{(i_1)}} \dots \int_0^{T^{(i_k)}} e^{-2\pi i n K (y_{i_1} s_{i_1} + \dots + y_{i_k} s_{i_k})} \beta_H(\alpha, m, \mathbf{T}_{i_1, \dots, i_k, s}) ds_{i_k} \dots ds_{i_1} \Big) \\ &= e^{-2\pi i \sum_{j=1}^d y_j n K T^{(j)}} \beta_H(\alpha, m, \mathbf{T}) + III. \end{aligned}$$

2 Then term III contains 0 to $d - 1$ th iterated integrals on the restricted rectangles
3 (from 0 to $d - 1$ -th) containing $e^{-2\pi i n K y_d T^{(d)}}$ outside of iterated integrals.

4 For the second part, we apply induction hypothesis again for restricted rectangle
5 $\mathbf{T}_{d,s}$. Then,

$$\begin{aligned}
II &= 2\pi\iota n K y_d \int_0^{T^{(d)}} e^{-2\pi\iota y_d n K s_d} \left[e^{-2\pi\iota \sum_{j=1}^{d-1} y_j n K T^{(j)}} \beta_H(\alpha, m, \mathbf{T}_{d,s}) \right] ds_d \\
&+ \sum_{k=1}^{d-1} \sum_{1 \leq i_1 < \dots < i_k \leq d-1} (2\pi\iota n K y_d) \prod_{j=1}^k (2\pi\iota n K y_{i_j}) e^{-2\pi\iota n K (\sum_{l \notin \{i_1, \dots, i_k\}} y_l T^{(l)})} \\
&\times \int_0^{T^{(d)}} \left(\int_0^{T^{(i_1)}} \dots \int_0^{T^{(i_k)}} ds_{i_k} \dots ds_{i_1} \right) ds_d \\
&\times e^{-2\pi\iota n K (y_{i_1} s_{i_1} + \dots + y_{i_k} s_{i_k} + y_d s_d)} \beta_H(\alpha, m, \mathbf{T}_{i_1, \dots, i_k, d, s}).
\end{aligned}$$

1 The term II consist of 1 to d -th iterated integrals on the (from 1 to d -th) restricted
2 rectangles containing $e^{-2\pi\iota n K y_d s_d}$ inside of iterated integrals. Thus, by rearranging
3 terms II and III , we obtain all the terms in the expression (67). \square

4 **6.2. Extensions of domain.** In this subsection, the domain of functionals defined
5 on standard rectangle $\Gamma_{\mathbf{T}}^X$ extends to the class \mathfrak{R} (see Definition 1.1). Furthermore,
6 the functional associated with the analytic norm extends to holomorphic function
7 on a complex domain.

8 *Proof of Corollary 1.4.* Firstly, we can extend our functional to class $(Q_y^{d,Y})_* \Gamma_{\mathbf{T}}^X$ for
9 any $y \in \mathbb{R}^d$ by invariance property (Lemma 3.6). Similarly, by Lemma 6.3, Bufetov
10 functional defined on the standard rectangles extends to the class of generalized
11 rectangle $(\phi_{t,z}^Z) \circ P_{\mathbf{t}}^{d,\alpha}(m)$. Since the flow generated by Z commutes with other
12 actions P and Q , for any standard rectangle $\Gamma = \Gamma(m)$ with a fixed point $m \in M$,
13 we have $(\phi_z^Z)_* \Gamma(m) = \Gamma(\phi_z^Z(m))$ for any $z \in \mathbb{R}$. Therefore, by combining with the
14 invariance under the action Q from Lemma 3.6, the domain of Bufetov functional
15 extends to the class \mathfrak{R} . \square

For any $R > 0$, the *analytic norm* defined for all $\mathbf{c} \in \ell^2$ as

$$\|\mathbf{c}\|_{\omega,R} = \sum_{n \neq 0} \sum_{i=1}^{\mu(n)} e^{nR} |c_{i,n}|.$$

16 Let Ω_R denote the subspace of $\mathbf{c} \in \ell^2$ such that $\|\mathbf{c}\|_{\omega,R}$ is finite.

Lemma 6.6. *For $\mathbf{c} \in \Omega_R$ and $\mathbf{T} \in \mathbb{R}_+^d$, the function*

$$\beta_{\mathbf{c}}(\alpha, Q_y^{d,Y} \circ \phi_z^Z(m), \mathbf{T}), \quad (y, z) \in \mathbb{R}^d \times \mathbb{T}$$

17 *extends to a holomorphic function in the domain*

$$(69) \quad D_{R,T} := \{(y, z) \in \mathbb{C}^d \times \mathbb{C}/\mathbb{Z} \mid \sum_{i=1}^d |\operatorname{Im}(y_i)| T^{(i)} + |\operatorname{Im}(z)| < \frac{R}{2\pi K}\}.$$

The following bound holds: for any $R' < R$ there exists a constant $C > 0$ such that, for all $(y, z) \in D_{R',T}$ we have

$$\begin{aligned}
&|\beta_{\mathbf{c}}(\alpha, Q_y^{d,Y} \circ \phi_z^Z(m), \mathbf{T})| \\
&\leq C_{R,R'} \|\mathbf{c}\|_{\omega,R} (L + \operatorname{vol}(U(\mathbf{T}))^{1/2} (1 + E_M(a, \mathbf{T})) (1 + K \sum_{i=1}^d |\operatorname{Im}(y_i)| T^{(i)}).
\end{aligned}$$

Proof. By Lemma 6.5 and (64),

$$\begin{aligned} \beta_{\mathbf{c}}(\alpha, \mathbf{Q}_y^{d,Y} \circ \phi_z^Z(m), \mathbf{T}) &= e^{(z-2\pi i \sum_{j=1}^d y_j n K T^{(j)})} \beta_H(\alpha, m, \mathbf{T}) \\ &+ \sum_{k=1}^d \sum_{1 \leq i_1 < \dots < i_k \leq d} \prod_{j=1}^k (2\pi i n K y_{i_j}) e^{-2\pi i n K (\sum_{l \notin \{i_1, \dots, i_k\}} y_l T^{(l)})} \\ &\times e^{2\pi i n K z} \int_0^{T^{(i_1)}} \dots \int_0^{T^{(i_k)}} e^{-2\pi i n K (y_{i_1} s_{i_1} + \dots + y_{i_k} s_{i_k})} \beta_H(\alpha, m, \mathbf{T}_{i_1, \dots, i_k, s}) ds_{i_k} \dots ds_{i_1}. \end{aligned}$$

As a consequence, by Lemma 3.7 for each variable $(y_i, z) \in \mathbb{C} \times \mathbb{C}/\mathbb{Z}$, Then for the rank d -action, by induction, for $(y, z) \in \mathbb{C}^d \times \mathbb{C}/\mathbb{Z}$ we have

$$\begin{aligned} &|\beta_{\mathbf{c}}(\alpha, \mathbf{Q}_y^{d,Y} \circ \phi_z^Z(x), \mathbf{T})| \\ &\leq (L + \text{vol}(U(\mathbf{T}))^{1/2} (1 + E_M(a, \mathbf{T}))) \left(C_1 \sum_{n \neq 0} \sum_{i=1}^{\mu(n)} e^{nR} |c_{i,n}| e^{2\pi |Im(z - \sum_{i=1}^d T^{(i)} y_i)| nK} \right. \\ &+ \sum_{k=1}^d C_k \left(\sum_{1 \leq i_1 < \dots < i_k \leq d} \prod_{j=1}^k (|Im(y_{i_j})| T^{(i_j)}) \right. \\ &\times \left. \sum_{n \neq 0} \sum_{i=1}^{\mu(n)} n |c_{i,n}| e^{2\pi (|Im(z)| + \sum_{j=1}^k T^{(i_j)} |Im(y_{i_j})|) nK} \right) \Big). \end{aligned}$$

- 1 Therefore, the functional $\beta_{\mathbf{c}}(\alpha, \mathbf{Q}_y^{d,Y} \circ \phi_z^Z(m), \mathbf{T})$ is bounded by a series of holo-
 2 morphic functions on $\mathbb{C}^d \times \mathbb{C}/\mathbb{Z}$ and it converges uniformly on compact subsets of
 3 domain $D_{R,T}$. Thus it is holomorphic on the set. \square

4 7. MEASURE ESTIMATION FOR BOUNDED-TYPE

5 In this section, we prove a measure estimation of Bufetov functional with bounded-
 6 type α . This result is a generalization of §11 in [FK20b].

7 Let \mathcal{O}_r denote the space of holomorphic functions on the ball $B_{\mathbb{C}}(0, r) \subset \mathbb{C}^n$.

Theorem 7.1. [Bru99, Theorem 1.9] For any $f \in \mathcal{O}_r$, there is a constant $d := d_f(r) > 0$ such that for any convex set $D \subset B_{\mathbb{R}}(0, 1) := B_{\mathbb{C}}(0, 1) \cap \mathbb{R}^n$, for any measurable subset $U \subset D$

$$\sup_D |f| \leq \left(\frac{4n \text{Leb}(D)}{\text{Leb}(U)} \right)^d \sup_U |f|.$$

8 We say that a holomorphic function f defined in a disk is p -valent if it assumes
 9 no value more than p -times there. We also say that f is θ -valent if it is a constant.

Definition 7.2. [Bru99, Def 1.6] Let \mathcal{L}_t denote the set of one-dimensional complex affine spaces $L \subset \mathbb{C}^n$ such that $L \cap B_{\mathbb{C}}(0, t) \neq \emptyset$. For $f \in \mathcal{O}_r$, the number

$$\nu_f(t) := \sup_{L \in \mathcal{L}_t} \{\text{valency of } f \mid L \cap B_{\mathbb{C}}(0, t) \neq \emptyset\}$$

10 is called the *valency* of f in $B_{\mathbb{C}}(0, t)$.

11 By Proposition 1.7 of [Bru99], for any $f \in \mathcal{O}_r$ with finite valency $\nu_f(t)$ for any
 12 $t \in [1, r)$, there is a constant $c := c(r) > 0$ such that

$$(70) \quad d_f(r) \leq c \nu_f\left(\frac{1+r}{2}\right).$$

Lemma 7.3. *[FK20b, Lemma 10.3] Let $R > r > 1$. For any normal family $\mathcal{F} \subset \mathcal{O}_R$, assume that no functions in $\overline{\mathcal{F}} = \emptyset$ are constant along a one-dimensional complex line. Then we have*

$$\sup_{f \in \mathcal{F}} \nu_f(r) < \infty.$$

Lemma 7.4. *Let $L > 0$ and $\mathcal{B} \subset DC(L)$ be a bounded subset. Given $R > 0$, for all $\mathbf{c} \in \Omega_R$ and all $\mathbf{T}^{(i)} > 0$, denote $\mathcal{F}(\mathbf{c}, \mathbf{T})$ by the family of real analytic functions of the variable $y \in [0, 1]^d$ and*

$$\mathcal{F}(\mathbf{c}, \mathbf{T}) := \{\beta_{\mathbf{c}}(\alpha, \Phi_{\alpha, x}(\xi_{y, z}), \mathbf{T}) \mid (\alpha, x, z) \in \mathcal{B} \times M \times \mathbb{T}\}.$$

Then there exist $\mathbf{T}_{\mathcal{B}} := (\mathbf{T}_{\mathcal{B}}^{(i)})$ and $\rho_{\mathcal{B}} > 0$, such that for every (R, \mathbf{T}) with $R/\mathbf{T}^{(i)} \geq \rho_{\mathcal{B}}$, $\mathbf{T}^{(i)} \geq \mathbf{T}_{\mathcal{B}}^{(i)}$ and for all $\mathbf{c} \in \Omega_R \setminus \{0\}$, we have

$$(71) \quad \sup_{f \in \mathcal{F}(\mathbf{c}, \mathbf{T})} \nu_f < \infty.$$

Proof. Since $\mathcal{B} \subset \mathfrak{M}$ is bounded, for each time $t_i \in \mathbb{R}$ and $1 \leq i \leq g$,

$$0 < t_{\mathcal{B}}^{\min} = \min_i \inf_{\alpha \in \mathcal{B}} t_{\text{Ret}, i, \alpha} \leq \max_i \sup_{\alpha \in \mathcal{B}} t_{\text{Ret}, i, \alpha} = t_{\mathcal{B}}^{\max} < \infty.$$

For any $\alpha \in \mathcal{B}$ and $x \in M$, the map $\Phi_{\alpha, x} : [0, 1]^d \times \mathbb{T} \rightarrow \prod_{i=1}^d [0, t_{\alpha, i}] \times \mathbb{T}$ in (53) extends to a complex analytic diffeomorphism $\hat{\Phi}_{\alpha, x} : \mathbb{C}^d \times \mathbb{C}/\mathbb{Z} \rightarrow \mathbb{C}^d \times \mathbb{C}/\mathbb{Z}$. By Lemma 6.6, it follows that for fixed $z \in \mathbb{T}$, real analytic function $\beta_{\mathbf{c}}(\alpha, \Phi_{\alpha, x}(\xi_{y, z}), \mathbf{T})$ extends to a holomorphic function defined on a region

$$H_{\alpha, m, R, t} := \{y \in \mathbb{C}^d \mid \sum_{i=1}^d |\text{Im}(y_i)| \leq h_{\alpha, m, R, t}\}.$$

By boundedness of the set $\mathcal{B} \subset \mathfrak{M}$, it follows that

$$\inf_{(\alpha, x) \in \mathcal{B} \times M} h_{\alpha, m, R, t} := h_{R, T} > 0.$$

We remark that the function $h_{\alpha, m, R, t}$ and its lower bound $h_{R, T}$ can be obtained from the formula (53) for the polynomial $\Phi_{\alpha, x}$ and the definition of the domain $D_{R, T}$ in the formula (69).

For every $r > 1$, there exists $\rho_{\mathcal{B}} > 1$ such that for every (R, \mathbf{T}) with $R/\mathbf{T}^{(i)} > \rho_{\mathcal{B}}$,

$$(72) \quad \beta_{\mathbf{c}}(\alpha, \Phi_{\alpha, x}(\xi_{y, z}), \mathbf{T}) \in \mathcal{O}_r$$

as a function of $y \in \mathbb{T}^d$.

By Lemma 6.6, the family $\mathcal{F}(\mathbf{c}, \mathbf{T})$ is uniformly bounded and normal. By Lemma 5.4 for the non-zero L^2 -lower bound of functionals, for sufficiently large pair \mathbf{T} , no sequence from $\mathcal{F}(\mathbf{c}, \mathbf{T})$ can converge to a constant. Therefore, by Lemma 7.3 for the family $\mathcal{F} = \mathcal{F}(\mathbf{c}, \mathbf{T})$, the main statement follows. \square

We derive measure estimates of Bufetov functionals on the rectangular domain.

Lemma 7.5. *Let $\alpha \in DC$ such that the forward orbit of \mathbb{R}^d -action $\{r_t[\alpha]\}_{t \in \mathbb{R}_+^d}$ is contained in a compact set of \mathfrak{M}_g . There exist $R, C, \delta > 0$ and $T_0 \in \mathbb{R}_+^d$ such that, for every $\mathbf{c} \in \Omega_R \setminus \{0\}$, $T \geq T_0$ and for every $\epsilon > 0$, we have*

$$\text{vol}(\{m \in M \mid |\beta_{\mathbf{c}}(\alpha, m, \mathbf{T})| \leq \epsilon \text{vol}(U(\mathbf{T}))^{1/2}\}) \leq C\epsilon^\delta.$$

Proof. Since $\alpha \in DC$ and the orbit $\{r_t[\alpha]\}_{t \in \mathbb{R}_+}$ is contained in a compact set, there exists $L > 0$ such that $r_t[\alpha] \in DC(L)$ for all $t \in \mathbb{R}_+^d$. Then, we choose $\mathbf{T}_0 \in \mathbb{R}^d$ and $R > 0$ from the conclusion of Lemma 7.4. By the scaling property,

$$\beta_{\mathbf{c}}(\alpha, m, \mathbf{T}) = \left(\frac{\text{vol}(U(\mathbf{T}))}{\text{vol}(U(\mathbf{T}_0))} \right)^{1/2} \beta_{\mathbf{c}}(g_{\log(\mathbf{T}/\mathbf{T}_0)}[\alpha], m, \mathbf{T}_0).$$

By Fubini's theorem, it suffices to estimate

$$\text{Leb}(\{y \in [0, 1]^d \mid |\beta_{\mathbf{c}}(\alpha, \Phi_{\alpha, x}(\xi_{y, z}), \mathbf{T}_0)| \leq \epsilon\}).$$

Let $\delta^{-1} := c(r) \sup_{f \in \mathcal{F}(\mathbf{c}, \mathbf{T}_0)} \nu_f(\frac{1+r}{2}) < \infty$ as in (70) and (71). By Lemma 5.4, we have

$$\inf_{(\alpha, x, z) \in \mathcal{B} \times M \times \mathbb{T}} \sup_{y \in [0, 1]^d} |\beta_{\mathbf{c}}(\alpha, \Phi_{\alpha, x}(\xi_{y, z}), \mathbf{T}_0)| > 0$$

so that the functional is not trivial. By Theorem 7.1 for the unit ball $D = B_{\mathbb{R}}(0, 1)$ and setting

$$U = \{y \in [0, 1]^d \mid |\beta_{\mathbf{c}}(\alpha, \Phi_{\alpha, x}(\xi_{y, z}), \mathbf{T}_0)| \leq \epsilon\},$$

by the bound in (70) for $d_f(r)$, there exists a constant $C > 0$ such that for all $\epsilon > 0$ and $(\alpha, x, z) \in \mathcal{B} \times M \times \mathbb{T}$,

$$\text{Leb}(\{y \in [0, 1]^d \mid |\beta_{\mathbf{c}}(\alpha, \Phi_{\alpha, x}(\xi_{y, z}), \mathbf{T}_0)| \leq \epsilon\}) \leq C\epsilon^\delta.$$

1 Then the statement follows from the Fubini theorem. \square

Corollary 7.6. *Let α be as in the previous Lemma 7.5. There exist $R, C, \delta > 0$ and $\mathbf{T}_0 \in \mathbb{R}_+^d$ such that, for every $\mathbf{c} \in \Omega_R \setminus \{0\}$, $\mathbf{T} \geq \mathbf{T}_0$ and for every $\epsilon > 0$, we have*

$$\text{vol}\left(\{m \in M \mid |\langle \mathcal{P}_{U(\mathbf{T})}^{d, \alpha} m, \omega_{\mathbf{c}} \rangle| \leq \epsilon \text{vol}(U(\mathbf{T}))^{1/2}\}\right) \leq C\epsilon^\delta.$$

2 **Acknowledgement.** The author deeply appreciates Giovanni Forni for valuable
3 suggestions to improve the draft. He acknowledges Rodrigo Treviño and Corinna
4 Ulcigrai for giving several comments. He is also grateful to Oliver Butterley, Jacky
5 Jia Chong, Krzysztof Frączek, Osama Khail, and Lucia D. Simonelli for fruitful dis-
6 cussions. This work was initiated when the author visited the Institut de Mathema-
7 tiques de Jussieu-Paris Rive Gauche in Paris, France. He acknowledges invitation
8 and hospitality during the visit. Lastly, the author is thankful to the referee for
9 helpful comments and suggestions for improvement in the presentation of this work.
10 This research was partially supported by the NSF grant DMS 1600687 and by the
11 Centre of Excellence “Dynamics, mathematical analysis and artificial intelligence”
12 at Nicolaus Copernicus University in Toruń.

REFERENCES

- 14 [AB18] Alexander Adam and Viviane Baladi. *Horocycle averages on closed manifolds and transfer*
15 *operators*. arXiv preprint arXiv:1809.04062 (2018).
16 [AFRU21] Artur Avila, Giovanni Forni, Davide Ravotti, and Corinna Ulcigrai, *Mixing for smooth*
17 *time-changes of general nilflows*, Advances in Mathematics **385** (2021), 107759.
18 [AFU11] Artur Avila, Giovanni Forni, and Corinna Ulcigrai, *Mixing for the time-changes of*
19 *heisenberg nilflows*, Journal of Differential Geometry **89** (2011), no. 3, 369–410.
20 [B19] Viviane Baladi. *There are no deviations for the ergodic averages of Giulietti–Liverani horo-*
21 *cycle flows on the two-torus*. Ergodic Theory and Dynamical Systems (2019): 1–14.
22 [Bru99] Alexander Brudnyi, *On local behavior of analytic functions*, Journal of Functional Anal-
23 *ysis* **2** (1999), no. 169, 481–493.

- 1 [Buf10] Alexander I Bufetov, *Hölder cocycles and ergodic integrals for translation flows on flat*
2 *surfaces*, Electronic Research Announcements **17** (2010), 34–42.
- 3 [Buf13] ———, *Limit theorems for suspension flows over Vershik automorphisms*. Russian Math-
4 ematical Surveys **68.5** (2013): 789.
- 5 [Buf14a] ———, *Finitely-additive measures on the asymptotic foliations of a markov compactum*,
6 Moscow Mathematical Journal **14** (2014), no. 2, 205–224.
- 7 [Buf14b] ———, *Limit theorems for translation flows*, Ann. of Math. (2) **179** (2014), no. 2,
8 431–499. MR 3152940
- 9 [BF14] Alexander Bufetov and Giovanni Forni, *Limit theorems for horocycle flows*, Ann. Sci. Éc.
10 Norm. Supér. (4) **47** (2014), no. 5, 851–903. MR 3294619
- 11 [BS13] Alexander I Bufetov and Boris Solomyak. *Limit theorems for self-similar tilings*. Commu-
12 nications in Mathematical Physics **319.3** (2013): 761–789.
- 13 [BS20] Oliver Butterley and Lucia D Simonelli, *Parabolic flows renormalized by partially hyper-*
14 *bolic maps*, Bollettino dell’Unione Matematica Italiana (2020), 1–20.
- 15 [CF15] Salvatore Cosentino and Livio Flaminio, *Equidistribution for higher-rank Abelian actions*
16 *on Heisenberg nilmanifolds*, J. Mod. Dyn. **9** (2015), 305–353. MR 3426826
- 17 [CM16] Francesco Cellarosi and Jens Marklof, *Quadratic weyl sums, automorphic functions and*
18 *invariance principles*, Proceedings of the London Mathematical Society **113** (2016), no. 6,
19 775–828.
- 20 [DS17] Dmitry Dolgopyat and Omri Sarig. *Temporal distributional limit theorems for dynamical*
21 *systems*. Journal of Statistical Physics 166.3-4 (2017): 680–713.
- 22 [FT15] Frédéric Faure and Masato Tsujii. *Prequantum transfer operator for symplectic Anosov*
23 *diffeomorphism*. Asterisque 375 (2015).
- 24 [FGL19] Frédéric Faure, Sébastien Gouëzel, and Erwan Lanneau, *Ruelle spectrum of linear*
25 *pseudo-anosov maps*, Journal de l’École polytechnique—Mathématiques **6** (2019), 811–877.
- 26 [FF06] ———, *Equidistribution of nilflows and applications to theta sums*, Ergodic Theory Dy-
27 nam. Systems **26** (2006), no. 2, 409–433. MR 2218767
- 28 [FF14] ———, *On effective equidistribution for higher step nilflows*, arXiv preprint
29 arXiv:1407.3640 (2014).
- 30 [FFT16] Livio Flaminio, Giovanni Forni, and James Tanis, *Effective equidistribution of twisted*
31 *horocycle flows and horocycle maps*, Geom. Funct. Anal. **26** (2016), no. 5, 1359–1448.
32 MR 3568034
- 33 [For02] Giovanni Forni, *Deviation of ergodic averages for area-preserving flows on surfaces of*
34 *higher genus*, Annals of Mathematics **155** (2002), no. 1, 1–103.
- 35 [For20a] ———, *On the equidistribution of unstable curves for pseudo-anosov diffeomorphisms*
36 *of compact surfaces*, arXiv preprint arXiv:2007.03144 (2020).
- 37 [For20b] ———, *Ruelle resonances from cohomological equations*, preprint arXiv:2007.03116
38 (2020).
- 39 [FK20a] Giovanni Forni and Adam Kanigowski, *Multiple mixing and disjointness for time changes*
40 *of bounded-type heisenberg nilflows*, Journal de l’École polytechnique—Mathématiques **7**
41 (2020), 63–91.
- 42 [FK20b] ———, *Time-changes of heisenberg nilflows*, Asterisque **416** (2020), 253–299.
- 43 [GG04] Friedrich Götze and Mikhail Gordin, *Limiting distributions of theta series on siegel half-*
44 *spaces*, St. Petersburg Mathematical Journal **15** (2004), no. 1, 81–102.
- 45 [GM14] Jory Griffin and Jens Marklof, *Limit theorems for skew translations.*, Journal of Modern
46 Dynamics **8** (2014), no. 2.
- 47 [GS14] Alexander Gorodnik and Ralf Spatzier, *Exponential mixing of nilmanifold automorphisms*,
48 Journal d’Analyse Mathématique **123** (2014), no. 1, 355–396.
- 49 [GS15] ———, *Mixing properties of commuting nilmanifold automorphisms*, Acta Mathematica
50 **215** (2015), no. 1, 127–159.
- 51 [Ka03] Katok, Anatolij Borisovič. Combinatorial constructions in ergodic theory and dynamics.
52 No. 30. American Mathematical Soc., 2003.
- 53 [Kim21] Minsung Kim, *Effective equidistribution for generalized higher step nilflows*, Ergodic
54 Theory and Dynamical Systems (2021), 1–60.
- 55 [KK95] Anatole Katok and Svetlana Katok, *Higher cohomology for abelian groups of toral auto-*
56 *morphisms*, Ergodic Theory and Dynamical Systems **15** (1995), no. 3, 569–592.
- 57 [KM99] D. Y. Kleinbock and G. A. Margulis, *Logarithm laws for flows on homogeneous spaces*,
58 Invent. Math. **138** (1999), no. 3, 451–494. MR 1719827

- 1 [Mar99] Jens Marklof, *Limit theorems for theta sums*, Duke mathematical journal **97** (1999),
2 no. 1, 127–154.
- 3 [MM07] David Mumford and C Musili, *Tata lectures on theta. i (modern birkhäuser classics)*,
4 Birkhäuser Boston Incorporated, 2007.
- 5 [MMY10] Stefano Marmi, Pierre Moussa, and J-C. Yoccoz. *Affine interval exchange maps with a*
6 *wandering interval*. Proceedings of the London Mathematical Society 100.3 (2010): 639-669.
- 7 [MNN07] David Mumford, Madhav Nori, and Peter Norman, *Tata lectures on theta iii*, vol. 43,
8 Springer, 2007.
- 9 [MUY20] Stefano Marmi, Corinna Ulcigrai, and Jean-Christophe Yoccoz. *On Roth type condi-*
10 *tions, duality and central Birkhoff sums for iem*. Asterisque **416** (2020): 65-132.
- 11 [GL19] Paolo Giulietti and Carlangelo Liverani, *Parabolic dynamics and anisotropic Banach*
12 *spaces*, Journal of the European Mathematical Society (2019).
- 13 [Rav18] Davide Ravotti, *Mixing for suspension flows over skew-translations and time-changes of*
14 *quasi-abelian filiform nilflows*, Ergodic Theory and Dynamical Systems (2018), 1–30.
- 15 [Rav20] ———, *Quantitative equidistribution of horocycle push-forwards of transverse arcs*,
16 L’Enseignement Mathématique **66.1** (2020), 135-150.
- 17 [Rav21] D. Ravotti. *Asymptotics and limit theorems for horocycle ergodic integrals a la Ratner*.
18 arXiv preprint arXiv:2107.02090 (2021).
- 19 [S09a] Shah, Nimish A. *Limiting distributions of curves under geodesic flow on hyperbolic mani-*
20 *folds*. Duke Mathematical Journal 148.2 (2009): 251-279.
- 21 [S09b] ———, *Asymptotic evolution of smooth curves under geodesic flow on hyperbolic mani-*
22 *folds*. Duke Mathematical Journal 148.2 (2009): 281-304.
- 23 [Sul82] Dennis Sullivan, *Disjoint spheres, approximation by imaginary quadratic numbers, and*
24 *the logarithm law for geodesics*, Acta Math. **149** (1982), no. 3-4, 215–237. MR 688349
- 25 [Tol78] R. Tolimieri, *Heisenberg manifolds and theta functions*, Transactions of the American
26 Mathematical Society **239** (1978), 293–319.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, MD 20742, USA

27 *Current address:* FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, NICOLAUS COPER-
NICUS UNIVERSITY, CHOPINA 12/18, 87-100 TORUŃ, POLAND
28 *Email address:* mkim16@mat.umk.pl