Unified pipeline: implemented boundaries and current evidence

One raw-data registry, one participant fold plan, shared train/apply ComBat.
Ridge and GINE use identical paired observations for DWI, fMRI and fusion.

Implemented

- Training-only edge QC, harmonization and normalization
- Inner model / sparsity selection and independent GINE stopping
- Inner-OOF reference calibration; global and cohort-aware cBAG
- Original outer checkpoints, private embeddings, gradient contributions
- Runtime audits, resumable tasks and source snapshots

Run classification: development evaluation — external confirmation still required

Outer / inner / stopping boundaries

Raw DWI log1p / signed fMRI Fisher-z → fixed participant outer split

OUTER TRAINING CN
Inner train / validation split
↓ GINE only: split inner training into fit / stopping
Fit preprocessing on fit only; choose epoch on stopping
↓ Refit preprocessing/model on all inner training
Evaluate inner validation; choose configuration
↓ Calibrate using inner OOF predictions
Refit outer model for inner-selected epoch budget

OUTER HELD-OUT PARTICIPANTS
Frozen preprocessing → raw age prediction
↓ Compute BAG using actual age
↓ Apply frozen training-only calibration

No test age in feature transformations.
No test fold in hyperparameters / sparsity / epochs.
No clinical, genetic or metabolic predictor features.

Harmonization decision follows age-preservation evidence

Primary development candidate: training-age mean-only ComBat
Held-out transformation: x_harmonized = x − training residual SD × batch offset

Training age helps separate age and acquisition offsets; held-out age is never required.
No-ComBat remains a prespecified control. Batch-only location/scale remains a diagnostic sensitivity option.
Prior modality-wide MAE: DWI none 6.02 / mean-only 6.11; fMRI none 7.43 / mean-only 7.49.
Those historical development comparisons are not the new matched six-model benchmark.

Conservative population: unresolved pairs are not averaged

Population	Scans	Participants	CN scans
DWI	919	670	723
fMRI	575	447	441
paired	318	252	237

Pairing: 318 retained; 250 age_conflict; 7 insufficient_paired_CN_support.
Same-key / age / sex / diagnosis agreement is checked; acquisition/clinical dates still require verification.

GINE comparisons and scientific outputs

Capacity: 1 × 16, 2 × 32, current 3 × 64; 2 × 32 residual ablation.
Graph sparsity is selected inside inner validation. Signed fMRI edges are retained; fusion uses modality-union topology and concatenated node profiles.

Learning curves: evaluation-mode training / stopping loss and MAE; selected epochs recorded.
Separate runs vary training participant fractions with fixed outer test folds.

Bias correction: inspect global and cohort-aware OOF cBAG versus age within every cohort.

Interpretation: original outer checkpoints, fold-specific embeddings and gradient-times-input.
Anatomical naming requires verified ROI order.

Results: completion status is part of the result

Mode	Input	Model	Completed folds	Mean fold MAE
age_mean	DWI	gine	5	7.846
age_mean	fMRI	gine	5	9.388
age_mean	fusion	gine	5	8.123
none	DWI	gine	5	7.495
none	fMRI	gine	5	9.441
none	fusion	gine	5	8.054

Status: complete. Mean-fold values are descriptive; use the report for pooled OOF metrics and cohort breakdowns.

Remaining data validation and clinical analysis gates

Verify HABS diagnosis provenance, ADRC visit links/protocols, age discrepancies and ROI order.

Clinical inventory: CN/MCI/AD, APOE / APOE4, sex, T2D, HbA1c and continuous age. Source codes, units and clinical timing must be verified before testing.

Clinical analyses: group distributions and complete-case multivariable associations with cohort, age and sex; participant-cluster uncertainty; explicit exploratory multiplicity correction.

Embeddings: train-fit PCA per outer model, held-out overlays; never pool unaligned fold coordinates.

No claim of fully removed scanner effects, preserved biology, clinical biomarkers or unseen-site generalization.