

# MAT426: Advanced Calculus

**Miraj Samarakkody**

Tougaloo College

04/01/2025

## 2.41 Theorem

### Theorem

If a set  $E$  in  $\mathbb{R}^k$  has one of the following three properties, then it has the other two:

- (a)  $E$  is closed and bounded.
- (b)  $E$  is compact.
- (c) Every infinite subset of  $E$  has a limit point in  $E$ .

Proof:

## 2.42 Theorem

### Theorem - Weierstrass

Every bounded infinite subset of  $\mathbb{R}^k$  has a limit point in  $\mathbb{R}^k$ .

# Perfect Sets

## 2.43 Theorem

Let  $P$  be a non-empty perfect set in  $\mathbb{R}^k$ . Then  $P$  is uncountable.

# Corollary

## Corollary

Every interval  $[a, b]$  is uncountable. In particular, the set of all real numbers is uncountable.

# The Cantor Set

Cantor set is a perfect set in  $\mathbb{R}^1$  which contain no segment.

# The Cantor Set

Cantor set is a perfect set in  $\mathbb{R}^1$  which contain no segment.

Let  $E_0$  be the interval  $[0, 1]$ .



# The Cantor Set

Cantor set is a perfect set in  $\mathbb{R}^1$  which contain no segment.

Let  $E_0$  be the interval  $[0, 1]$ .

Remove the segment  $\left(\frac{1}{3}, \frac{2}{3}\right)$ , and let  $E_1$  be the union of the intervals

$$\left[0, \frac{1}{3}\right], \left[\frac{2}{3}, 1\right]$$

# The Cantor Set

Cantor set is a perfect set in  $\mathbb{R}^1$  which contain no segment.

Let  $E_0$  be the interval  $[0, 1]$ .

Remove the segment  $\left(\frac{1}{3}, \frac{2}{3}\right)$ , and let  $E_1$  be the union of the intervals

$$\left[0, \frac{1}{3}\right], \left[\frac{2}{3}, 1\right]$$

Remove the middle thirds of these intervals, and let  $E_2$  be the union of the intervals

$$\left[0, \frac{1}{9}\right], \left[\frac{2}{9}, \frac{3}{9}\right], \left[\frac{6}{9}, \frac{7}{9}\right], \left[\frac{8}{9}, 1\right]$$

# The Cantor set

Continuing in this way, we obtain a sequence of compact sets  $E_n$ , such that

1.  $E_1 \supset E_2 \supset E_3 \supset \dots$
2.  $E_n$  is the union of  $2^n$  intervals, each of length  $3^{-n}$ .

The set

$$P = \bigcap_{n=1}^{\infty} E_n$$

is called the **Cantor set**.

# Connected Sets

# Connected Sets

## 2.45 Definition

Two subsets  $A$  and  $B$  of a metric space  $X$  are said to be separated if both  $A \cap \overline{B}$  and  $\overline{A} \cap B$  are empty.

# Connected Sets

## 2.45 Definition

Two subsets  $A$  and  $B$  of a metric space  $X$  are said to be separated if both  $A \cap \overline{B}$  and  $\overline{A} \cap B$  are empty.

A set  $E \subset X$  is said to be connected if  $E$  is not a union of two non-empty separated sets.

# Connected Sets

## 2.45 Definition

Two subsets  $A$  and  $B$  of a metric space  $X$  are said to be separated if both  $A \cap \overline{B}$  and  $\overline{A} \cap B$  are empty.

A set  $E \subset X$  is said to be connected if  $E$  is not a union of two non-empty separated sets.

Separated sets are disjoint, but disjoint sets need not to be separated.

# Connected Sets

## 2.45 Definition

Two subsets  $A$  and  $B$  of a metric space  $X$  are said to be separated if both  $A \cap \overline{B}$  and  $\overline{A} \cap B$  are empty.

A set  $E \subset X$  is said to be connected if  $E$  is not a union of two non-empty separated sets.

Separated sets are disjoint, but disjoint sets need not to be separated.

## Example

The segment  $[0, 1]$  and the segment  $(1, 2)$ .



## 2.47 Theorem

A subset  $E$  of the real line  $\mathbb{R}^1$  is connected if and only if it has the following property:

If  $x \in E, y \in E$ , and  $x < z < y$ , then  $z \in E$ .