

MAT426: Advanced Calculus - Numerical Sequences and Series

Convergent Sequences

Miraj Samarakkody

Tougaloo College

Updated: April 6, 2025

Convergent Sequences

3.1 Definition

A sequence $\{p_n\}$ in a metric space X is said to **converge** if there is a point $p \in X$ with the following property:

For every $\epsilon > 0$ there is an integer N such that $n \geq N$ implies that $d(p_n, p) < \epsilon$.

Convergent Sequences

3.1 Definition

A sequence $\{p_n\}$ in a metric space X is said to **converge** if there is a point $p \in X$ with the following property:

For every $\epsilon > 0$ there is an integer N such that $n \geq N$ implies that $d(p_n, p) < \epsilon$.

► We say that $\{p_n\}$ converges to p , or that p is the limit of p_n .

Convergent Sequences

3.1 Definition

A sequence $\{p_n\}$ in a metric space X is said to **converge** if there is a point $p \in X$ with the following property:

For every $\epsilon > 0$ there is an integer N such that $n \geq N$ implies that $d(p_n, p) < \epsilon$.

- ▶ We say that $\{p_n\}$ converges to p , or that p is the limit of p_n .
- ▶ We write $p_n \rightarrow p$ or $\lim_{n \rightarrow \infty} p_n = p$.

Convergent Sequences

3.1 Definition

A sequence $\{p_n\}$ in a metric space X is said to **converge** if there is a point $p \in X$ with the following property:

For every $\epsilon > 0$ there is an integer N such that $n \geq N$ implies that $d(p_n, p) < \epsilon$.

- ▶ We say that $\{p_n\}$ converges to p , or that p is the limit of p_n .
- ▶ We write $p_n \rightarrow p$ or $\lim_{n \rightarrow \infty} p_n = p$.
- ▶ If $\{p_n\}$ does not converge, it is said to *diverge*.

Convergent Sequences

3.1 Definition

A sequence $\{p_n\}$ in a metric space X is said to **converge** if there is a point $p \in X$ with the following property:

For every $\epsilon > 0$ there is an integer N such that $n \geq N$ implies that $d(p_n, p) < \epsilon$.

- ▶ We say that $\{p_n\}$ converges to p , or that p is the limit of p_n .
- ▶ We write $p_n \rightarrow p$ or $\lim_{n \rightarrow \infty} p_n = p$.
- ▶ If $\{p_n\}$ does not converge, it is said to *diverge*.
- ▶ Convergent sequence depends not only on $\{p_n\}$ but also on X
Ex: Sequence $1/n$ in \mathbb{R}^1 and in set of all positive real numbers with $d(x, y) = |x - y|$

Convergent Sequences

- ▶ The set of all points p_n ($n = 1, 2, 3, \dots$) is the range of $\{p_n\}$.
- ▶ The range of a sequence may be finite set, or may be infinite.
- ▶ The sequence $\{p_n\}$ is said to be bounded if its range is bounded.

Example

Consider the following sequences of complex numbers ($X = \mathbb{R}^2$):

Example

Consider the following sequences of complex numbers ($X = \mathbb{R}^2$):

- ▶ If $s_n = 1/n$, then $\lim_{n \rightarrow \infty} s_n = 0$; the range is infinite and the sequence is bounded.

Example

Consider the following sequences of complex numbers ($X = \mathbb{R}^2$):

- ▶ If $s_n = 1/n$, then $\lim_{n \rightarrow \infty} s_n = 0$; the range is infinite and the sequence is bounded.
- ▶ If $s_n = n^2$, the sequence $\{s_n\}$ is unbounded, is divergent, and has infinite range.

Example

Consider the following sequences of complex numbers ($X = \mathbb{R}^2$):

- ▶ If $s_n = 1/n$, then $\lim_{n \rightarrow \infty} s_n = 0$; the range is infinite and the sequence is bounded.
- ▶ If $s_n = n^2$, the sequence $\{s_n\}$ is unbounded, is divergent, and has infinite range.
- ▶ If $s_n = 1 + [(-1)^n/n]$, the sequence $\{s_n\}$ converges to 1, is bounded, and has infinite range.

Example

Consider the following sequences of complex numbers ($X = \mathbb{R}^2$):

- ▶ If $s_n = 1/n$, then $\lim_{n \rightarrow \infty} s_n = 0$; the range is infinite and the sequence is bounded.
- ▶ If $s_n = n^2$, the sequence $\{s_n\}$ is unbounded, is divergent, and has infinite range.
- ▶ If $s_n = 1 + [(-1)^n/n]$, the sequence $\{s_n\}$ converges to 1, is bounded, and has infinite range.
- ▶ If $s_n = i^2$, the sequence $\{s_n\}$ is divergent, is bounded, and has finite range.

Example

Consider the following sequences of complex numbers ($X = \mathbb{R}^2$):

- ▶ If $s_n = 1/n$, then $\lim_{n \rightarrow \infty} s_n = 0$; the range is infinite and the sequence is bounded.
- ▶ If $s_n = n^2$, the sequence $\{s_n\}$ is unbounded, is divergent, and has infinite range.
- ▶ If $s_n = 1 + [(-1)^n/n]$, the sequence $\{s_n\}$ converges to 1, is bounded, and has infinite range.
- ▶ If $s_n = i^2$, the sequence $\{s_n\}$ is divergent, is bounded, and has finite range.
- ▶ If $s_n = 1$ ($n = 1, 2, 3, \dots$), then $\{s_n\}$ converges to 1, is bounded, and has finite range.

Example

Consider the following sequences of complex numbers ($X = \mathbb{R}^2$):

- ▶ If $s_n = 1/n$, then $\lim_{n \rightarrow \infty} s_n = 0$; the range is infinite and the sequence is bounded.
- ▶ If $s_n = n^2$, the sequence $\{s_n\}$ is unbounded, is divergent, and has infinite range.
- ▶ If $s_n = 1 + [(-1)^n/n]$, the sequence $\{s_n\}$ converges to 1, is bounded, and has infinite range.
- ▶ If $s_n = i^2$, the sequence $\{s_n\}$ is divergent, is bounded, and has finite range.
- ▶ If $s_n = 1$ ($n = 1, 2, 3, \dots$), then $\{s_n\}$ converges to 1, is bounded, and has finite range.
- ▶ If $s_n = 1$ ($n = 1, 2, 3, \dots$), then $\{s_n\}$ converges 1, is bounded, and has finite range.

Theorem

Let $\{p_n\}$ be a sequence in a metric space X .

- ▶ $\{p_n\}$ converges to $p \in X$ if and only if every neighborhood of p contains p_n for all but finitely many n .

Theorem

Let $\{p_n\}$ be a sequence in a metric space X .

- ▶ $\{p_n\}$ converges to $p \in X$ if and only if every neighborhood of p contains p_n for all but finitely many n .
- ▶ If $p \in X$, $p' \in X$, and if $\{p_n\}$ converges to p and to p' , then $p' = p$.

Theorem

Let $\{p_n\}$ be a sequence in a metric space X .

- ▶ $\{p_n\}$ converges to $p \in X$ if and only if every neighborhood of p contains p_n for all but finitely many n .
- ▶ If $p \in X$, $p' \in X$, and if $\{p_n\}$ converges to p and to p' , then $p' = p$.
- ▶ If $\{p_n\}$ converges, then $\{p_n\}$ is bounded.

Theorem

Let $\{p_n\}$ be a sequence in a metric space X .

- ▶ $\{p_n\}$ converges to $p \in X$ if and only if every neighborhood of p contains p_n for all but finitely many n .
- ▶ If $p \in X$, $p' \in X$, and if $\{p_n\}$ converges to p and to p' , then $p' = p$.
- ▶ If $\{p_n\}$ converges, then $\{p_n\}$ is bounded.
- ▶ If $E \subset X$ and if p is a limit point of E , then there is a sequence $\{p_n\}$ in E such that $p = \lim_{n \rightarrow \infty} p_n$.