

MAT426: Advanced Calculus

Miraj Samarakkody

Tougaloo College

04/01/2025

2.41 Theorem

Theorem

If a set E in \mathbb{R}^k has one of the following three properties, then it has the other two:

- (a) E is closed and bounded.
- (b) E is compact.
- (c) Every infinite subset of E has a limit point in E .

Proof:

2.42 Theorem

Theorem - Weierstrass

Every bounded infinite subset of \mathbb{R}^k has a limit point in \mathbb{R}^k .

Perfect Sets

2.43 Theorem

Let P be a non-empty perfect set in \mathbb{R}^k . Then P is uncountable.

Corollary

Corollary

Every interval $[a, b]$ is uncountable. In particular, the set of all real numbers is uncountable.

The Cantor Set

Cantor set is a perfect set in \mathbb{R}^1 which contain no segment.

The Cantor Set

Cantor set is a perfect set in \mathbb{R}^1 which contain no segment.

Let E_0 be the interval $[0, 1]$.

The Cantor Set

Cantor set is a perfect set in \mathbb{R}^1 which contain no segment.

Let E_0 be the interval $[0, 1]$.

Remove the segment $\left(\frac{1}{3}, \frac{2}{3}\right)$, and let E_1 be the union of the intervals

$$\left[0, \frac{1}{3}\right], \left[\frac{2}{3}, 1\right]$$

The Cantor Set

Cantor set is a perfect set in \mathbb{R}^1 which contain no segment.

Let E_0 be the interval $[0, 1]$.

Remove the segment $\left(\frac{1}{3}, \frac{2}{3}\right)$, and let E_1 be the union of the intervals

$$\left[0, \frac{1}{3}\right], \left[\frac{2}{3}, 1\right]$$

Remove the middle thirds of these intervals, and let E_2 be the union of the intervals

$$\left[0, \frac{1}{9}\right], \left[\frac{2}{9}, \frac{3}{9}\right], \left[\frac{6}{9}, \frac{7}{9}\right], \left[\frac{8}{9}, 1\right]$$

The Cantor set

Continuing in this way, we obtain a sequence of compact sets E_n , such that

1. $E_1 \supset E_2 \supset E_3 \supset \dots$
2. E_n is the union of 2^n intervals, each of length 3^{-n} .

The set

$$P = \bigcap_{n=1}^{\infty} E_n$$

is called the **Cantor set**.