

# MAT426: Advanced Calculus - Numerical Sequences and Series

## Convergent Sequences [1]

**Miraj Samarakkody**

Tougaloo College

Updated: April 17, 2025

# Convergent Sequences

## 3.1 Definition

A sequence  $\{p_n\}$  in a metric space  $X$  is said to **converge** if there is a point  $p \in X$  with the following property:

For every  $\epsilon > 0$  there is an integer  $N$  such that  $n \geq N$  implies that  $d(p_n, p) < \epsilon$ .

# Convergent Sequences

## 3.1 Definition

A sequence  $\{p_n\}$  in a metric space  $X$  is said to **converge** if there is a point  $p \in X$  with the following property:

For every  $\epsilon > 0$  there is an integer  $N$  such that  $n \geq N$  implies that  $d(p_n, p) < \epsilon$ .

► We say that  $\{p_n\}$  converges to  $p$ , or that  $p$  is the limit of  $p_n$ .

# Convergent Sequences

## 3.1 Definition

A sequence  $\{p_n\}$  in a metric space  $X$  is said to **converge** if there is a point  $p \in X$  with the following property:

For every  $\epsilon > 0$  there is an integer  $N$  such that  $n \geq N$  implies that  $d(p_n, p) < \epsilon$ .

- ▶ We say that  $\{p_n\}$  converges to  $p$ , or that  $p$  is the limit of  $p_n$ .
- ▶ We write  $p_n \rightarrow p$  or  $\lim_{n \rightarrow \infty} p_n = p$ .

# Convergent Sequences

## 3.1 Definition

A sequence  $\{p_n\}$  in a metric space  $X$  is said to **converge** if there is a point  $p \in X$  with the following property:

For every  $\epsilon > 0$  there is an integer  $N$  such that  $n \geq N$  implies that  $d(p_n, p) < \epsilon$ .

- ▶ We say that  $\{p_n\}$  converges to  $p$ , or that  $p$  is the limit of  $p_n$ .
- ▶ We write  $p_n \rightarrow p$  or  $\lim_{n \rightarrow \infty} p_n = p$ .
- ▶ If  $\{p_n\}$  does not converge, it is said to *diverge*.

# Convergent Sequences

## 3.1 Definition

A sequence  $\{p_n\}$  in a metric space  $X$  is said to **converge** if there is a point  $p \in X$  with the following property:

For every  $\epsilon > 0$  there is an integer  $N$  such that  $n \geq N$  implies that  $d(p_n, p) < \epsilon$ .

- ▶ We say that  $\{p_n\}$  converges to  $p$ , or that  $p$  is the limit of  $p_n$ .
- ▶ We write  $p_n \rightarrow p$  or  $\lim_{n \rightarrow \infty} p_n = p$ .
- ▶ If  $\{p_n\}$  does not converge, it is said to *diverge*.
- ▶ Convergent sequence depends not only on  $\{p_n\}$  but also on  $X$   
Ex: Sequence  $1/n$  in  $\mathbb{R}^1$  and in set of all positive real numbers with  $d(x, y) = |x - y|$

# Convergent Sequences

- ▶ The set of all points  $p_n$  ( $n = 1, 2, 3, \dots$ ) is the range of  $\{p_n\}$ .
- ▶ The range of a sequence may be finite set, or may be infinite.
- ▶ The sequence  $\{p_n\}$  is said to be bounded if its range is bounded.

## Example

Consider the following sequences of complex numbers ( $X = \mathbb{R}^2$ ):



## Example

Consider the following sequences of complex numbers ( $X = \mathbb{R}^2$ ):

- ▶ If  $s_n = 1/n$ , then  $\lim_{n \rightarrow \infty} s_n = 0$ ; the range is infinite and the sequence is bounded.

## Example

Consider the following sequences of complex numbers ( $X = \mathbb{R}^2$ ):

- ▶ If  $s_n = 1/n$ , then  $\lim_{n \rightarrow \infty} s_n = 0$ ; the range is infinite and the sequence is bounded.
- ▶ If  $s_n = n^2$ , the sequence  $\{s_n\}$  is unbounded, is divergent, and has infinite range.

## Example

Consider the following sequences of complex numbers ( $X = \mathbb{R}^2$ ):

- ▶ If  $s_n = 1/n$ , then  $\lim_{n \rightarrow \infty} s_n = 0$ ; the range is infinite and the sequence is bounded.
- ▶ If  $s_n = n^2$ , the sequence  $\{s_n\}$  is unbounded, is divergent, and has infinite range.
- ▶ If  $s_n = 1 + [(-1)^n/n]$ , the sequence  $\{s_n\}$  converges to 1, is bounded, and has infinite range.

## Example

Consider the following sequences of complex numbers ( $X = \mathbb{R}^2$ ):

- ▶ If  $s_n = 1/n$ , then  $\lim_{n \rightarrow \infty} s_n = 0$ ; the range is infinite and the sequence is bounded.
- ▶ If  $s_n = n^2$ , the sequence  $\{s_n\}$  is unbounded, is divergent, and has infinite range.
- ▶ If  $s_n = 1 + [(-1)^n/n]$ , the sequence  $\{s_n\}$  converges to 1, is bounded, and has infinite range.
- ▶ If  $s_n = i^2$ , the sequence  $\{s_n\}$  is divergent, is bounded, and has finite range.

## Example

Consider the following sequences of complex numbers ( $X = \mathbb{R}^2$ ):

- ▶ If  $s_n = 1/n$ , then  $\lim_{n \rightarrow \infty} s_n = 0$ ; the range is infinite and the sequence is bounded.
- ▶ If  $s_n = n^2$ , the sequence  $\{s_n\}$  is unbounded, is divergent, and has infinite range.
- ▶ If  $s_n = 1 + [(-1)^n/n]$ , the sequence  $\{s_n\}$  converges to 1, is bounded, and has infinite range.
- ▶ If  $s_n = i^2$ , the sequence  $\{s_n\}$  is divergent, is bounded, and has finite range.
- ▶ If  $s_n = 1$  ( $n = 1, 2, 3, \dots$ ), then  $\{s_n\}$  converges to 1, is bounded, and has finite range.

## Example

Consider the following sequences of complex numbers ( $X = \mathbb{R}^2$ ):

- ▶ If  $s_n = 1/n$ , then  $\lim_{n \rightarrow \infty} s_n = 0$ ; the range is infinite and the sequence is bounded.
- ▶ If  $s_n = n^2$ , the sequence  $\{s_n\}$  is unbounded, is divergent, and has infinite range.
- ▶ If  $s_n = 1 + [(-1)^n/n]$ , the sequence  $\{s_n\}$  converges to 1, is bounded, and has infinite range.
- ▶ If  $s_n = i^2$ , the sequence  $\{s_n\}$  is divergent, is bounded, and has finite range.
- ▶ If  $s_n = 1$  ( $n = 1, 2, 3, \dots$ ), then  $\{s_n\}$  converges to 1, is bounded, and has finite range.
- ▶ If  $s_n = 1$  ( $n = 1, 2, 3, \dots$ ), then  $\{s_n\}$  converges 1, is bounded, and has finite range.

# Theorem

Let  $\{p_n\}$  be a sequence in a metric space  $X$ .

- ▶  $\{p_n\}$  converges to  $p \in X$  if and only if every neighborhood of  $p$  contains  $p_n$  for all but finitely many  $n$ .

# Theorem

Let  $\{p_n\}$  be a sequence in a metric space  $X$ .

- ▶  $\{p_n\}$  converges to  $p \in X$  if and only if every neighborhood of  $p$  contains  $p_n$  for all but finitely many  $n$ .
- ▶ If  $p \in X$ ,  $p' \in X$ , and if  $\{p_n\}$  converges to  $p$  and to  $p'$ , then  $p' = p$ .



# Theorem

Let  $\{p_n\}$  be a sequence in a metric space  $X$ .

- ▶  $\{p_n\}$  converges to  $p \in X$  if and only if every neighborhood of  $p$  contains  $p_n$  for all but finitely many  $n$ .
- ▶ If  $p \in X$ ,  $p' \in X$ , and if  $\{p_n\}$  converges to  $p$  and to  $p'$ , then  $p' = p$ .
- ▶ If  $\{p_n\}$  converges, then  $\{p_n\}$  is bounded.

# Theorem

Let  $\{p_n\}$  be a sequence in a metric space  $X$ .

- ▶  $\{p_n\}$  converges to  $p \in X$  if and only if every neighborhood of  $p$  contains  $p_n$  for all but finitely many  $n$ .
- ▶ If  $p \in X$ ,  $p' \in X$ , and if  $\{p_n\}$  converges to  $p$  and to  $p'$ , then  $p' = p$ .
- ▶ If  $\{p_n\}$  converges, then  $\{p_n\}$  is bounded.
- ▶ If  $E \subset X$  and if  $p$  is a limit point of  $E$ , then there is a sequence  $\{p_n\}$  in  $E$  such that  $p = \lim_{n \rightarrow \infty} p_n$ .

### 3.3 Theorem

Suppose  $\{s_n\}, \{t_n\}$  are complex sequences, and  $\lim_{n \rightarrow \infty} s_n = s$ ,  $\lim_{n \rightarrow \infty} t_n = t$ . Then:

### 3.3 Theorem

Suppose  $\{s_n\}, \{t_n\}$  are complex sequences, and  $\lim_{n \rightarrow \infty} s_n = s$ ,  $\lim_{n \rightarrow \infty} t_n = t$ . Then:

►  $\lim_{n \rightarrow \infty} (s_n + t_n) = s + t$

### 3.3 Theorem

Suppose  $\{s_n\}, \{t_n\}$  are complex sequences, and  $\lim_{n \rightarrow \infty} s_n = s$ ,  $\lim_{n \rightarrow \infty} t_n = t$ . Then:

- ▶  $\lim_{n \rightarrow \infty} (s_n + t_n) = s + t$
- ▶  $\lim_{n \rightarrow \infty} cs_n = cs$  for any number  $c$ .

### 3.3 Theorem

Suppose  $\{s_n\}, \{t_n\}$  are complex sequences, and  $\lim_{n \rightarrow \infty} s_n = s$ ,  $\lim_{n \rightarrow \infty} t_n = t$ . Then:

- ▶  $\lim_{n \rightarrow \infty} (s_n + t_n) = s + t$
- ▶  $\lim_{n \rightarrow \infty} cs_n = cs$  for any number  $c$ .
- ▶  $\lim_{n \rightarrow \infty} (c \pm s_n) = c \pm s$  for any number  $c$ .

### 3.3 Theorem

Suppose  $\{s_n\}, \{t_n\}$  are complex sequences, and  $\lim_{n \rightarrow \infty} s_n = s$ ,  $\lim_{n \rightarrow \infty} t_n = t$ . Then:

- ▶  $\lim_{n \rightarrow \infty} (s_n + t_n) = s + t$
- ▶  $\lim_{n \rightarrow \infty} cs_n = cs$  for any number  $c$ .
- ▶  $\lim_{n \rightarrow \infty} (c \pm s_n) = c \pm s$  for any number  $c$ .
- ▶  $\lim_{n \rightarrow \infty} (s_n t_n) = st$

### 3.3 Theorem

Suppose  $\{s_n\}, \{t_n\}$  are complex sequences, and  $\lim_{n \rightarrow \infty} s_n = s$ ,  $\lim_{n \rightarrow \infty} t_n = t$ . Then:

- ▶  $\lim_{n \rightarrow \infty} (s_n + t_n) = s + t$
- ▶  $\lim_{n \rightarrow \infty} cs_n = cs$  for any number  $c$ .
- ▶  $\lim_{n \rightarrow \infty} (c \pm s_n) = c \pm s$  for any number  $c$ .
- ▶  $\lim_{n \rightarrow \infty} (s_n t_n) = st$
- ▶  $\lim_{n \rightarrow \infty} (1/t_n) = 1/t$  provided  $t_n \neq 0$  and  $t \neq 0$ .



## 3.4 Theorem

- Suppose  $\mathbf{x}_n \in \mathbb{R}^k$  ( $n = 1, 2, 3, \dots$ ) and

$$\mathbf{x}_n = (\alpha_{1,n}, \dots, \alpha_{k,n}).$$

Then  $\{\mathbf{x}_n\}$  converges to  $\mathbf{x} = (\alpha_1, \dots, \alpha_k)$  if and only if  $\lim_{n \rightarrow \infty} \alpha_{i,n} = \alpha_i$  for  $i = 1, 2, \dots, k$ .

# Subsequences

# Definition

## 3.5 Definitions

Given a sequence  $\{p_n\}$ , consider a sequence  $n_k$  of positive integers, such that  $n_1 < n_2 < n_3 \dots$ . Then the sequence  $\{p_{n_l}\}$  is called a subsequence of  $\{p_n\}$ . If  $\{p_{n_l}\}$  converges, its limit is called a subsequential limit of  $\{p_n\}$ .

# Definition

## 3.5 Definitions

Given a sequence  $\{p_n\}$ , consider a sequence  $n_k$  of positive integers, such that  $n_1 < n_2 < n_3 \dots$ . Then the sequence  $\{p_{n_l}\}$  is called a subsequence of  $\{p_n\}$ . If  $\{p_{n_l}\}$  converges, its limit is called a subsequential limit of  $\{p_n\}$ .

It is clear that  $\{p_n\}$  converges to  $p$  if and only if every subsequence of  $\{p_n\}$  converges to  $p$ .

# Theorem

## 3.6 Theorem

- ▶ If  $\{p_n\}$  is a sequence in a metric space  $X$ , then some subsequence of  $p_n$  converges to a point of  $X$ .

# Theorem

## 3.6 Theorem

- ▶ If  $\{p_n\}$  is a sequence in a metric space  $X$ , then some subsequence of  $p_n$  converges to a point of  $X$ .
- ▶ Every bounded sequence in  $\mathbb{R}^k$  contains a convergent subsequences.

# Theorem

## 3.7 Theorem

The subsequential limit of a sequence  $\{p_n\}$  in a metric space  $X$  form a closed subset of  $X$ .

# References



Walter Rudin.

*Principles of Mathematical Analysis.*

McGraw-Hill, New York, 3rd edition, 1976.