

MAT434: Theory of Mathematical Statistics

Miraj Samarakkody

Tougaloo College

02/12/2025

Refer the operation table.

Refer the operation table.
What is closure property?

Refer the operation table.
What is closure property?
Identity

Refer the operation table.

What is closure property?

Identity

Inverse

Refer the operation table.

What is closure property?

Identity

Inverse

Commutative or Abelian

The Dihedral Group

The dihedral group of order $2n$ is often called the group of symmetries of a regular n -gon.

Groups

Let G be a set. A binary operation on G is a function that assigns each ordered pair of elements of G an element of G .

Groups

Let G be a set. A binary operation on G is a function that assigns each ordered pair of elements of G an element of G .

Definition

Let G be a set together with a binary operation that assigns to each ordered pair (a, b) of elements of G an element in G denoted by ab . The G is a group if the following three properties are satisfied.

Groups

Let G be a set. A binary operation on G is a function that assigns each ordered pair of elements of G an element of G .

Definition

Let G be a set together with a binary operation that assigns to each ordered pair (a, b) of elements of G an element in G denoted by ab . The G is a group if the following three properties are satisfied.

1. **Associativity:** $(ab)c = a(bc)$ for all $a, b \in G$.
2. **Identity:** There is an element e (called identity) in G such that $ae = ea = a$ for all $a \in G$.
3. **Inverses:** For each element $a \in G$, there is an element $b \in G$ such that $ab = ba = e$.

If for any $a, b \in G$, $ab = ba$ then group is an abelian group.

Examples

1. The set of integers \mathbb{Z} , the set of rational numbers \mathbb{Q} , and the set of real numbers \mathbb{R} , are all groups under ordinary addition.

Examples

1. The set of integers \mathbb{Z} , the set of rational numbers \mathbb{Q} , and the set of real numbers \mathbb{R} , are all groups under ordinary addition.
2. The subset $\{1, -1, i, -i\}$ of the complex numbers under multiplication is a group.

Examples

1. The set of integers \mathbb{Z} , the set of rational numbers \mathbb{Q} , and the set of real numbers \mathbb{R} , are all groups under ordinary addition.
2. The subset $\{1, -1, i, -i\}$ of the complex numbers under multiplication is a group.
3. The set S of positive irrational numbers together with 1 under multiplication is now a group.

Examples

1. The set of integers \mathbb{Z} , the set of rational numbers \mathbb{Q} , and the set of real numbers \mathbb{R} , are all groups under ordinary addition.
2. The subset $\{1, -1, i, -i\}$ of the complex numbers under multiplication is a group.
3. The set S of positive irrational numbers together with 1 under multiplication is now a group.
4. A rectangular array of the form $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is called a 2×2 matrix. The set of all 2×2 matrices with real entries is a group under componentwise addition.

$$\begin{bmatrix} a_1 & b_1 \\ c_1 & d_1 \end{bmatrix} + \begin{bmatrix} a_2 & b_2 \\ c_2 & d_2 \end{bmatrix} = \begin{bmatrix} a_1 + a_2 & b_1 + b_2 \\ c_1 + c_2 & d_1 + d_2 \end{bmatrix}$$

Examples Cont.

5. The set $\mathbb{Z}_n = \{0, 1, \dots, n-1\}$ for $n \geq 1$ is a group under addition modulo n . This group is usually referred to as the group of integers modulo n .

Examples Cont.

5. The set $\mathbb{Z}_n = \{0, 1, \dots, n-1\}$ for $n \geq 1$ is a group under addition modulo n . This group is usually referred to as the group of integers modulo n .
6. The set \mathbb{R}^* of non-zero real numbers is a group under ordinary multiplication. The identity is 1 and the inverse of a is $1/a$.

Examples Cont.

5. The set $\mathbb{Z}_n = \{0, 1, \dots, n-1\}$ for $n \geq 1$ is a group under addition modulo n . This group is usually referred to as the group of integers modulo n .
6. The set \mathbb{R}^* of non-zero real numbers is a group under ordinary multiplication. The identity is 1 and the inverse of a is $1/a$.
7. The set of all invertible $n \times n$ matrices with real entries is a group under matrix multiplication. This group is denoted by $GL(n, \mathbb{R})$.

$$GL(2, \mathbb{R}) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mid ad - bc \neq 0 \text{ for } a, b, c, d \in \mathbb{R} \right\}$$

Examples Cont.

5. The set $\mathbb{Z}_n = \{0, 1, \dots, n-1\}$ for $n \geq 1$ is a group under addition modulo n . This group is usually referred to as the group of integers modulo n .
6. The set \mathbb{R}^* of non-zero real numbers is a group under ordinary multiplication. The identity is 1 and the inverse of a is $1/a$.
7. The set of all invertible $n \times n$ matrices with real entries is a group under matrix multiplication. This group is denoted by $GL(n, \mathbb{R})$.

$$GL(2, \mathbb{R}) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mid ad - bc \neq 0 \text{ for } a, b, c, d \in \mathbb{R} \right\}$$

What happens when $ad - bc = 0$?