

# MAT434: Theory of Mathematical Statistics

**Miraj Samarakkody**

Tougaloo College

03/28/2025

## 2.3 Functions of a Random Variable

# Proposition C

## Proposition

Let  $Z = F(X)$ ; then  $Z$  has a uniform distribution of  $[0, 1]$ .

Proof:

# Proposition D

## Proposition

Let  $U$  be uniform on  $[0, 1]$ , and let  $X = F^{-1}(U)$ . Then the cdf of  $X$  is  $F$ .

Proof:

# Joint Distributions

## 3.1 Introduction

### Definition

Two or more random variables defined on the same sample space are said to have a joint probability structure.

## 3.1 Introduction

### Definition

Two or more random variables defined on the same sample space are said to have a joint probability structure.

### Applications

- ▶ In ecological studies. For an example when we count several species, one species is often the prey of another. i.e. the number of predators will be related to the number of prey.

## 3.1 Introduction

### Definition

Two or more random variables defined on the same sample space are said to have a joint probability structure.

### Applications

- ▶ In ecological studies. For an example when we count several species, one species is often the prey of another. i.e. the number of predators will be related to the number of prey.
- ▶ The joint probability distribution of the  $x$ ,  $y$ , and  $z$  components of wind velocity can be experimentally measured in studies of atmospheric turbulence.



## 3.1 Introduction

### Definition

Two or more random variables defined on the same sample space are said to have a joint probability structure.

### Applications

- ▶ In ecological studies. For an example when we count several species, one species is often the prey of another. i.e. the number of predators will be related to the number of prey.
- ▶ The joint probability distribution of the  $x$ ,  $y$ , and  $z$  components of wind velocity can be experimentally measured in studies of atmospheric turbulence.
- ▶ The joint distribution of the values of various physiological variables in a population of patients is often of interest in medical studies.

## 3.1 Introduction

### Definition

Two or more random variables defined on the same sample space are said to have a joint probability structure.

### Applications

- ▶ In ecological studies. For an example when we count several species, one species is often the prey of another. i.e. the number of predators will be related to the number of prey.
- ▶ The joint probability distribution of the  $x$ ,  $y$ , and  $z$  components of wind velocity can be experimentally measured in studies of atmospheric turbulence.
- ▶ The joint distribution of the values of various physiological variables in a population of patients is often of interest in medical studies.
- ▶ A model for the joint distribution of age and length in a population of fish can be used to estimate the age distribution from the length distribution.

# Introduction

The joint behavior of two random variables,  $X$  and  $Y$ , is determined by the cumulative distribution function

$$F(x, y) = P(X \leq x, Y \leq y)$$

regardless of whether  $X$  and  $Y$  are continuous or discrete.


# Introduction

The joint behavior of two random variables,  $X$  and  $Y$ , is determined by the cumulative distribution function

$$F(x, y) = P(X \leq x, Y \leq y)$$

regardless of whether  $X$  and  $Y$  are continuous or discrete.


The cdf gives the probability that the point  $(X, Y)$  belongs to a semi-infinite rectangle in the plane.



# Introduction

The probability that  $(X, Y)$  belongs to a rectangle if

$$P(x_1 \leq X \leq x_2, y_1 \leq Y \leq y_2) = F(x_2, y_2) - F(x_2, y_1) - F(x_1, y_2) + F(x_1, y_1)$$



# Discrete Random Variables

Suppose that  $X$  and  $Y$  are discrete random variables defined on the same sample space and that they take on values  $x_1, x_2, \dots$ , and  $y_1, y_2, \dots$  respectively.

# Discrete Random Variables

Suppose that  $X$  and  $Y$  are discrete random variables defined on the same sample space and that they take on values  $x_1, x_2, \dots$ , and  $y_1, y_2, \dots$  respectively.

Their **joint frequency function**  $p(x, y)$  is

$$p(x_i, y_j) = P(X = x_i, Y = y_j)$$

## Example

A fair coin is tossed three times and let  $X$  denote the number of heads on the first toss and  $Y$  the total number of heads. Find joint frequency function of  $X$  and  $Y$ .



## Example

A fair coin is tossed three times and let  $X$  denote the number of heads on the first toss and  $Y$  the total number of heads. Find joint frequency function of  $X$  and  $Y$ .

Find the frequency function of  $Y$  ( $p_Y$ ) from the joint frequency function.

## Example

A fair coin is tossed three times and let  $X$  denote the number of heads on the first toss and  $Y$  the total number of heads. Find joint frequency function of  $X$  and  $Y$ .

Find the frequency function of  $Y$  ( $p_Y$ ) from the joint frequency function.

The  $p_Y$  is called the **marginal frequency function** of  $Y$ .

# Marginal Frequency Function

Let  $X$  and  $Y$  are discrete random variables defined on the same sample space. The marginal function of  $X$  can be written as

$$p_X(x) = \sum_i p(x, y_i)$$

In the similar way, we can write the marginal frequency function for  $Y$ .