# MAT414 - Modern Algebra - Permutation Groups Cycle Notation [1]

### Miraj Samarakkody

Tougaloo College

Updated - April 13, 2025

# Cycle Notation

Write the followings in the cyclic notations:

$$\alpha = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 2 & 1 & 4 & 6 & 5 & 3 \end{bmatrix} \quad \beta = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 5 & 3 & 1 & 6 & 2 & 4 \end{bmatrix}$$

# Cycle Notation

Write the followings in the cyclic notations:

$$\alpha = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 2 & 1 & 4 & 6 & 5 & 3 \end{bmatrix} \quad \beta = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 5 & 3 & 1 & 6 & 2 & 4 \end{bmatrix}$$

Find  $\alpha\beta$ .

### Properties of Permutations

### Theorem 5.1 - Products of Disjoint Cycles

Every permutation of a finite set can be written as a cycle or as a product of disjoint cycles.

### Disjoint Cycles Commute

If the pair of cycles  $\alpha = (a_1, a_2, \dots, a_m)$  and  $\beta = (b_1, b_2, \dots, b_n)$  have no entries in common, then  $\alpha\beta = \beta\alpha$ .

#### Order of a Permutation

The order of a permutation of a finite set written in disjoint cycle form is the least common multiple of the lengths of the cycles.

#### Order of a Permutation

The order of a permutation of a finite set written in disjoint cycle form is the least common multiple of the lengths of the cycles.

#### Find

- **▶** |(132)(45)|
- **►** |(1432)(56)|
- ► |(123)(456)(78)|
- **▶** |(123)(145)|

Determine the orders of the elements of  $S_7$ .

Determine the number of elements in  $S_7$  of order 12.

Determine the number of elements in  $S_7$  of order 3.

Product of 2-Cycles

Every permutation of in  $S_n$ , n > 1, is a product of 2-cycles.

$$(1 \ 2 \ 3 \ 4 \ 5) =$$
 $(1 \ 6 \ 3 \ 2)(4 \ 5 \ 7) =$ 

#### Lemma

In  $S_n$ , if  $\epsilon=\beta_1\beta_2\beta_3\ldots\beta_r$ , where the  $\beta_i$ 's are 2-cycles, then r is even.

### Always Even or Always Odd

If a permutation  $\alpha$  can be expressed as a product of an even (odd) number of 2-cycles, then every decomposition of  $\alpha$  into a product of 2-cycles must have an an even (odd) number of 2-cycles.

In symbols, if

$$\alpha = \beta_1 \beta_2 \dots \beta_r$$
 and  $\alpha = \gamma_1 \gamma_2 \dots \gamma_s$ ,

where the  $\beta$ 's and  $\gamma$ 's are 2-cycles, then r and s are both even or both odd.

#### Even and Odd Permutations

#### Definition

A permutation that can be expressed as a product of an even number of 2—cycles is called an **even permutation**. A permutation that can be expressed as a product of an odd number of 2—cycles is called an **odd permutation**.

# Even Permutations Form a Group

#### Theorem 5.6

The set of all even permutations of  $S_n$  is a subgroup of  $S_n$  and is denoted by  $A_n$ .

# Alternating Group of Degree *n*

#### Definition


The alternating group of degree n, denoted  $A_n$ , is the set of all even permutations of  $S_n$ .

### **Theorem**

Theorem 5.7 For n > 1,  $A_n$  has order n!/2.

### Example 10 - Rotations of a Tetrahedron

The 12 rotations of a regular tetrahedron can be conveniently described with the elements of  $A_4$ .



### Example 10 - Rotations of a Tetrahedron

The 12 rotations of a regular tetrahedron can be conveniently described with the elements of  $A_4$ .


Table 5.1 The Alternating Group A, of Even Permutations of {1, 2, 3, 4}

(In this table, the permutations of  $A_4$  are designated as  $\alpha_1, \alpha_2, \ldots, \alpha_{12}$  and an entry k inside the table represents  $\alpha_k$ . For example,  $\alpha_3$   $\alpha_8 = \alpha_6$ .)

	$\alpha_{1}$	$\alpha_2$	$\alpha_3$	$\alpha_4$	$\alpha_{5}$	$\alpha_6$	$\alpha_7$	$\alpha_8$	$\alpha_9$	$\alpha_{_{10}}$	$\alpha_{11}$	$\alpha_{12}$
$(1) = \alpha_1$	1	2	3	4	5	6	7	8	9	10	11	12
$(12)(34) = \alpha_2$	2	1	4	3	6	5	8	7	10	9	12	11
$(13)(24) = \alpha_3$	3	4	1	2	7	8	5	6	11	12	9	10
$(14)(23) = \alpha_4$	4	3	2	1	8	7	6	5	12	11	10	9
$(123) = \alpha_5$	5	8	6	7	9	12	10	11	1	4	2	3
$(243) = \alpha_6$	6	7	5	8	10	11	9	12	2	3	1	4
$(142) = \alpha_7$	7	6	8	5	11	10	12	9	3	2	4	1
$(134) = \alpha_8$	8	5	7	6	12	9	11	10	4	1	3	2
$(132) = \alpha_9$	9	11	12	10	1	3	4	2	5	7	8	6
$(143) = \alpha_{10}$	10	12	11	9	2	4	3	1	6	8	7	5
$(234) = \alpha_{11}$	11	9	10	12	3	1	2	4	7	5	6	8
$(124) = \alpha_{12}$	12	10	9	11	4	2	1	3	8	6	5	7

### **Applications**

Many molecules with chemical formulas of the form  $AB_4$ , such as methane  $(CH_4)$  and carbon tetrachloride  $(CCI_4)$ , have  $A_4$  as thier symmetry group.



An intersting application of permutations is in the field of cryptography.

- An intersting application of permutations is in the field of cryptography.
- cryptography is the science of encoding and decoding messages.

- An intersting application of permutations is in the field of cryptography.
- cryptography is the science of encoding and decoding messages.
- ► The process of encoding a message is called encryption, and the process of decoding a message is called decryption.

- An intersting application of permutations is in the field of cryptography.
- cryptography is the science of encoding and decoding messages.
- ► The process of encoding a message is called encryption, and the process of decoding a message is called decryption.
- First known cryptosystme is th Caesar cipher.

- An intersting application of permutations is in the field of cryptography.
- cryptography is the science of encoding and decoding messages.
- ► The process of encoding a message is called encryption, and the process of decoding a message is called decryption.
- First known cryptosystme is th Caesar cipher.
- ▶ The Caesar cipher is a substitution cipher, which means that each letter in the plaintext is replaced by a letter some fixed number of positions down the alphabet.

► It was invented in 1974 by Hungarian architect and professor of architecture Ernő Rubik.

- ► It was invented in 1974 by Hungarian architect and professor of architecture Ernő Rubik.
- ► The cube has 6 faces, each with 9 stickers of one of 6 solid colors.

- It was invented in 1974 by Hungarian architect and professor of architecture Ernő Rubik.
- ➤ The cube has 6 faces, each with 9 stickers of one of 6 solid colors.
- ► The cube can be rotated about its axes, and the goal is to return the cube to its original state after it has been scrambled.

- It was invented in 1974 by Hungarian architect and professor of architecture Ernő Rubik.
- ► The cube has 6 faces, each with 9 stickers of one of 6 solid colors.
- ► The cube can be rotated about its axes, and the goal is to return the cube to its original state after it has been scrambled.
- ► The cube has 43, 252, 003, 274, 489, 856, 000 possible configurations.

- It was invented in 1974 by Hungarian architect and professor of architecture Ernő Rubik.
- ► The cube has 6 faces, each with 9 stickers of one of 6 solid colors.
- ► The cube can be rotated about its axes, and the goal is to return the cube to its original state after it has been scrambled.
- ► The cube has 43, 252, 003, 274, 489, 856, 000 possible configurations.
- ► God's number is 20, which means that any configuration of the cube can be solved in 20 moves or less.

			1 4	2 top	3 5						
			6	7	8						
9 12	10 left	11 13	17 20	18 front	19 21	25 28	26 right	27 29	33 36	34 rear	35 37
14	15	16	22	23	24	30	31	32	38	39	40
			41	42	43						
			44	bottom	45						
			46	47	48						

### References



Joseph A. Gallian.

Contemporary Abstract Algebra.

Cengage Learning, 9th edition, 2017.