

MAT414 - Modern Algebra

Miraj Samarakkody

Tougaloo College

03/24/2025

Cyclic Groups

Theorem 4.2 - $\langle a^k \rangle = \langle a^{\gcd(n,k)} \rangle$ and $|a^k| = n/\gcd(n, k)$

Let a be an element of order n in a group and let k be a positive integer. Then $\langle a^k \rangle = \langle a^{\gcd(n,k)} \rangle$ and $|a^k| = n/\gcd(n, k)$

Cyclic Groups

Theorem 4.2 - $\langle a^k \rangle = \langle a^{\gcd(n,k)} \rangle$ and $|a^k| = n/\gcd(n, k)$

Let a be an element of order n in a group and let k be a positive integer. Then $\langle a^k \rangle = \langle a^{\gcd(n,k)} \rangle$ and $|a^k| = n/\gcd(n, k)$

Proof steps:

We let $d = \gcd(n, k)$

- In the first part, we have to prove $\langle a^k \rangle \subseteq \langle a^{\gcd(n,k)} \rangle$ and $\langle a^k \rangle \supseteq \langle a^{\gcd(n,k)} \rangle$

Cyclic Groups

Theorem 4.2 - $\langle a^k \rangle = \langle a^{\gcd(n,k)} \rangle$ and $|a^k| = n/\gcd(n, k)$

Let a be an element of order n in a group and let k be a positive integer. Then $\langle a^k \rangle = \langle a^{\gcd(n,k)} \rangle$ and $|a^k| = n/\gcd(n, k)$

Proof steps:

We let $d = \gcd(n, k)$

- ▶ In the first part, we have to prove $\langle a^k \rangle \subset \langle a^{\gcd(n,k)} \rangle$ and $\langle a^{\gcd(n,k)} \rangle \subset \langle a^k \rangle$
- ▶ Let $d = \gcd(n, k)$

Cyclic Groups

Theorem 4.2 - $\langle a^k \rangle = \langle a^{\gcd(n,k)} \rangle$ and $|a^k| = n/\gcd(n, k)$

Let a be an element of order n in a group and let k be a positive integer. Then $\langle a^k \rangle = \langle a^{\gcd(n,k)} \rangle$ and $|a^k| = n/\gcd(n, k)$

Proof steps:

We let $d = \gcd(n, k)$

- ▶ In the first part, we have to prove $\langle a^k \rangle \subset \langle a^{\gcd(n,k)} \rangle$ and $\langle a^{\gcd(n,k)} \rangle \subset \langle a^k \rangle$
- ▶ Let $d = \gcd(n, k)$
- ▶ Write $d = ns + kt$ for some integers s, t

Cyclic Groups

Theorem 4.2 - $\langle a^k \rangle = \langle a^{\gcd(n,k)} \rangle$ and $|a^k| = n/\gcd(n, k)$

Let a be an element of order n in a group and let k be a positive integer. Then $\langle a^k \rangle = \langle a^{\gcd(n,k)} \rangle$ and $|a^k| = n/\gcd(n, k)$

Proof steps:

We let $d = \gcd(n, k)$

- ▶ In the first part, we have to prove $\langle a^k \rangle \subset \langle a^{\gcd(n,k)} \rangle$ and $\langle a^{\gcd(n,k)} \rangle \subset \langle a^k \rangle$
- ▶ Let $d = \gcd(n, k)$
- ▶ Write $d = ns + kt$ for some integers s, t
- ▶ Here we show that $|a^d| = n/d$ and then $|a^k| = n/d$.